

EXPONENTIAL GENUS PROBLEMS IN ONE-RELATOR PRODUCTS OF GROUPS

A.J. DUNCAN

ABSTRACT. Exponential equations in free groups were studied initially by Lyndon and Schützenberger [26] and then by Comerford and Edmunds [5]. Comerford and Edmunds showed that the problem of determining whether or not the class of quadratic exponential equations have solution is decidable, in finitely generated free groups. In this paper we show that for finite systems of quadratic exponential equations decidability passes, under certain hypotheses, from the factor groups to free products and one-relator products.

1. INTRODUCTION

In [26] Lyndon and Schützenberger studied the equation

$$a^M = b^N c^P,$$

where a, b and c are fixed elements of a free group and M, N and P are variables which may take integer variables. This is an example of an exponential equation. In fact it is shown in [26] that this equation has no solution with M, N and P all greater than 1, unless a, b and c are all powers of a common element. Comerford and Edmunds [5] studied more general exponential equations and showed that the problem of determining whether or not the class of quadratic exponential equations have solution is decidable, in finitely generated free groups. The definition of Comerford and Edmunds is that adopted here.¹ In fact Lyndon also introduced exponential groups (on which an associative ring acts by exponentiation) which he used to characterise the solution sets of equations in 1 variable in free groups [24], [25]. Lyndon considered the word problem in an exponential group and this can also be regarded as an exponential equation, but using a definition which allows more general values of exponents than that used here and in [5]. Exponential groups have been widely studied (see for example [1], [2], [12], [29], [21], [22], [20]). In particular these groups play an important role in work on Tarskii's conjectures on the elementary theory of free groups (see [21], [22], [34] and [33]). In this paper we consider exponential equations in the setting of one-relator products of groups. We show that for finite systems of quadratic exponential equations decidability is, under some extra hypotheses, inherited by free products and one-relator products.

Received by the editors February 1, 2008.

2000 *Mathematics Subject Classification*. Primary 20F65 20F05 20F10; Secondary 20F06 57M07.

Research partially supported by EPSRC grants B/91/RFH/1148 and GR/N28313.

¹In [5] the word “parametric” is used instead of “exponential”.

Let G be a group given by a recursive presentation $\langle C|R \rangle$, write $\bar{u}$ for the element of G represented by $u \in F(C)$ and let X be a recursive set. An equation over G with variables in X is an expression of the form

$$(1.1) \quad w = 1,$$

where w is an element of $F(C)$. A solution to (1.1) is a homomorphism $\phi : F(C \dot{\cup} X) \rightarrow G$ such that $\phi(w) = 1$ and $\phi(c) = \bar{c}$, for all $c \in C$. A class of equations over G is said to be solvable (or decidable) if there exists an algorithm which will determine whether or not a given equation of the class has a solution. As it is well known that the word problem is not solvable in certain finitely presented groups, in general equations are not solvable. However in suitably restricted classes of groups large classes of equations are known to be solvable. One of the most striking results in this direction is Makanin's: that finite systems of equations are solvable in finitely generated free groups [27]. Furthermore Rips and Sela [31] have generalised this result to torsion-free hyperbolic groups.

An equation is quadratic if every variable which occurs at all does so exactly twice. For example, if $u, v \in F(C)$, and $x, y, z \in X$, for then the equations $u = 1$, $ux^{-1}vxy^2 = 1$ and $ux^{-1}vx[y, z] = 1$ are all quadratic. These examples are all of the form

$$\prod_{i=1}^n x_i^{-1} u_i x_i \prod_{i=1}^t [x_{i+n}, x_{i+n+t}] \prod_{i=1}^p x_{i+n+2t}^2 = 1,$$

with $u_i \in F(C)$ and $x_i \in X$. In fact all quadratic equations are equivalent to at least one such equation: where equations with the same solution sets are deemed equivalent. The *genus* of an n -tuple $\bar{u}_1, \dots, \bar{u}_n$ of elements of G is the set of pairs of integers (t, p) such that the equation above has solution. The genus problem is that of determining whether or not a given pair (t, p) belongs to the genus of a given n -tuple of elements of G . It is known (see Example 5.7 and [7], [8] and [9]) that decidability of the genus problem is inherited from the factor groups by one-relator products, under suitable hypotheses on the factors and the relator.

In this paper we generalise some of the results of Example 5.7 mentioned above to quadratic exponential equations. If λ and μ are unknowns, $u, v \in F(C)$ and $x, y, z \in X$ then the expressions $u^\lambda = 1$, $u^\lambda x^{-1} v^\mu xy^2 = 1$ and $u^\lambda x^{-1} v^\mu x[y, z] = 1$ are all examples of quadratic exponential equations. A solution to one of these equations consists of a map $\alpha : \{\lambda, \mu\} \rightarrow \mathbb{Z}$, say $\alpha(\lambda) = l$, $\alpha(\mu) = m$, and a homomorphism $\phi : F(C) \rightarrow G$ such that ϕ is a solution to the quadratic equation $u^l = 1$, $u^l x^{-1} v^m xy^2 = 1$ or $u^l x^{-1} v^m x[y, z] = 1$, as appropriate. Usually some conditions are imposed on the values that λ and μ can take. For instance in the exponential equations above it is reasonable to insist that $\lambda, \mu \neq 0$. More generally solutions with $\lambda > l$ and $\mu > m$ may be of interest. We leave the formal definition to Section 3 but roughly speaking a quadratic exponential equation consists of an expression, similar to one of those above, together with a finite system of equations, congruences and inequalities on $\lambda, \mu, \dots$, which we call a parameter system. We restrict to linear parameter systems, since as shown by Matiyasevič, Hilbert's tenth problem is unsolvable for non-linear systems [28]. In [5] an algorithm is constructed which will determine whether or not a given quadratic exponential equation has a solution in the case where G is a finitely generated free group (see Example 5.1).

Let A and B be groups. A *one-relator product* of A and B is a group of the form $G = (A * B)/N(s)$, where $A * B$ is the free product, $s \in A * B$ is a cyclically reduced

word of length at least 2, and $N(s)$ is the normal closure of s in $A * B$. The exponential genus problem is a natural generalisation of the genus problem, described above, from quadratic to quadratic exponential equations. We show that under suitable conditions the decidability of the exponential genus problem is inherited by G from the factor groups A and B . In order to prove this result a stronger hypothesis on A and B has been imposed: namely that simultaneous systems of quadratic exponential equations are solvable in A and B , in the following sense. Let $Q_1, \dots, Q_m, Q_{m+1}, \dots, Q_{m+n}$ be quadratic exponential equations subject to a system of parameters $\mathcal{L}$. If Q_i is an exponential equation over A , for $i = 1, \dots, m$ and over B , for $i = m+1, \dots, m+n$ then we regard these equations as simultaneous. For example take $a \in A$ and $b \in B$, where Q_1 is $a^\lambda = 1$ and Q_2 is $b^\mu = 1$, subject to the parameter system consisting of $\lambda > 0$, $\mu > 0$ and $\lambda = 3\mu - 7$. If there is an algorithm to decide whether or not such systems of simultaneous equations have a simultaneous solution then we say that quadratic exponential equations are solvable over (A, B) . This gives rise to a corresponding simultaneous definition of exponential genus (Definition 3.4). We prove that if quadratic exponential equations are solvable over (A, B) then quadratic exponential equations defined over G are also solvable, as long as the relator is a sufficiently high power and not of a particularly pathogenic type (see Corollary 5.12). In fact we show that simultaneous systems of quadratic equations defined over one-relator groups of the required form are also solvable (see Corollary 5.10). This follows from the main result of the paper, Theorem 5.8, which is that if the exponential genus problem is solvable in a class of groups then it is also solvable in one-relator products, of the required form, of these groups. All these results hold if we replace one-relator products with free products. This means in particular that we obtain the result of [5], that quadratic exponential equations are solvable in finitely generated free groups (see Corollary 5.11).

The paper is structured as follows. In Sections 2 and 3 we describe exponential quadratic equations and exponential genus. In Section 4 we describe how, given an exponential quadratic equation we can form a finite set, called a resolution, of corresponding equations in a normal form, which we call special. The original equation has a solution if and only if some element of the resolution has a solution so this allows us to restrict attention to special equations. In Section 5 we describe in more detail the decision problems under consideration. In Section 3.1 we state the main result of the paper, Theorem 5.8, and show how it is a consequence of the more technical Theorem 5.9. The remainder of the paper is devoted to proving Theorem 5.9. We use pictures which are described here following, mainly, Howie [16], [17] and [18]. The use of pictures in group theory goes back to Rourke [32] (see also Fenn [11]) and was developed to apply to one-relator products by Short [35]. The theory has been further developed and applied to various aspects of geometric group theory in, for example, [3], [10], [19] and [30]. Here we extend use of pictures made in [7], [8] and [9]. We introduce pictures and establish some of their basic properties in Section 6. In particular we describe the link between solutions of equations and pictures. A picture can be thought of as a graph on a compact surface Σ with boundary. Each boundary component is assigned a label and these labels correspond to the group elements which occur in a quadratic equation. A solution to the quadratic equation arises from a picture and vice-versa. In Section 7 we define corridors which are later used to bound the degree of exponents occurring

in certain minimal solutions to exponential equations. In Sections 8 and 9 we assign angle and then curvature to vertices and regions of pictures. These are used together with corridors to bound the size of pictures that correspond to solutions to equations. To obtain this bound we analyse the curvature of regions of our pictures and show, in Sections 10, 11 and 12 that non-negative curvature occurs only at the boundary and in certain well defined configurations, see Figures 89 and 90 for example. This allows us, in Section 13, to bound the number of edges of a picture corresponding to a special equation, suitably minimised. We complete the proof of Theorem 5.9 in Section 14.

It seems likely that the results proved here should hold with at least some of the restrictions on the relator removed. In fact it would be surprising if the conditions on the factor groups and relators of Example 5.7 were not sufficient to prove Theorem 5.8. Indeed I believe that it is possible to push the proof given here to obtain results more like those of Example 5.7 without very much more effort. However, given the length of the proof, I do not feel that any increase in its complexity is desirable, especially for such minor gains. A more systematic approach to the arguments of Section 6 onwards might considerably shorten the argument and allow some unnecessary hypotheses to be removed.

2. QUADRATIC WORDS

Let X and D be sets indexed by $\mathbb{N}$. A word with *coefficients* in D and *variables* in X is a finite (possibly empty) sequence of elements of $(D \dot{\cup} X)^{\pm 1}$. Suppose $w = w_1 \cdots w_n$ is a word, where $w_i \in (D \dot{\cup} X)^{\pm 1}$, and let $a \in D \dot{\cup} X$. If $w_i = a^\varepsilon$, where $\varepsilon = \pm 1$, then w_i is called an *occurrence* of a in w , and a is said to *occur* in w .

Define

$$\begin{aligned} L_X(w) &= \{x \in X : x \text{ occurs in } w\}, \\ L_D(w) &= \{d \in D : d \text{ occurs in } w\} \text{ and} \\ L(w) &= L_D(w) \dot{\cup} L_X(w). \end{aligned}$$

Given $a \in D \dot{\cup} X$ define $O_a(w)$ to be the number of occurrences of a in w . The word w with coefficients in D and variables in X is *quadratic* if $O_x(w) = 2$, for all $x \in L_X(w)$, and $O_d(w) = 1$, for all $d \in L_D(w)$. A list of $k \geq 1$ words $\mathbf{w} = (w_1, \dots, w_k)$ with coefficients in D and variables in X is called a *system* of words, of *dimension* k , if $L(w_i) \cap L(w_j) = \emptyset$, whenever $i \neq j$. If $\mathbf{w} = (w_1, \dots, w_k)$ is a system of words then we define $L_D(\mathbf{w}) = \cup_{i=1}^k L_D(w_i)$, $L_X(\mathbf{w}) = \cup_{i=1}^k L_X(w_i)$ and $L(\mathbf{w}) = L_D(\mathbf{w}) \dot{\cup} L_X(\mathbf{w})$. An element $a \in L(\mathbf{w})$ is said to *occur* in $\mathbf{w}$. The *length* of $\mathbf{w}$ is $|\mathbf{w}| = \sum_{i=1}^k |w_i|$ and $\mathbf{w}$ is *quadratic* if w_i is quadratic, for $i = 1, \dots, k$.

A word w in an alphabet $A \dot{\cup} A^{-1}$ is *cyclically reduced* if no cyclic permutation of w has a subword of the form aa^{-1} or $a^{-1}a$, with $a \in A$. In [15] Grigorchuk and Lysionok take a cyclically reduced quadratic word $w = a_1^{\varepsilon_1} \cdots a_n^{\varepsilon_n}$, with coefficients in D and variables in X , where $a_i \in D \dot{\cup} X$ and $\varepsilon_i = \pm 1$, and describe a surface $\Sigma(w)$ associated to w . This is done as follows. Let I denote the unit interval $I = [0, 1]$ oriented from 0 to 1. Choose a partition $0 = p_0, \dots, p_n = 1$ of I into n sub-intervals of equal length. Label the sub-interval $[p_{i-1}, p_i]$ with a_i and orient it from p_{i-1} to p_i if $\varepsilon_i = 1$ and in the opposite sense otherwise. Identify p_0 and p_n to form a labelled oriented complex B homeomorphic to S^1 . Let Δ be a disk with boundary $\delta\Delta$ and identify $\delta\Delta$ with B . Then $\Sigma(w)$ is the quotient space of

Δ formed by identifying sub-intervals of B which have the same label, respecting the orientation of sub-intervals. The result is a compact surface $\Sigma(w)$, which has boundary if and only if $L_D(w)$ is non-empty. Furthermore $\Sigma(w)$ is orientable if and only if every element of $L_X(w)$ occurs once with positive exponent and once with negative exponent. We denote the number of boundary components of $\Sigma(w)$ by $\beta(w)$. Let $\Sigma'(w)$ be the surface obtained from $\Sigma(w)$ by capping off each boundary component with a disk. Define the *genus* of w , $\text{genus}(w) = \{(t, p) \in \mathbb{N}^2 : \Sigma(w) \text{ is homeomorphic to the connected sum of } t \text{ torii and } p \text{ projective planes}\}$.

If $\Sigma(w)$ has boundary component β then β is divided into oriented sub-intervals labelled by elements of D . Take an end point of one of these intervals as base point of β . If $\Sigma(w)$ is orientable we may choose an orientation on β such that, travelling round β in the direction of orientation every sub-interval $[p_{i-1}, p_i]$ that is encountered is traversed from p_{i-1} to p_i . Otherwise we fix one of the two possible orientations of β . Then traversing β once, in the direction of orientation, starting from the base point, and recording the labels as they occur, with positive exponent if the sub-interval orientation agrees with that of β and negative exponent otherwise, we obtain a word which we call a *boundary label* of β . If $\Sigma(w)$ has boundary components $\beta_1, \dots, \beta_n$ with boundary labels $b_1, \dots, b_n$, respectively, then $(b_1, \dots, b_n)$ is called a *boundary labels list* for w of *length* n , and $(w; b_1, \dots, b_n)$ is called a *labelled quadratic word*. Note that it is implicit in the definition of labelled quadratic word that w is cyclically reduced.

Let $F = F(D \dot{\cup} X)$ denote the free group on the set $D \dot{\cup} X$ and let $\text{Aut}_f F$ denote the subgroup of the automorphism group of F which consists of those elements which fix all but finitely many elements of $D \dot{\cup} X$. Given an automorphism $\theta \in \text{Aut}_f(F)$ let $\widetilde{u\theta}$ denote the reduced word representing $u\theta \in F$. Let $(w; b_1, \dots, b_n)$ be a labelled quadratic word. If $\widetilde{w\theta}$ is a quadratic word, $\Sigma(w)$ is homeomorphic to $\Sigma(\widetilde{w\theta})$ and $\widetilde{w\theta}$ has a boundary labels list $(\widetilde{b_1\theta}, \dots, \widetilde{b_n\theta})$, then θ is called an *admissible transformation* of $(w; b_1, \dots, b_n)$ to $(\widetilde{w\theta}; \widetilde{b_1\theta}, \dots, \widetilde{b_n\theta})$.

Given non-negative integers ξ, n, t, p define the quadratic word

$$q(\xi, n, t, p) = \prod_{i=1+\xi}^{n+\xi} x_i^{-1} d_i x_i \prod_{i=1+\xi}^{t+\xi} [x_{i+n}, x_{i+n+t}] \prod_{i=1+\xi}^{p+\xi} x_{i+n+2t}^2.$$

Let n be a non-negative integer. A *partition* of n of *length* k is a k -tuple $(n_1, \dots, n_k)$ of integers such that $n_i \geq 0$ and $\sum_{i=1}^k n_i = n$. Let $\mathbf{n} = (n_1, \dots, n_k)$, $\mathbf{t} = (t_1, \dots, t_k)$ and $\mathbf{p} = (p_1, \dots, p_k)$ be partitions of length $k > 0$, of integers n, t and p respectively, such that $n_j + t_j + p_j > 0$, for $1 \leq j \leq k$. Then we say that $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ is a *positive 3-partition* of (n, t, p) (of *length* k). If $\mathbf{h}$ is any partition, of length k of a non-negative integer h and $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ is a positive 3-partition then we say that $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ is a *positive 4-partition* of (h, n, t, p) . Given partitions $\mathbf{h}' = (h'_1, \dots, h'_l)$, $\mathbf{n}' = (n'_1, \dots, n'_l)$, $\mathbf{t}' = (t'_1, \dots, t'_l)$ and $\mathbf{p}' = (p'_1, \dots, p'_l)$ such that $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ is a positive 4-partition, we say that $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ if there exist integers $j_1, \dots, j_l$ such that $1 \leq j_1 < \dots < j_l \leq k$ with $h'_i \leq h_{j_i}$, $n'_i \leq n_{j_i}$, $t'_i \leq t_{j_i}$ and $p'_i \leq p_{j_i}$, for $i = 1, \dots, l$. If, in addition, $t'_i = t_{j_i}$ and $p'_i = p_{j_i}$, for $i = 1, \dots, l$, we say that $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ is a *sub-partition* of $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ and write $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq_g (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.

Given a positive 3-partition $(\mathbf{n}, \mathbf{t}, \mathbf{p})$, of length k as above, define

$$\begin{aligned} \xi_1 &= \xi_1(\mathbf{n}, \mathbf{t}, \mathbf{p}) = 0 \text{ and} \\ (2.1) \quad \xi_{j+1} &= \xi_{j+1}(\mathbf{n}, \mathbf{t}, \mathbf{p}) = \xi_j + n_j + 2t_j + p_j, \text{ for } j = 1, \dots, k-1. \end{aligned}$$

Next define the system of quadratic words

$$(2.2) \quad \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p}) = (q(\xi_j, n_j, t_j, p_j) : j = 1, \dots, k).$$

Note that $q(\xi_j, n_j, t_j, p_j)$ has boundary labels list $(d_{1+\xi_j}, \dots, d_{n_j+\xi_j})$. A system $\mathbf{w} = (w_1, \dots, w_l)$ of quadratic words is in *standard form* if $\mathbf{w} = \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p})$, for some positive 3-partition $(\mathbf{n}, \mathbf{t}, \mathbf{p})$.

Suppose that $\mathbf{w} = (w_1, \dots, w_k)$ is a system of quadratic words. We define the surface associated to $\mathbf{w}$ to be the compact surface

$$\Sigma(\mathbf{w}) = \bigcup_{i=1}^k \Sigma(w_i).$$

Assume that w_i has boundary labels list $\mathbf{b}^i = (b_1^i, \dots, b_{n_i}^i)$ of length n_i . We denote $((w_1; \mathbf{b}^1), \dots, (w_k; \mathbf{b}^k))$ by $(\mathbf{w}; \mathbf{b})$. We call $(\mathbf{w}; \mathbf{b})$ a *system of labelled quadratic words* which we say is in *standard form* if $\mathbf{w}$ is in standard form. Given $\theta \in \text{Aut}_f(F)$, let $\widetilde{\theta(\mathbf{b}^i)}$ denote $(\widetilde{\theta(b_1^i)}, \dots, \widetilde{\theta(b_{n_i}^i)})$ and $(\widetilde{\theta(\mathbf{w})}; \widetilde{\theta(\mathbf{b})})$ denote $((\widetilde{\theta(w_1)}; \widetilde{\theta(\mathbf{b}^1)}), \dots, (\widetilde{\theta(w_k)}; \widetilde{\theta(\mathbf{b}^k)}))$. Then θ is called an *admissible transformation* of $(\mathbf{w}; \mathbf{b})$ to $(\widetilde{\theta(\mathbf{w})}; \widetilde{\theta(\mathbf{b})})$ if θ is an admissible transformation of $(w_i; \mathbf{b}^i)$ to $(\widetilde{\theta(w_i)}; \widetilde{\theta(\mathbf{b}^i)})$, for $i = 1, \dots, k$, and $L(\theta(x)) \cap L(\theta(y)) = \emptyset$ whenever $x \in L(w_i)$, $y \in L(w_j)$ and $i \neq j$.

The following proposition can be proved exactly as the case $k = 1$ of [15, Proposition 2.1].

Proposition 2.1 (cf. [15, Proposition 2.1]). *Using the notation above, let $(\mathbf{w}; \mathbf{b})$ be a system of labelled quadratic words, of dimension k , let $n_i = \beta(w_i)$ and let $(t_i, p_i) \in \text{genus}(w_i)$, for $i = 1, \dots, k$. Let $\mathbf{n} = (n_1, \dots, n_k)$, $\mathbf{t} = (t_1, \dots, t_k)$ and $\mathbf{p} = (p_1, \dots, p_k)$. Then there exists an automorphism $\eta \in \text{Aut}_f(F(D \dot{\cup} X))$ such that η is an admissible transformation of $(\mathbf{w}; \mathbf{b})$ to $(\mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p}); \mathbf{d})$ and $\eta(F(D)) = F(D)$. Furthermore η can be effectively constructed.*

3. QUADRATIC EXPONENTIAL EQUATIONS

Let I be a recursive set and, for all $i \in I$, let $\langle C_i | R_i \rangle$ be a recursive presentation for a group H_i and let $H = *_{i \in I} H_i$. Let J be a recursively enumerable set and, for all $j \in J$, let $s_j \in H$, of free-product length at least 2, and let N be the normal closure of $\{s_j\}_{j \in J}$ in H .

Let $C = \dot{\cup}_{i \in I} C_i$, $R = \dot{\cup}_{i \in I} R_i$ and, for each $j \in J$, choose $\tilde{s}_j \in F(C)$ such that $\tilde{s}_j$ represents $s_j \in H$. Let $\tilde{\mathbf{s}} = \{\tilde{s}_j\}_{j \in J}$. Then H has a presentation $\langle C | R \rangle$ and $G = H/N$ has a presentation $\langle C | R \cup \tilde{\mathbf{s}} \rangle$, which we call the *natural* presentations of H and G , respectively. Given groups H_i with fixed presentations as above and groups H and G as described we shall always assume H and G are given by fixed natural presentations. By a *word* in H we mean an element of H written in free product normal form. The *length*, $l(h)$, of an element h of H refers to the free product length of a word in the normal form for H representing h (as opposed to the length as an element of $F(C)$). If u and v are words in H and u is a subword of some cyclic permutation of v then we say that u is a *cyclic subword* of v .

A *quadratic equation over G* is a pair of the form $(w = 1, \beta)$, where w is a quadratic word with coefficients in D and variables in X , and β is a map $\beta : X \rightarrow G$.

$L_D(w) \longrightarrow H$. A *solution* to the quadratic equation $(w = 1, \beta)$ consists of a homomorphism $\phi : F(L(w)) \longrightarrow H$ such that $\phi(w) \in N$ and $\phi(d) = \beta(d)$, for all $d \in L_D(w)$. In this section we define a generalisation of quadratic equations to what we shall call quadratic exponential equations.

We assume that H_i has solvable word problem, for all $i \in I$, and consequently H has solvable word problem: that is, that there exists an algorithm which, given $w \in F(C)$ will determine whether or not $w = 1$ in H and, if so, will output a product $\sum_{k=1}^n u_k^{-1} r_k^{\varepsilon_k} u_k = w$ in $F(C)$, where $u_k \in F(C)$, $r_k \in R$ and $\varepsilon_k = \pm 1$. Given this assumption we may formulate equations and decision problems for $\langle C | R \cup \tilde{s} \rangle$ in terms of words in the free product H instead of words in $F(C)$.

Let $\Lambda = \{\lambda_1, \dots\}$ be a set indexed by $\mathbb{N}$, which we call a set of *parameters*. Let N be the free $\mathbb{Z}$ -module on Λ and let M be the free $\mathbb{Z}$ -module $\mathbb{Z} \oplus N$. Elements of M are regarded as linear polynomials over Λ . An ordered pair (h, m) , $h \in H$, $m \in M$, $m \notin \mathbb{Z}$, h cyclically reduced is called a *proper exponential H -letter* and the set of all such ordered pairs is denoted $(H, \Lambda)_0$. An ordered pair (h, n) , $h \in H$, $l(h) = n \in \mathbb{Z}$ is called a *degenerate exponential H -letter* and the set of all such ordered pairs is denoted $(H, \Lambda)_1$. The set $(H, \Lambda)_0 \cup (H, \Lambda)_1$ is called the set of *exponential H -letters* and denoted H^Λ . We regard elements of $F(H^\Lambda)$ as reduced words on $H^\Lambda \cup (H^\Lambda)^{-1}$. Given $h \in H$ we identify h with $(h, l(h))$, so as a set $H \subset H^\Lambda$. An element $(h, f) \in H^\Lambda$ such that $h \in H_i$, for some $i \in I$, is called a *minor element* of H^Λ .

If $h \in H$ and $r \in \mathbb{Z}$ with $l(h) \geq r \geq 0$, define $\iota(h, r)$ to be the maximal initial subword of h of length r and $\tau(h, r)$ to be the maximal terminal subword of h of length r . Given $a \in H$, with $l(a) = m$, and $\alpha \in \mathbb{Z}$, let $q, r \in \mathbb{Z}$, such that $|\alpha| = qm + r$ and $0 \leq r < m$, and define

$$a \wedge \alpha = \begin{cases} a^q \iota(a, r) & \text{if } \alpha \geq 0 \\ a^{-q} \iota(a^{-1}, r) & \text{if } \alpha < 0 \end{cases}.$$

We have $a \wedge ql(a) = a^q$ so, if $l(a) > 1$ and $\alpha \neq 0$, then $a^\alpha \neq a \wedge \alpha$. Also if a is cyclically reduced

$$\begin{aligned} (a \wedge \alpha)^{-1} &= \begin{cases} (\iota(a, r)^{-1} \tau(a, m-r)^{-1}) \wedge \alpha & \text{if } \alpha \geq 0 \\ (\tau(a, r) \iota(a, m-r)) \wedge (-\alpha) & \text{if } \alpha < 0 \end{cases}, \\ &= \begin{cases} (\tau(a, m-r) \iota(a, r)) \wedge (-\alpha) & \text{if } \alpha \geq 0 \\ (\tau(a^{-1}, m-r) \iota(a^{-1}, r)) \wedge \alpha & \text{if } \alpha < 0 \end{cases}. \end{aligned}$$

Suppose a is cyclically reduced, $|\alpha| = ql(a) + r$, $0 \leq r < l(a)$, $a = a_0 a_1$, with $l(a_0) = r$, $l(a_1) = l(a) - r$, and $a = b_0 b_1$ with $l(b_1) = r$, $l(b_0) = l(a) - r$. Then

$$a \wedge \alpha = \begin{cases} a^q a_0 & \text{if } \alpha \geq 0 \\ a^{-q} b_1^{-1} & \text{if } \alpha < 0 \end{cases},$$

and

$$a^{-1} \wedge \alpha = \begin{cases} a^{-q} b_1^{-1} & \text{if } \alpha \geq 0 \\ a^q a_0 & \text{if } \alpha < 0 \end{cases},$$

so $a \wedge (-\alpha) = a^{-1} \wedge \alpha$.

A *parameter system* over Λ is a finite system of linear diophantine equations, congruences and inequalities in the variables Λ . A *solution* to the parameter system $\mathcal{L}$ is a retraction α from M to $\mathbb{Z}$ (i.e. a $\mathbb{Z}$ -module homomorphism from M onto $\mathbb{Z}$, fixing $\mathbb{Z}$ pointwise) such that substitution of $\alpha(\lambda)$ for λ , for each $\lambda \in \Lambda$, solves

$\mathcal{L}$. A parameter system is *consistent* if it has a solution. Let α be a $\mathbb{Z}$ -module retraction from M onto $\mathbb{Z}$. We define the *parameter evaluation homomorphism* $\hat{\alpha} : F(H^\Lambda) \longrightarrow H$ corresponding to α , by setting $\hat{\alpha}(h, f) = h \wedge \alpha(f)$, for $(h, f) \in H^\Lambda$.

Next we generalise the notation for presentations of H and G to collections of such groups and presentations. Let I , H_i , $\langle C_i | R_i \rangle$, $\langle C | R \rangle$ and H be defined as above. Let K be a recursive subset of $\mathbb{N}$ and let $X_k \subseteq I$ be a recursive set, for all $k \in K$. Define

$$(3.1) \quad H_{*,k} = *_{j \in X_k} H_j, \text{ for } k \in K.$$

Let $\mathbf{w} = (w_1, \dots, w_k)$ be a system of cyclically reduced quadratic words, with coefficients in D and variables in X , let $\beta : F(L_D(\mathbf{w})) \longrightarrow F(H^\Lambda)$ be a homomorphism and let $a_1, \dots, a_k$ be elements of K . If $\beta(d) \in F(H_{*,a_i}^\Lambda)$, for all $d \in L_D(w_i)$, $i = 1, \dots, k$, then the ordered pair $(\mathbf{w}, \beta)$ is called a *quadratic exponential word* over $((X_k)_{k \in K}; (H_i)_{i \in I})^\Lambda$ with *basis* $(a_1, \dots, a_k)$.

We define a *quadratic exponential equation* over $((X_k)_{k \in K}; (H_i)_{i \in I})^\Lambda$ with *basis* $(a_1, \dots, a_k)$ to be a pair $(\mathbf{w} = 1, \beta)$, where $(\mathbf{w}, \beta)$ is a quadratic exponential word over $((X_k)_{k \in K}; (H_i)_{i \in I})^\Lambda$ with basis $(a_1, \dots, a_k)$.

For each $k \in K$ let $\mathbf{s}_k$ be a recursively enumerable subset of $H_{*,k}$, such that each element $s \in \mathbf{s}_k$ satisfies $l(s) \geq 2$, and define

$$(3.2) \quad G_k = H_{*,k} / N_k,$$

where N_k is the normal closure of $\mathbf{s}_k$ in $H_{*,k}$. We allow the possibility that $\mathbf{s}_k$ is empty, so it may be that $G_k = H_{*,k}$. Given a quadratic exponential equation $W = (\mathbf{w} = 1, \beta)$ over $((X_k)_{k \in K}; (H_i)_{i \in I})^\Lambda$ with basis $(a_1, \dots, a_k)$ a *solution* to W in $(G_k)_{k \in K}$ is a pair (ϕ, α) such that α is a retraction from M to $\mathbb{Z}$, $\phi : F(L(\mathbf{w})) \longrightarrow H$ is a homomorphism, $\phi(a) \in H_{*,a_i}$, for all $a \in L(w_i)$, $\phi(w_i) \in N_{a_i}$, for $i = 1, \dots, k$, and $\phi(d) = \hat{\alpha}(\beta(d))$, for all $d \in L_D(\mathbf{w})$.

If W is an equation over $((X_k)_{k \in K}; (H_i)_{i \in I})^\Lambda$ for which solutions are sought in $(G_k)_{k \in K}$ then we say that W has *environment*

$$((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda.$$

To specify an equation fully its environment and basis must be given, but we omit these when the meaning is clear. If, in the definition of the $H_{*,k}$, the set K has a unique element s and $X_s = I$ then we refer to quadratic equations with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$ as equations over H^Λ with solutions in G .

Let $u \in F(H^\Lambda)$ and let $\mathcal{L}$ be a consistent parameter system. We say u is *constrained* by $\mathcal{L}$ if $\mathcal{L}$ implies that $f \equiv k \pmod{l(a)}$, for some $k \in \mathbb{Z}$, for all proper exponential H -letters $(a, f) \in H^\Lambda$ occurring in u , with $l(a) \geq 2$. A *constrained quadratic exponential equation* consists of a triple $(\mathbf{w} = 1, \beta, \mathcal{L})$, where $(\mathbf{w} = 1, \beta)$ is a quadratic exponential equation, $\mathcal{L}$ is a consistent parameter system and $\beta(d)$ is constrained by $\mathcal{L}$, for all $d \in L_D(\mathbf{w})$. A quadratic exponential equation $(\mathbf{w} = 1, \beta)$ or a constrained quadratic exponential equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ is in *standard form* if $\mathbf{w}$ is in standard form. A *solution* to the constrained quadratic exponential equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ is a solution (ϕ, α) to $(\mathbf{w} = 1, \beta)$ such that α is a solution to $\mathcal{L}$.

An element $(a, f) \in H^\Lambda$ is said to *occur* in the system of quadratic words $(\mathbf{w}, \beta)$ if (a, f) occurs in $\beta(d)$ for some $d \in L_D(\mathbf{w})$. A polynomial $f \in M$ is said to *occur* in the quadratic exponential equation $(\mathbf{w} = 1, \beta)$, or in the constrained quadratic exponential equation $(\mathbf{w} = 1, \beta, \mathcal{L})$, if $(a, f) \in H^\Lambda$ occurs in $(\mathbf{w}, \beta)$, for some $a \in H$. Also a polynomial f is said to *occur* in the parameter system $\mathcal{L}$ if $\mathcal{L}$ contains an

equation $f = 0$ or a congruence $f \equiv 0 \pmod{k}$, for some $k \in \mathbb{Z}$. A polynomial f is said to *occur* in the constrained quadratic exponential equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ if f occurs either in $(\mathbf{w} = 1, \beta)$ or in $\mathcal{L}$. An element $\lambda \in \Lambda$ is said to *occur* in $f \in M$ if the coefficient of λ in f is non-zero. Also λ is said to *occur* in a quadratic exponential equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ if λ occurs in an element $f \in M$ which occurs in $(\mathbf{w} = 1, \beta, \mathcal{L})$.

The H -length of a exponential H -letter (a, f) is $l_H(a, f) = l(a \wedge f)$. The H -length of a reduced word $u = (a_1, f_1)^{\varepsilon_1} \dots (a_n, f_n)^{\varepsilon_n} \in F(H^\Lambda)$, where $\varepsilon_i = \pm 1$, is defined to be

$$(3.3) \quad l_H(u) = \sum_{i=1}^n l_H(a_i, f_i).$$

The *exponential length* of u is defined to be

$$(3.4) \quad |u|_\Lambda = n - |\{i : 1 \leq i \leq n \text{ and } (a_i, f_i) \text{ is degenerate}\}|.$$

If $\alpha : M \rightarrow \mathbb{Z}$ is a retraction the *exponent length* of α on u is

$$(3.5) \quad |u|_\alpha = \sum_{i=1}^n |\alpha(f_i)|.$$

The H^Λ -length of u is defined to be

$$(3.6) \quad |u| = n.$$

Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation, with $\mathbf{w} = (w_1, \dots, w_k)$. Then the H -length, *exponential coordinates*, *exponential length* and H^Λ -length of W , are

$$\begin{aligned} l_H(W) &= \sum_{d \in L_D(\mathbf{w})} l_H(\beta(d)), \\ (W)_\Lambda &= \left(\sum_{d \in L_D(w_1)} |\beta(d)|_\Lambda, \dots, \sum_{d \in L_D(w_k)} |\beta(d)|_\Lambda \right), \\ |W|_\Lambda &= \sum_{d \in L_D(\mathbf{w})} |\beta(d)|_\Lambda \text{ and} \\ |W| &= \sum_{d \in L_D(\mathbf{w})} |\beta(d)|, \end{aligned}$$

respectively.

A solution (ϕ, α) to W is said to be *exponent-minimal* if, given any solution (ϕ', ρ) to W , the inequality

$$\sum_{d \in L_D(\mathbf{w})} |\beta(d)|_\rho \geq \sum_{d \in L_D(\mathbf{w})} |\beta(d)|_\alpha$$

holds.

Let $z \in *_{i \in I} H_i$, with $l(z) = n$, $z = u_1 \dots u_n$, where $u_j \in H_{i_j}$, for some $i_j \in I$ and $j = 1, \dots, n$. Define $\text{supp}(z)$ the *support* of z to be $\{i_1, \dots, i_n\}$, so $z \in *_{j \in \text{supp}(z)} H_j$. We define $\text{supp}(z, f)$ the *support* of $(z, f) \in H^\Lambda$ to be $\text{supp}(z)$. If $w \in F(H^\Lambda)$, $w = w_1^{\varepsilon_1} \dots w_n^{\varepsilon_n}$, with $w_j \in H^\Lambda$ and $\varepsilon_j = \pm 1$, we define $\text{supp}(w)$ the *support* of w to be

$\cup_{j=1}^n \text{supp}(w_j)$. Let $\mathbf{w} = (w_1, \dots, w_n)$ be a system of words. If $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ is a constrained quadratic exponential equation with environment

$$(3.7) \quad ((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$$

and basis $(a_1, \dots, a_k)$ then we define the *support* of W

$$\text{supp}(W) = (X'_{a_1}, \dots, X'_{a_k}),$$

where

$$X'_{a_i} = \bigcup \{ \text{supp}(\beta(d)) : d \in L_D(w_i) \} \cup \bigcup \{ \text{supp}(s) : s \in \mathbf{s}_{a_i} \}.$$

Define $I' = \cup_{i=1}^k X'_{a_i}$, $H'_{*,a_i} = *_{j \in X'_{a_i}} H_j$, N'_{a_i} to be the normal closure of $\mathbf{s}_{a_i} \in H'_{*,a_i}$ and $G'_{a_i} = H'_{*,a_i} / N'_{a_i}$, for $i = 1, \dots, k$. Then W is a constrained quadratic exponential equation with environment

$$(3.8) \quad ((X'_{a_j})_{j=1, \dots, k}; (H_i)_{i \in I'}; (G'_{a_j})_{j=1, \dots, k})^\Lambda$$

and basis $(a_1, \dots, a_k)$. Every solution to W with environment (3.8) is a solution to W with environment (3.7). The converse also holds as in the next lemma.

Lemma 3.1. *Every solution to W with environment (3.7) determines canonically a solution to W with environment (3.8).*

Proof. Let (ϕ, α) be a solution to W with environment (3.7). Fix $i \in \{1, \dots, k\}$, suppose $X'_{a_i} = \{i_1, \dots, i_s\}$ and set $X'' = X_{a_i} \setminus X'_{a_i}$. Let

$$A = *_{j=1}^s H_{i_j} \text{ and } B = *_{j \in X''} H_j.$$

Denote by f the canonical map $f : A * B \longrightarrow A$ such that $\ker(f)$ is the normal closure of B in $A * B$. Then N_{a_i} is by definition the normal closure of $\mathbf{s}_{a_i}$ in $A * B$ and $\mathbf{s}_{a_i} \subseteq A$, so f maps N_{a_i} into N'_{a_i} . Define $\phi' : L(w_i) \longrightarrow A$ by $\phi'(a) = f \circ \phi(a)$, for all $a \in L(w_i)$. If $d \in L_D(w_i)$ then $\phi(d) = \hat{\alpha}\beta(d)$ and $\text{supp}(\beta(d)) \subseteq X'_{a_i}$ so $\phi(d) \in A$. Hence $\phi'(d) = \hat{\alpha}\beta(d)$, for all $d \in L_D(w_i)$. Also $\phi'(w_i) \in N'_{a_i}$ as $\phi(w_i) \in N_{a_i}$. As $L(w_i) \cap L(w_j) = \emptyset$, when $i \neq j$, we can define ϕ' in this way simultaneously over $L(w_i)$, for $i = 1, \dots, k$. Then (ϕ', α) is a solution to W with environment (3.8).

Example 3.2. Let $I = \mathbb{N}$ and in each example below let $H_i = \langle C_i | R_i \rangle$ be trivial unless C_i and R_i are given explicitly.

- (i) Let $K = \{1, 2\}$, $X_1 = \{2\}$ and $X_2 = \{1, 2\}$. Let $C_i = \{c_{i1}, c_{i2}\}$, for $i = 1, 2$, $R_1 = \{c_{11}^3, c_{12}^3\}$, $R_2 = \{c_{21}^2, c_{22}^2\}$, $\mathbf{s}_1 = \emptyset$ and $\mathbf{s}_2 = \{c_{11}c_{22}c_{21}c_{12}c_{22}c_{21}\}$. Then $H_{*,1} = H_2$, $H_{*,2} = H_1 * H_2$, $G_1 = H_2$ and $G_2 = H_{*,2} / \langle \langle \mathbf{s}_2 \rangle \rangle$. Let $w_1 = d_1$, $w_2 = d_2x_3^{-1}d_3x_3$ and $\mathbf{w} = (w_1, w_2)$. Let $\beta(d_1) = (c_{22}, \lambda_1)$, $\beta(d_2) = (c_{12}c_{22}c_{21}, 2)(c_{11}c_{22}c_{21}, f_1)(c_{11}c_{22}, 2)$, and $\beta(d_3) = (c_{21}c_{22}c_{11}^2, f_2)(c_{22}c_{11}^2, 2)$, where $f_1 = \lambda_1 - 2$ and $f_2 = \lambda_1 - \lambda_2$. Let $\mathcal{L}_0$ consist of the congruences $\lambda_i \equiv 0 \pmod{2}$ and the inequalities $\lambda_i > 0$, for $i = 1, 2$. Let $\mathcal{L}_1$ consist of the same congruences and the inequalities $\lambda_1 > 4$, $\lambda_2 > 2$. Then $W_i = (\mathbf{w} = 1, \beta, \mathcal{L}_i)$ is a quadratic exponential equation with environment $(X_1, X_2; (H_i)_{i \in I}; G_1, G_2)^\Lambda$ and basis $(1, 2)$. These equations also have environment $(X_1, X_2; H_1, H_2; G_1, G_2)^\Lambda$. Let α_0 be a retraction of M to $\mathbb{Z}$ such that $\alpha_0(\lambda_1) = 4$ and $\alpha_0(\lambda_2) = 4$. Let $\phi_0(x_3) = c_{11}c_{22}$ and set $\phi_0(d_i) = \hat{\alpha}_0\beta(d_i)$, for $i = 1, 2, 3$. Then (ϕ_0, α_0) is a solution to W_0 but not W_1 . Let α_1 be such that $\alpha_1(\lambda_1) = 6$ and $\alpha_1(\lambda_2) = 4$. Let $\phi_1(x_3) = c_{11}c_{22}$ and set $\phi_1(d_i) = \hat{\alpha}_1\beta(d_i)$, for $i = 1, 2, 3$. Then (ϕ_1, α_1) is a solution to W_0 and W_1 .

- (ii) Let C_i, R_i, H_i be as in Example 3.2(i), for $i = 1, 2$ and let C_j, R_j be arbitrary for $j \geq 3$. Let $K = \mathbb{N}$ and $X_1 = \{1, 2\}$, $X_2 = \{1, 2, 3\}$ and $X_j = \{j, j+1\}$, for $j \geq 3$, so $H_{*,1} = H_1 * H_2$, $H_{*,2} = H_1 * H_2 * H_3$ and $H_{*,j} = H_j * H_{j+1}$, $j \geq 3$. Let $\mathbf{s}_i$ be as in Example 3.2(i), for $i = 1, 2$ and let $\mathbf{s}_j = \emptyset$, for $j \geq 3$. For $i = 0$ or 1 , the equation W_i of Example 3.2(i) is an equation with environment $((X_i)_{i \in K}; (H_i)_{i \in I}; (G_i)_{i \in K})^\Lambda$ and basis $(1, 2)$, with G_i defined as usual. The support of W_i is (X'_1, X'_2) , where X'_i is the set X_i of Example 3.2(i). Therefore W_i is an equation with environment $(X'_1, X'_2; H_1, H_2; G'_1, G'_2)^\Lambda$ and basis $(1, 2)$, for G'_i defined as usual. Moreover W_i has a solution with respect to the first named environment if and only if it has a solution with respect to the latter.

The next example illustrates equations corresponding to some common group theoretic questions.

Example 3.3. Let $I = \mathbb{N}$, and let C_i, R_i (and H_i) be given, for all $i \in I$. Let $K = \{1\}$, X_1 be a recursive subset of I and $\mathbf{s}_1 \subseteq H_{*,1}$ and G_1 be defined as usual. Let $\mathbf{w} = (w_1)$ and let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$, where w_1, β and $\mathcal{L}$ are described below. The equation W will in each case have environment $(X_1; (H_i)_{i \in I}; G_1)^\Lambda$ and basis (1).

- (i) Let $w_1 = d_1$ and $\beta(d_1) = (h_1, \lambda_1)$, where $h_1 \in H_{*,1}$. Assume $\mathcal{L}$ contains the subsystem $\mathcal{L}_0$ consisting of the inequality $\lambda_1 \neq 0$ and the congruence $\lambda_1 \equiv 0 \pmod{l(h_1)}$. As no elements of X occur in W , given a solution α to $\mathcal{L}$ we may determine whether there is a solution (ϕ, α) to W by setting $\phi = \hat{\alpha}\beta$ and evaluating $\phi(w_1)$. If there is a solution then we have $h_1^m = 1$ in G_1 , where $ml(h_1) = \alpha(\lambda_1)$. When $\mathcal{L} = \mathcal{L}_0$ finding a solution is equivalent to showing that h_1 has finite order.
- (ii) With w_1 and $\mathcal{L}$ as in the previous case let $\beta(d_1) = (h_1, \lambda_1)(h_2, l(h_2))$, where $h_1, h_2 \in H_{*,1}$. Again, given a solution α to $\mathcal{L}$ we may determine whether there is a solution (ϕ, α) to W by setting $\phi = \hat{\alpha}\beta$ and evaluating $\phi(w_1)$. If there is a solution then we have $h_1^m = h_2$ in G_1 , with m as before a non-zero integer.
- (iii) Let $w_1 = d_1 x_2^{-1} d_2 x_2$ and $\beta(d_i) = (h_i, \lambda_i)$, where $h_i \in H_{*,1}$, for $i = 1, 2$. Assume $\mathcal{L}$ contains the subsystem $\mathcal{L}_0$ consisting of the inequalities $\lambda_i \neq 0$ and the congruences $\lambda_i \equiv 0 \pmod{l(h_i)}$, for $i = 1, 2$. If there is a solution then h_1^m is conjugate to h_2^n in G_1 , for some non-zero integers m and n . When $\mathcal{L} = \mathcal{L}_0$ finding a solution is equivalent to showing that some power of h_1 is conjugate to a power of h_2 . If $\mathcal{L}$ is the union of $\mathcal{L}_0$ with the equation $\lambda_2 = l(h_2)$ then finding a solution is equivalent to showing that some non-trivial power of h_1 is conjugate to h_2 .

3.1. Quadratic equations and genus. Let I and K be recursive sets and for each $k \in K$ let X_k be a recursive subset of I . Let $H_{*,k}$ be the free-product defined in (3.1), for all $k \in K$. Let $\mathbf{s}_k$ be a recursively enumerable subset of $H_{*,k}$ and define G_k as in (3.2). Let $\mathcal{L}$ be a consistent parameter system. Let $\mathbf{z} = (z_1, \dots, z_n)$ be an n -tuple of elements of $F(H^\Lambda)$ such that z_i is constrained by $\mathcal{L}$ and $\text{supp}(z_i) \subseteq X_{a_i}$, for some $a_i \in K$, for $i = 1, \dots, n$. The set of all such $\mathbf{z}$ is denoted by

$$(\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}).$$

Let $\mathbf{n} = (n_1, \dots, n_k)$, $\mathbf{t} = (t_1, \dots, t_k)$ and $\mathbf{p} = (p_1, \dots, p_k)$ be k -tuples of integers such that $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ is a positive 3-partition of (n, t, p) , for some integers t and p .

Define $\xi_j = \xi_j(\mathbf{n}, \mathbf{t}, \mathbf{p})$, for $1 \leq j \leq k$, as in (2.1) and define

$$(3.9) \quad \begin{aligned} \mu_1 &= \mu_1(\mathbf{n}) = 0 \text{ and} \\ \mu_{j+1} &= \mu_{j+1}(\mathbf{n}) = \mu_j + n_j, \text{ for } j = 1, \dots, k. \end{aligned}$$

If $\mathbf{z} \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L})$ and there exist $a_1, \dots, a_k \in K$ such that $\text{supp}(z_i) \subseteq X_{a_j}$, for all i with $1 + \mu_j \leq i \leq \mu_{j+1}$ and for $j = 1, \dots, k$, then we say that $\mathbf{z}$ is *supported* over $\mathbf{n}$ with *basis* $(a_1, \dots, a_k)$.

Assume that $\mathbf{z}$ is supported over $\mathbf{n}$ with basis $(a_1, \dots, a_k)$. Define a map $\beta = \beta(\mathbf{z}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ from the subset $\cup_{j=1}^k \{d_{\xi_j+1}, \dots, d_{\xi_j+n_j}\}$ of D to $F(H^\Lambda)$ by

$$(3.10) \quad \beta(d_{\xi_j+i}) = z_{\mu_j+i}, \text{ for } j = 1, \dots, k,$$

and let $\mathbf{q} = \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p})$. The *quadratic exponential equation associated to* $(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ is defined to be the equation $(\mathbf{q} = 1, \beta, \mathcal{L})$ which we denote by $Q = Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Note that Q is an equation with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$ and basis $(a_1, \dots, a_k)$.

Definition 3.4. The $\mathcal{L}$ -genus of $\mathbf{z}$ in $(G_k : k \in K)$, denoted $\mathcal{L}\text{-genus}(\mathbf{z})$ is the set of positive 3-partitions $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ such that $\mathbf{z}$ is supported over $\mathbf{n}$ and Q has a solution.

Now let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition. We shall denote by

$$(3.11) \quad (\dot{\cup}_{k \geq 1} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$$

the set of n -tuples $\mathbf{z} \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L})$ such that $\mathbf{z}$ is supported over $\mathbf{n}$, and $\sum_{i=1}^{n_j} |z_{\mu_j+i}|_\Lambda \leq h_j$, for $j = 1, \dots, k$. (Note that given the latter condition the j th exponential coordinate h'_j of Q satisfies $h'_j \leq h_j$, for all j .) If the set K has a unique element, s say, and $X_s = I$ then we write

$$(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$$

instead of (3.11) and to simplify notation we sometimes use the latter even though K has more than one element.

Given $z = \prod_{i=1}^\eta (h_i, f_i) \in F(H^\Lambda)$ define

$$(3.12) \quad \omega(z) = \sum_{i=1}^\eta l(h_i)(l(h_i) - 1).$$

Define

$$(3.13) \quad |\mathbf{s}| = \max\{l(s) : s \in \mathbf{s}_j, j \in K\},$$

$$(3.14) \quad W_0(\mathbf{z}) = \sum_{i=1}^n \omega(z_i), \text{ where } \omega(z_i) \text{ is defined in (3.12),}$$

$$(3.15) \quad W_1(\mathbf{z}) = \sum_{i=1}^n |z_i|, \text{ where } |z_i| \text{ is defined in (3.6),}$$

$$(3.16) \quad W_2(\mathbf{z}) = \max\{l(h) : (h, f) \text{ occurs in } z_i, \text{ for some } i\} \text{ and}$$

$$(3.17) \quad W_3(\mathbf{z}) = \max\{|\mathbf{s}|, W_2^2(\mathbf{z})\}.$$

We use W_i instead of $W_i(\mathbf{z})$ when appropriate.

Next define

$$(3.18) \quad M(j) = \begin{cases} W_2^2 + 1, & \text{if } j = 0 \\ W_3 W_2^2 (W_2^2 + 2|\mathbf{s}|^2(|\mathbf{s}|/2 + 1)^2), & \text{if } j = 1 \\ 8|\mathbf{s}|^{12} W_2^2 + |\mathbf{s}|, & \text{if } j = 2 \end{cases}$$

and

$$(3.19) \quad M(\mathbf{z}) = \max\{M(j) : j = 0, 1, 2\}.$$

Let the H^Λ -length of z_i be $|z_i| = \eta_i$, for $i = 1, \dots, n$. Define

$$(3.20) \quad \begin{aligned} \widetilde{\mu_1} &= 0 \text{ and} \\ \widetilde{\mu_{j+1}} &= \widetilde{\mu_j} + \eta_j, \text{ for } j = 1, \dots, n \end{aligned}$$

(so $\widetilde{\mu_{n+1}} = W_1$). Write

$$z_j = \prod_{i=1}^{\eta_j} (h_{i+\widetilde{\mu_j}}, f_{i+\widetilde{\mu_j}}),$$

for $j = 1, \dots, n$. Consider the list

$$(h_1, f_1), \dots, (h_{W_1}, f_{W_1})$$

of letters occuring in the above expressions for $z_1, \dots, z_n$. We call the p th element (h_p, f_p) of this list the p th *letter* of $\mathbf{z}$, for $1 \leq p \leq W_1$.

Let $\lambda_1, \dots, \lambda_{|W_1|}$ be distinct elements of Λ . For each p such that (h_p, f_p) is a proper exponential H -letter define

$$\overline{(h_p, f_p)} = (h_p, \lambda_p) \in H^\Lambda.$$

For each p such that (h_p, f_p) is a degenerate exponential H -letter define

$$\overline{(h_p, f_p)} = (h_p \wedge f_p, f_p) \in H^\Lambda.$$

For $j = 1, \dots, n$, let

$$\overline{z_j} = \prod_{i=1}^{\eta_j} \overline{(h_{i+\mu_j}, f_{i+\mu_j})},$$

and define $\overline{\mathbf{z}} = (\overline{z_1}, \dots, \overline{z_n})$. Given p such that (h_p, f_p) is a proper exponential H^Λ -letter let k_p be the unique integer such that $\mathcal{L}$ implies $f_p \equiv k_p \pmod{l(h_p)}$ and $0 \leq k_p < l(h_p)$. Define the *homogeneous* system of parameters $\mathcal{H} = \mathcal{H}(\mathbf{z})$ associated to $\mathbf{z}$ and $\mathcal{L}$ to be the parameter system:

$$\lambda_p - 1 \geq 0,$$

$$\lambda_p - k_p \equiv 0 \pmod{l(h_p)},$$

for all p with $1 \leq p \leq W_1$ such that (h_p, f_p) is a proper exponential H -letter and $l(h_p) > 1$. Then $\overline{\mathbf{z}}$ is constrained by $\mathcal{H}(\mathbf{z})$ and supported over $\mathbf{n}$ and

$$\sum_{i=1}^n |z_i|_\Lambda = \sum_{i=1}^n |\overline{z_i}|_\Lambda,$$

so $\overline{\mathbf{z}} \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{H}, \mathbf{n}, \mathbf{h})$.

Definition 3.5. Let $\mathbf{z} \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$. The *homogeneous* equation associated to $\mathbf{z}$ is the quadratic exponential equation $Q_{\mathcal{H}} = Q_{\mathcal{H}}(\overline{\mathbf{z}}, \mathcal{H}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ associated to $(\overline{\mathbf{z}}, \mathcal{H}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.

Example 3.6. With $I, C_i, R_i, K, X_k, \mathbf{s}_k, H_{*,k}$, and G_k as in Example 3.2(i) let

$$\begin{aligned} z_1 &= (c_{22}, \lambda_1), \\ z_2 &= (c_{12}c_{22}c_{21}, 2)(c_{11}c_{22}c_{21}, f_1)(c_{11}c_{22}, 2), \text{ and} \\ z_3 &= (c_{21}c_{22}c_{11}^2, f_2)(c_{22}c_{11}^2, 2), \end{aligned}$$

where $f_1 = \lambda_1 - 2$ and $f_2 = \lambda_1 - \lambda_2$. Let $\mathcal{L}_0$ and $\mathcal{L}_1$ be as in Example 3.2(i). Then $\mathbf{z} = (z_1, z_2, z_3)$ is constrained by $\mathcal{L}_i$, $i = 1, 2$. Let $n = 3$, $t = p = 0$ and $\mathbf{n} = (1, 2)$. As $\text{supp}(z_1) = X_1$ and $\text{supp}(z_i) = X_2$, $i = 1, 2$, $\mathbf{z}$ is supported over $\mathbf{n}$ with basis $(1, 2)$. Since $t = p = 0$ we have $\mathbf{t} = \mathbf{p} = (0, 0)$, $\mu_j = \xi_j$, for $j = 0, 1, 2$ and $\mu_1 = 0$, $\mu_2 = 1$ and $\mu_3 = 3$. The map β from $\{d_1, d_2, d_3\}$ to $F(H^\Lambda)$ satisfies $\beta(d_i) = z_i$, for $i = 1, 2, 3$. We have $\mathbf{q} = (q_1, q_2)$, where $q_1 = x_1^{-1}d_1x_1$ and $q_2 = x_2^{-1}d_2x_2x_3^{-1}d_3x_3$. Let $Q_i = Q(\mathbf{z}, \mathcal{L}_i, \mathbf{n}, \mathbf{t}, \mathbf{p})$, for $i = 1, 2$ and let W_i be as given in Example 3.2(i). A solution to Q_i is also a solution to W_i . A solution (ϕ, α) to W_i gives rise to solutions for Q_i : a solution is obtained by setting $\phi(x_1)$ equal to any element of $H_{*,1}$ and $\phi(x_2) = 1 \in H_{*,2}$. Since we found solutions for W_1 and W_2 in Example 3.2(i) it follows that $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}_i\text{-genus}(\mathbf{z})$, for $i = 1, 2$.

We have $\omega(z_1) = 0$, $\omega(z_2) = 6$ and $\omega(z_3) = 4$ and $|\mathbf{s}| = 4$ so $W_0 = 10$. We have $|z_1| = 1$, $|z_2| = 3$ and $|z_3| = 2$ so $W_1 = 6$. As $W_2 = 2$ we have $W_3 = 4$. We have $M(0) = 4$, $M(1) = 576$ and $M(2) = 131074$ so $M(\mathbf{z}) = 131074$.

We have $\widetilde{\mu}_1 = 0$, $\widetilde{\mu}_2 = 1$, $\widetilde{\mu}_3 = 4$ and $\widetilde{\mu}_4 = 6$ so

$$\begin{aligned} \bar{z}_1 &= (c_{22}, \lambda_1), \\ \bar{z}_2 &= (c_{12}c_{22}c_{21}, 2)(c_{11}c_{22}c_{21}, \lambda_3)(c_{11}c_{22}, 2), \text{ and} \\ \bar{z}_3 &= (c_{21}c_{22}c_{11}^2, \lambda_6)(c_{22}c_{11}^2, 2), \end{aligned}$$

Then $\mathcal{H}$ consists of the inequalities $\lambda_p - 1 \geq 0$, for $p = 1, 3, 6$, and the congruences $\lambda_p \equiv 0 \pmod{2}$, for $p = 3, 6$. The homogeneous equation associated to $\mathbf{z}$ is $Q_{\mathcal{H}} = (\mathbf{q} = 1, \beta_{\mathcal{H}}, \mathcal{H})$, where $\beta_{\mathcal{H}}(d_i) = \bar{z}_i$. Define ϕ from $L(Q_{\mathcal{H}})$ to $H_{*,1} \dot{\cup} H_{*,2}$ by $\phi(x_i) = 1$, $i = 1, 2$ and $\phi(x_3) = c_{11}c_{22}$. Define α to be the retraction such that $\alpha(\lambda_1) = 6$, $\alpha(\lambda_3) = 4$ and $\alpha(\lambda_6) = 2$. Then (ϕ, α) is a solution to $Q_{\mathcal{H}}$. Compare this solution to the solution (ϕ_1, α_1) to W_i given in Example 3.2(i). We have obtained ϕ from ϕ_1 and set $\alpha(\lambda_p)$ equal to the value $\alpha_1(g_p)$, where g_p is the polynomial occuring in the p th letter of $\mathbf{z}$. The same construction fails using α_0 from Example 3.2(i) because we obtain $\alpha(\lambda_6) = 0$.

Example 3.7. Let $I, C_i, R_i, K, X_k, \mathbf{s}_k, H_{*,k}$, and G_k as in Example 3.3. In this case equations have environment $(X_1; (H_i)_{i \in I}; G_1)^\Lambda$.

- (i) Let $\mathbf{n} = \mathbf{h} = (1)$. The set $(H_{*,1}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ consists of elements of the form $\mathbf{z} = (u_0(h, f)u_1)$, where u_i is a product of degenerate exponential letters, that is $u_i \in F((H_{*,1}, \Lambda)_0)$, for $i = 1, 2$. The equation associated to such $\mathbf{z}$ is $Q = ((x_1^{-1}d_1x_1) = 1, \beta, \mathcal{L})$, where $\beta(d_1) = u_0(h, f)u_1$. This is equivalent to the equation $Q' = ((d_1) = 1, \beta, \mathcal{L})$, in that solutions to Q are solutions to Q' and, given a solution to Q' any value of $\phi(x_1)$ gives a solution to Q . Suppose that $\mathbf{z} = ((h, f))$, $f = \lambda_1$ and $\mathcal{L}$ contains the subsystem $\mathcal{L}_0$ consisting of $\lambda_1 \neq 0$ and $\lambda_1 \equiv 0 \pmod{l(h)}$. Then Q' is the equation of Example 3.3(i). Hence $(\mathbf{n}, (0), (0)) \in \mathcal{L}_0\text{-genus}(\mathbf{z})$ if and only if h has finite order in G_1 .

On the other hand if $\mathbf{z} = ((h_1, \lambda_1)(h_2, l(h_2)))$ then, as in Example 3.3(ii), $(\mathbf{n}, (0), (0)) \in \mathcal{L}_0\text{-genus}(\mathbf{z})$ if and only if $h_1^m = h_2$, for some non-zero integer m .

- (ii) Let $\mathbf{n} = (1)$ and $\mathbf{h} = (2)$. The set $(H_{*,1}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ consists of elements of the form $\mathbf{z} = (z_1)$, where $z_1 \in H_{*,1}^\Lambda$, $z_1 = u_0 w_1 u_1 w_2 u_2$, with $u_i \in F((H_{*,1}, \Lambda)_0)$ and $w_i \in H_{*,1}^\Lambda$. The equation associated to such $\mathbf{z}$ is $Q = ((x_1^{-1} d_1 x_1) = 1, \beta, \mathcal{L})$, where $\beta(d_1) = z_1$. This is equivalent to the equation $Q' = ((d_1) = 1, \beta, \mathcal{L})$, as in the previous case. With $z_1 = (h_1, \lambda_1)(h_2, \lambda_2)$ and $\mathcal{L}_0$ consisting of the inequalities $\lambda_i \neq 0$ and the congruences $\lambda_i \equiv 0 \pmod{l(h_i)}$ it follows that $(\mathbf{n}, (0), (0)) \in \mathcal{L}_0\text{-genus}(\mathbf{z})$ if and only if a non-zero power of h_1 is equal to a non-zero power of h_2 .
- (iii) Let $\mathbf{n} = \mathbf{h} = (2)$. The set $(H_{*,1}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ consists of elements of the form $\mathbf{z} = (z_1, z_2)$, where $z_i \in H_{*,1}^\Lambda$, $z_i = u_i w_i v_i$, with $u_i, v_i \in F((H_{*,1}, \Lambda)_0)$ and either $w_i = (h_1, f_1)(h_2, f_2)$, with $f_1, f_2 \in M \setminus \mathbb{Z}$ and $w_{i-1} \in F((H_{*,1}, \Lambda)_0)$ or $w_1 = (h_1, f_1)$, $w_2 = (h_2, f_2)$, with $f_i \in M$ (and w_i may be trivial). The equation associated to such $\mathbf{z}$ is $Q = ((x_1^{-1} d_1 x_1 x_2^{-1} d_2 x_2) = 1, \beta, \mathcal{L})$, where $\beta(d_i) = z_i$. This is equivalent to the equation $Q' = ((d_1 x_2^{-1} d_2 x_2) = 1, \beta, \mathcal{L})$, in the sense of the previous case of the Example. With $\mathbf{z} = ((h_1, \lambda_1), (h_2, \lambda_2))$ and $\mathcal{L}_0$ as in Example 3.3(iii) it follows that $(\mathbf{n}, (0), (0)) \in \mathcal{L}_0\text{-genus}(\mathbf{z})$ if and only if a non-zero power of h_1 is conjugate to a non-zero power of h_2 .

4. RESOLUTIONS OF QUADRATIC EQUATIONS

Throughout this section the sets I , K and X_k and the groups H_i , $H_{*,k}$ and G_k are defined as in Section 3, (3.1) and (3.2). In particular we assume that H_i has solvable word problem, for all $i \in I$.

Given a k -tuple $(a_1, \dots, a_k)$ of integers we say that the s -tuple $(b_{i_1}, \dots, b_{i_s})$ of integers is a *sub-basis* of $(a_1, \dots, a_k)$ if $(b_1, \dots, b_k)$ is a permutation of $(a_1, \dots, a_k)$ and $1 \leq i_1 < i_2 < \dots < i_s \leq k$.

Definition 4.1. Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$ and basis $(a_1, \dots, a_k)$. Let R be a set $R = \{W_i = (\mathbf{w}_i = 1, \gamma_i, \mathcal{L}_i) : i = 1, \dots, n\}$ of constrained quadratic exponential equations such that W_i has environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$ and basis equal to a sub-basis of $(a_1, \dots, a_k)$, for $i = 1, \dots, n$. Then R is called a *resolution* of W if the following conditions hold.

- (i) If W has a solution then W_i has a solution, for some i .
- (ii) There is an algorithm which, given a solution to W_i , for some i , outputs a solution to W in $(G_k)_{k \in K}$.

A resolution R of W is said to have property P if every element of R has property P .

Theorem 4.2 (cf. [15, Proposition 2.2]). *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation. Then there exists an algorithm which, given input W , outputs a resolution of W in standard form.*

Proof. Let $\mathbf{w} = (w_1, \dots, w_k)$ and let $(b_1^i, \dots, b_{n_i}^i)$ be a boundary labels list for w_i , $i = 1, \dots, k$. Choose $\eta \in \text{Aut}_f(F)$ using Proposition 2.1, such that η is an

admissible transformation of $(\mathbf{w}, \mathbf{b})$ to $(\mathbf{v}, \mathbf{d})$, where $\mathbf{v} = \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p})$, with $\mathbf{n}$, $\mathbf{t}$ and $\mathbf{p}$ as described in Proposition 2.1, and $\eta(F(D)) = F(D)$.

Let $T = L(\mathbf{w})$ and define $U = \cup_{a \in T} L(\eta(a))$ so $L(\mathbf{v}) \subset U$. Since η fixes all but finitely many elements of $D \dot{\cup} X$ there is a finite set $V \subset D \dot{\cup} X$ such that $\eta(F(V)) = F(V)$ and $T \cup U \subset V$. Then $D \cap T \subset D \cap V$ so we may define a map $\beta^* : F(D \cap V) \longrightarrow F(H^\Lambda)$ with $\beta^*(d) = \beta(d)$, for all $d \in D \cap T$ and $\beta^*(d) = 1$, for all $d \notin D \cap T$.

As $\eta(F(D)) = F(D)$ we have $\eta(F(D \cap V)) = F(D \cap V)$, so we may define a map $\delta^* : F(D \cap V) \longrightarrow F(H^\Lambda)$ by $\delta^* = \beta^* \circ \eta^{-1}|_{F(D \cap V)}$. Define $\delta : F(L_D(\mathbf{v})) \longrightarrow F(H^\Lambda)$ by $\delta = \delta^*|_{F(L_D(\mathbf{v}))}$. If $d \in L_D(\mathbf{v})$ then $d = \eta(b_j^i)$, for appropriate i and j , so $\delta(d) = \delta^*(d) = \beta^* \eta^{-1}(d) = \beta^*(b_j^i) = \beta(b_j^i)$. It follows that $(\mathbf{v} = 1, \delta, \mathcal{L})$ is a constrained quadratic exponential equation with the same environment and basis as W .

Suppose (μ, α) is a solution to $(\mathbf{v} = 1, \delta, \mathcal{L})$. Let $T_i = L_X(w_i)$ and $U_i = \cup_{x \in T_i} L(\eta(x))$, for $i = 1, \dots, k$. Let $S_i = L_X(v_i)$ and $V_i = \cup_{x \in S_i} L(\eta^{-1}(x))$, for $i = 1, \dots, k$. Define a map $\mu^* : V \longrightarrow H$ by

$$\mu^*(a) = \begin{cases} \mu(a), & \text{if } a \in L(\mathbf{v}) \\ \hat{\alpha}\delta^*(a), & \text{if } a \in D \cap V \setminus L(\mathbf{v}) \\ \text{any element of } H_{*,a_i} & \text{if } a \in U_i \setminus L(\mathbf{v}) \\ \text{any element of } H & \text{otherwise} \end{cases} ,$$

and extend μ^* to a homomorphism from $F(V)$ to H . (Thus if $a \in D \cap V$ then $\mu^*(a) = \hat{\alpha}\delta^*(a)$.) If $d \in D \cap T$ then $\eta(d) \in F(D \cap U)$ so $\mu^*(\eta(d)) = \hat{\alpha}\delta^*(\eta(d)) = \hat{\alpha}\beta^*(d) = \hat{\alpha}\beta(d)$. Also $\mu^*\eta(a) \in H_{*,a_i}$, for all $a \in L(w_i)$, and $\mu^*\eta(w_i)$ is conjugate in H_{*,a_i} to $\mu^*\eta(w_i) = \mu\eta(w_i) = \mu(v_i) \in N_{a_i}$, for $i = 1, \dots, k$. It follows that, setting $\phi = \mu^* \circ \eta|_{F(T)}$, we have a solution (ϕ, α) to W .

Conversely suppose (ϕ, α) is a solution to W . Define a map $\phi^* : V \longrightarrow H$ by

$$\phi^*(a) = \begin{cases} \phi(a), & \text{if } a \in T \\ \hat{\alpha}\beta^*(a), & \text{if } a \in D \cap V \setminus T \\ \text{any element of } H_{*,a_i} & \text{if } a \in V_i \setminus T \\ \text{any element of } H & \text{otherwise} \end{cases} ,$$

and extend ϕ^* to a homomorphism $\phi^* : F(V) \longrightarrow H$. Define $\mu = \phi^* \circ \eta^{-1}|_{L(\mathbf{v})}$. Then $\mu(a) \in H_{*,a_i}$, for all $a \in L(v_i)$, and $\mu(v_i) = \phi^*\eta^{-1}(v_i)$ is conjugate to $\phi^*(w_i) \in N_{a_i}$, for $i = 1, \dots, k$. If $d \in L_D(\mathbf{v})$, we have $\mu(d) = \phi^*\eta^{-1}(d) = \hat{\alpha}\beta^*\eta^{-1}(d) = \hat{\alpha}\delta^*(d) = \hat{\alpha}\delta(d)$. Hence (μ, α) is a solution to $(\mathbf{v} = 1, \delta, \mathcal{L})$. It follows that $R = \{(\mathbf{v} = 1, \delta, \mathcal{L})\}$ is a resolution of W in standard form. Since $\eta, U_i, V_i, \beta^*, \delta^*, L(\mathbf{w})$ and $L(\mathbf{v})$ may all be effectively constructed there exists an algorithm as claimed.

The process of finding $\eta, \mathbf{v}, U, V, \beta^*, \delta^*, L(\mathbf{w})$ and $L(\mathbf{v})$ is called *reduction to standard form*.

Let u be an element of $F(H^\Lambda)$ such that u is constrained by a parameter system $\mathcal{L}$. We say $(u, \mathcal{L})$ has $H^\Lambda(i)$ -*redundancy* if condition i below holds and that $(u, \mathcal{L})$ has H^Λ -*redundancy* if $(u, \mathcal{L})$ has $H^\Lambda(i)$ -redundancy for some i , with $1 \leq i \leq 7$.

$i=1$: A generator $(h, f) \in H^\Lambda$ occurs in u with $h = 1 \in H$.

$i=2$: A generator $(h, f) \in H^\Lambda$ occurs in u and $\mathcal{L}$ implies that $f = 0$.

- $i=3$: A proper exponential H -letter (a, f) occurs in u with $a = a_0^s$, where a_0 is a cyclically reduced element of H , $s \in \mathbb{Z}$, $|s| > 1$ and $l(a) = m > 1$.
- $i=4$: A generator $(h, f) \in H^\Lambda$ occurs in u with exponent -1 .
- $i=5$: The word u contains a subword $(a_1, f_1) \cdots (a_n, f_n)$, where $(a_i, f_i) \in H^\Lambda$ and either
 - (a) $n > 1$ and $f_1, \dots, f_n \in \mathbb{Z}$ or
 - (b) there is at least one i such that (a_i, f_i) is a proper exponential H -letter and, for each $i = 1, \dots, n$, if (a_i, f_i) is a proper exponential H -letter then $\mathcal{L}$ implies that $f_i = m_i \in \mathbb{Z}$.
- $i=6$: The word u contains a subword $(a, f)(b, g)$, where $(a, f), (b, g) \in H^\Lambda$ at least one of which is a proper exponential H -letter, $l(a) = l(b) = m$, $\mathcal{L}$ implies that $f \equiv k(\pmod{m})$, for some k with $0 \leq k < m$, $a = g_1 \cdots g_m$, $g_i \in H_{k_i}$ and $b^\varepsilon = g_{k+1} \cdots g_m g_1 \cdots g_k$, with $\varepsilon = \pm 1$.
- $i=7$: The word u contains a subword $(a, f)(b, g)$, where $(a, f), (b, g) \in H^\Lambda$ at least one of which is a proper exponential H -letter, $l(a) = m$, $l(b) = n$, $a = a_1 \cdots a_m$, $b = b_1 \cdots b_n$, $a_i \in H_{k_i}$, $b_j \in H_{l_j}$, $\mathcal{L}$ implies that $f \equiv k(\pmod{m})$, for some k with $0 \leq k < m$, and $a_k = b_1^{-1} \in H$.

If $(u, \mathcal{L})$ has no H^Λ -redundancy it is said to be H^Λ -irredundant. The equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ is said to be H^Λ -redundant if $(\beta(d), \mathcal{L})$ is H^Λ -redundant, for some $d \in L_D(\mathbf{w})$, and H^Λ -irredundant otherwise.

If $s \in \mathbb{Z}$, with $1 \leq s \leq 7$, and $(u, \mathcal{L})$ is $H^\Lambda(i)$ -irredundant for $i = 1, \dots, s$ we say $(u, \mathcal{L})$ is $H^\Lambda(1, s)$ -irredundant. The equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ is $H^\Lambda(1, s)$ -irredundant if $(\beta(d), \mathcal{L})$ is $H^\Lambda(1, s)$ -irredundant, for all $d \in L_D(\mathbf{w})$.

Given a constrained quadratic exponential equation $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ and $d \in L_D(\mathbf{w})$, suppose $(\beta(d), \mathcal{L})$ has $H^\Lambda(i)$ -redundancy. For each i we define a set $R(W, i)$ of constrained quadratic exponential equations. When $i \neq 4$ the set $R(W, i) = \{W'\}$, where $W' = (\mathbf{w} = 1, \beta', \mathcal{L})$ with β' obtained from β by setting $\beta'(d') = \beta(d')$, if $d' \neq d$ and $\beta'(d)$ equal to an element of $F(H^\Lambda)$ dependent on i as follows.

- $i=1$ or 2 : Delete, from $\beta(d)$, all occurrences of the generator (h, f) at which redundancy occurs and freely reduce the resulting word, to give $\beta'(d)$.
- $i=3$: If $H^\Lambda(3)$ -redundancy occurs at a subword (a, f) of $\beta(d)$, where $a = a_0^s$, as in the description of $H^\Lambda(3)$ -redundancy above, then replace all occurrences of (a, f) in $\beta(d)$ by (a_0, f) , to give $\beta'(d)$. Note that, if $\mathcal{L}$ implies that $f \equiv k(\pmod{l(a)})$ then $\mathcal{L}$ implies that $f \equiv k(\pmod{l(a_0)})$.
- $i=5$: Suppose $H^\Lambda(5)$ -redundancy occurs at a subword $(a_1, f_1) \cdots (a_n, f_n)$ of $\beta(d)$. For i such that (a_i, f_i) is a proper exponential H -letter set $m_i \in \mathbb{Z}$ such that $\mathcal{L}$ implies $f_i = m_i$. For all other i set $m_i = f_i$. Set $c_i = a_i \wedge m_i$, for $i = 1, \dots, n$, to give $\beta'(d)$. Replace $(a_1, f_1) \cdots (a_n, f_n)$ in $\beta(d)$ by (c, h) , where $c = c_1 \cdots c_n$ and $h = l(c) \in \mathbb{Z}$.
- $i=6$: If $H^\Lambda(6)$ -redundancy occurs at a subword $(a, f)(b, g)$ of $\beta(d)$ then replace all occurrences of $(a, f)(b, g)$ in $\beta(d)$ by $(a, f + \varepsilon g)$, to give $\beta'(d)$. Note that, if $\mathcal{L}$ implies that $f \equiv k(\pmod{m})$ and $g \equiv l(\pmod{m})$, then $\mathcal{L}$ implies that $f + \varepsilon g \equiv k + \varepsilon l(\pmod{m})$.
- $i=7$: Suppose $H^\Lambda(7)$ -redundancy occurs at a subword $(a, f)(b, g)$ of $\beta(d)$, as described in the case $i=7$ above. We obtain $\beta'(d)$ from $\beta(d)$ as follows. If (a, f) and (b, g) are both proper exponential H -letters then replace all occurrences of $(a, f)(b, g)$ in $\beta(d)$ by $(a, f - 1)(b_2 \cdots b_n b_1, g - 1)$. If

(a, f) is a proper exponential H -letter but (b, g) is not then replace all occurrences of $(a, f)(b, g)$ in $\beta(d)$ by $(a, f-1)(\tau(b, g-1), g-1)$. If (b, g) is a proper exponential H -letter but (a, f) is not then replace all occurrences of $(a, f)(b, g)$ in $\beta(d)$ by $(\iota(a, f-1), f-1)(b_2 \cdots b_n b_1, g-1)$.

We consider finally the case where W has occurrence of $H^\Lambda(4)$ -redundancy.

$i=4$: Suppose $\beta(d) = w_0(h, f)^{-1}w_1$, where $w_i \in F(H^\Lambda)$, $(h, f) \in H^\Lambda$ and $\beta(d)$ is reduced as written. If (h, f) is a degenerate exponential H -letter then $f = l(h)$ and we define $\beta_1 : F(L_D(\mathbf{w})) \rightarrow H$ by $\beta_1(d) = w_0(h^{-1}, f)w_1$ and $\beta_1(d') = \beta(d')$, for $d' \neq d$. Then $R(W, 4) = \{(\mathbf{w} = 1, \beta_1, \mathcal{L})\}$.

On the other hand, if (h, f) is a proper exponential H -letter, let $m = l(h)$ and let $k \in \mathbb{Z}$, such that $0 \leq k < m$ and $\mathcal{L}$ implies that $f \equiv k \pmod{m}$. For $i = 1, 2$ define $\beta_i : F(L_D(\mathbf{w})) \rightarrow F(H^\Lambda)$ by $\beta_i(v) = \beta(v)$, if $v \neq d$,

$$\beta_1(d) = w_0([\iota(h, k)^{-1}\tau(h, m-k)^{-1}], f)w_1$$

and

$$\beta_2(d) = w_0([\tau(h, k)\iota(h, m-k)], -f)w_1.$$

Let $\mathcal{L}_1 = \mathcal{L} \cup \{f \geq 0\}$, let $\mathcal{L}_2 = \mathcal{L} \cup \{-f > 0\}$. Then any solution α to $\mathcal{L}$ is a solution to either $\mathcal{L}_1$ or $\mathcal{L}_2$ and, if α is a solution to $\mathcal{L}_i$, we have

$$\hat{\alpha}\beta(d') = \hat{\alpha}\beta_i(d'), \text{ for all } d' \in L_D(\mathbf{w}), i = 1, 2.$$

Define $W_i = (\mathbf{w} = 1, \beta_i, \mathcal{L}_i)$, for $i = 1, 2$ and set $R(W, 4) = \{W_1, W_2\}$.

Lemma 4.3. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation. Then there exists an algorithm which, given W , outputs an $H^\Lambda(1, 4)$ -irredundant resolution R of W in standard form.*

Proof. We may assume, using Theorem 4.2, that W is in standard form. The solvability of the word problem for H allows us to determine whether W contains an occurrence of $H^\Lambda(i)$ -redundancy, for some fixed i , $1 \leq i \leq 3$, and to find an occurrence if one exists. Replacing W by the unique equation of the set $R(W, i)$ we obtain an equation W' with fewer occurrences of $H^\Lambda(i)$ -redundancy than W . Repeating this process we eventually obtain an equation W' which is $H^\Lambda(i)$ -irredundant, for $i = 1, 2$ and 3. We may now eliminate an occurrence of $H^\Lambda(4)$ -redundancy by replacing $\{W'\}$ with $R(W', 4)$, if necessary. Note that, by construction, no element of $R(W', 4)$ contains an occurrence of $H^\Lambda(i)$ -redundancy, with $i = 1, 2$ or 3. We may repeat the removal of occurrences of $H^\Lambda(4)$ -redundancy on elements of $R(W', 4)$ to obtain a set R of $H^\Lambda(1, 4)$ -irredundant, constrained, quadratic exponential equations.

It remains to show that if W is a constrained quadratic exponential equation then $R(W, i)$ is a resolution of W , for $i = 1, 2, 3, 4$. This is easy to see if $i = 1$ or 2.

Suppose we have $H^\Lambda(3)$ -redundancy at the occurrence of (a, f) in $\beta(d) = w_0(a, f)^\varepsilon w_1$, where $\varepsilon = \pm 1$ and $d \in L_D(\mathbf{w})$. As W is constrained, $\mathcal{L}$ implies that $f \equiv k \pmod{m}$, for some integer k , with $0 \leq k < m$. Let a_0 be a cyclically reduced word and $s > 1$ an integer such that $a = a_0^s$. By definition of $H^\Lambda(3)$ -redundancy $l(a_0) > 0$. Set $n = l(a_0)$ and let $p, l \in \mathbb{Z}$ such that $k = pn + l$, with $0 \leq l < n$. Assume α is a solution to $\mathcal{L}$. Then, if $\alpha(f) \geq 0$, say $\alpha(f) = qsn + k = qm + k$, we have $\hat{\alpha}(a, f) = a^q \iota(a, k) = a_0^{sq} a_0^p \iota(a_0, l) = a_0 \wedge (qsn + pn + l) = \hat{\alpha}(a_0, f)$. On the other hand, if $\alpha(f) < 0$, say $-\alpha(f) = qm + k$, then $\hat{\alpha}(a, f) = a^{-q} \iota(a^{-1}, k) = a_0^{-sq} a_0^{-p} \iota(a_0^{-1}, l) = a_0 \wedge -(qsn + pn + l) = \hat{\alpha}(a_0, f)$. Hence, if α is a solution to $\mathcal{L}$

then $\hat{\alpha}\beta(d) = \hat{\alpha}\beta'(d)$, for all $d \in L_D(\mathbf{w})$. Therefore solutions of W and W' coincide and $R(W, 3)$ is a resolution of W .

When $H^\Lambda(4)$ -redundancy occurs at a exponential H -letter which is not proper, then the resulting set $R(W, 4)$ is clearly a resolution of W . In the remaining case a solution to W is also a solution to W_i , for $i = 1$ or 2 . Conversely given a solution to W_i it is a solution to W . It follows that $R(W, i)$ is a resolution of W , for $i = 1, 2, 3$ and 4 .

As $\mathbf{w}$ is the quadratic word appearing in every element of each of the resolutions used above to replace W , the resolution obtained is also in standard form.

The application of the algorithm of Lemma 4.3 to produce an $H^\Lambda(1, 4)$ -irredundant resolution of an equation is called $H^\Lambda(1, 4)$ -reduction.

The *dual* of a exponential H -letter (a, f) is

$$(a, f)^* = (a^{-1}, -f).$$

Note that $((a, f)^*)^* = (a, f)$. There is a well-defined map $\nu : H^\Lambda \rightarrow F(H^\Lambda)$ given by $\nu(a, f) = (a, f)^*$ and $\nu(b, g) = (b, g)$, if $(b, g) \neq (a, f)$. Such a map induces an endomorphism ν of $F(H^\Lambda)$, called the *dualization* at (a, f) .

We define the *deletion* at $(a, f) \in H^\Lambda$ to be the endomorphism of $F(H^\Lambda)$ defined by $\delta(a, f) = \delta((a, f)^*) = 1$ and $\delta(b, g) = (b, g)$, for all $(b, g) \notin \{(a, f), (a, f)^*\}$.

Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation. Dualization at (a, f) is *admissible* for W if (a, f) occurs in $\beta(d)$, for some $d \in L_D(\mathbf{w})$ and if $\mathcal{L} \cup \{-f > 0\}$ is consistent. Deletion at (a, f) is *admissible* for W if (a, f) occurs in $\beta(d)$, for some $d \in L_D(\mathbf{w})$ and if $\mathcal{L} \cup \{f = 0\}$ is consistent.

Define $\mathcal{M}(\mathbf{w}, \beta) = \{f \in M : (a, f) \text{ occurs in } (\mathbf{w}, \beta)\}$. Then W is called *positive* if it is $H^\Lambda(1, 4)$ -irredundant and $\mathcal{L}$ implies that $f > 0$, for all $f \in \mathcal{M}(\mathbf{w}, \beta)$.

Lemma 4.4. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be an $H^\Lambda(1, 4)$ -irredundant, consistent, quadratic exponential equation. Then there exists an algorithm which, when given W , outputs a positive resolution of W in standard form.*

Proof. If (a, f) is a degenerate exponential H -letter occurring in W and $f < 0$ then we may replace (a, f) by $(a^{-1}, -f)$. We may therefore restrict attention to proper exponential H -letters. Given $f \in \mathcal{M}(\mathbf{w}, \beta)$ define $H^\Lambda(\mathbf{w}, \beta, f) = \{(a, f) \in H^\Lambda : (a, f) \text{ occurs in } (\mathbf{w}, \beta)\}$. Suppose $f \in \mathcal{M}(\mathbf{w}, \beta)$ and assume that $H^\Lambda(\mathbf{w}, \beta, f) = \{(a_1, f), \dots, (a_n, f)\}$ and that $H^\Lambda(\mathbf{w}, \beta, -f) = \{(a_{n+1}, -f), \dots, (a_{n+p}, -f)\}$. If $\mathcal{L} \cup \{f < 0\}$ is consistent, let ν_i be the dualization at (a_i, f) , $i = 1, \dots, n+p$, define $\nu^f = \nu_n \circ \dots \circ \nu_1$ and set $W_-(f) = (\mathbf{w} = 1, \nu^f \circ \beta, \mathcal{L} \cup \{-f > 0\})$. If $\mathcal{L} \cup \{f = 0\}$ is consistent, let δ_i be the deletion at (a_i, f) . Define $\delta^f = \delta_{n+p} \circ \dots \circ \delta_n \circ \dots \circ \delta_1$ and set $W_0(f) = (\mathbf{w} = 1, \delta^f \circ \beta, \mathcal{L} \cup \{f = 0\})$. Finally, if $\mathcal{L} \cup \{f > 0\}$ is consistent, define $\nu^{-f} = \nu_{n+p} \circ \dots \circ \nu_{n+1}$ and set $W_+(f) = (\mathbf{w} = 1, \beta, \mathcal{L} \cup \{f > 0\})$. Let R_0 be the set consisting of whichever of $W_-(f)$, $W_0(f)$ and $W_+(f)$ are defined. As $\mathcal{L}$ is consistent R_0 is non-empty.

Note that for any solution α to $\mathcal{L}$ we have

$$\begin{aligned} \hat{\alpha}(a, f) &= a \wedge \alpha(f) &= a^{-1} \wedge -\alpha(f) \\ &= a^{-1} \wedge \alpha(-f) &= \hat{\alpha}(a^{-1}, -f) \\ &= \hat{\alpha}((a, f)^*). \end{aligned}$$

Hence (ϕ, α) is a solution to W with $\alpha(f) < 0$ if and only if (ϕ, α) is a solution to $W_-(f)$. Furthermore (ϕ, α) is a solution to W with $\alpha(f) > 0$ or $\alpha(f) = 0$ if and

only if (ϕ, α) is a solution to $W_+(f)$ or $W_0(f)$, respectively. Thus R_0 is a resolution of W and as $\mathcal{L}$ is linear there is an algorithm which constructs R_0 from input W .

Let $\mathcal{N}(W) = \{f \in \mathcal{M}(\mathbf{w}, \beta) : \mathcal{L} \cup \{f \leq 0\} \text{ is consistent}\}$ and suppose $f \in \mathcal{N}(W)$. By definition, $f \notin \mathcal{N}(W_+(f))$. Also $f \notin \mathcal{N}(W_0(f)) \cup \mathcal{N}(W_-(f))$ as $H^\Lambda(\mathbf{w}, \delta^f \circ \beta, f) = H^\Lambda(\mathbf{w}, \nu^f \circ \beta, f) = \emptyset$. Similarly $-f \notin \mathcal{N}(W_0(f)) \cup \mathcal{N}(W_-(f)) \cup \mathcal{N}(W_+(f))$. It follows that, for $W' \in R_0$, we have $|\mathcal{N}(W')| < |\mathcal{N}(W)|$.

No $H^\Lambda(i)$ -redundancy, with $i = 1, 3$ or 4 , is introduced in the process of forming the elements of R_0 . If $H^\Lambda(2)$ -redundancy occurs in some element W' of R_0 , then its subsequent removal does not increase $|\mathcal{N}(W')|$. Thus we may assume that R_0 is $H^\Lambda(1, 4)$ -irredundant and repeat the above process on elements of R_0 . Eventually a positive resolution of W is obtained. As in the proof of Lemma 4.3 we may assume that W and this resolution of W are both in standard form.

The process of forming a positive resolution is called *positive $H^\Lambda(1, 4)$ -reduction*.

Lemma 4.5. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained, quadratic exponential equation. Then there exists an algorithm which, when given W , outputs an H^Λ -irredundant, positive resolution of W in standard form.*

Proof. Lemma 4.4 implies that we may assume that W is positive. If W contains an occurrence of $H^\Lambda(5)$ -redundancy then the unique equation W' of $R(W, 5)$, described above, is positive. Clearly $R(W, 5)$ is a resolution of W and W' has fewer occurrences of $H^\Lambda(5)$ -redundancy than W . Furthermore the sum of exponential length and H^Λ -length decreases in passing from W to W' . Then W' may be $H^\Lambda(1, 4)$ -reduced, if necessary, to produce a resolution of W every element of which is positive and has smaller exponential length or H^Λ -length than W . We may continue this process to arrive at a resolution which is positive and $H^\Lambda(1, 5)$ -irredundant. We call the process of forming this resolution *positive $H^\Lambda(1, 5)$ -reduction*.

Note that the solvability of the word problem in H allows the identification of occurrences of $H^\Lambda(6)$ - and $H^\Lambda(7)$ -redundancy. Assume that W is an element of the positive $H^\Lambda(1, 5)$ -irredundant resolution obtained in the previous paragraph. Given an occurrence of $H^\Lambda(6)$ -redundancy we note that, as W is positive, the set $R(W, 6)$ defined above is a resolution of W . Also, the unique element of $R(W, 6)$ has smaller H^Λ -length than W . Hence we may form the resolution $R(W, 6)$, to each element of which we apply positive $H^\Lambda(1, 5)$ -reduction, if necessary, and repeat the process. As none of the operations involved increases H^Λ -length, this process terminates and the resulting resolution is positive and $H^\Lambda(1, 6)$ -irredundant. We call the process of forming this resolution *positive $H^\Lambda(1, 6)$ -reduction*.

Given W in the above resolution, suppose W contains an occurrence of $H^\Lambda(7)$ -redundancy at $(a, f)(b, g)$, as described in the definition of H^Λ -redundancy. The set $R(W, 7)$ is a resolution of W , as W is positive. Let W' be the unique element of this set. As above, we may form a positive $H^\Lambda(1, 6)$ -irredundant resolution R' , of W' , which is also a resolution of W . Elements of $R(W, i)$, where $i = 1, 2$ or 6 have smaller H^Λ -length than W . The element of $R(W, 3)$ has smaller H -length than W . In the case of $W' \in R(W, 5)$ the sum of the H^Λ -length and exponential length is smaller than that of W . No operation involved in positive $R(1, 6)$ -reduction increases any of H^Λ -length, H -length or exponential length. Furthermore once $H^\Lambda(4)$ -redundancy has been removed it is not reintroduced by any subsequent removal of $H^\Lambda(i)$ -redundancy, for any other i . Hence we may assume that in this

application of positive $H^\Lambda(1,6)$ -reduction to form R' we either reduce the sum of H^Λ -length, H -length and exponential length, or do nothing, and so introduce no new occurrence of $H^\Lambda(7)$ -redundancy. (Note that if $\mathcal{L}$ implies $f-1$ or $g-1$ is equal to 0 then the process of H^Λ -reduction removes (a, f) or (b, g) from the image of $L_D(\mathbf{w})$ under β .)

In forming R' we construct β' by replacing occurrences of $u_0 = (a, f)(b, g)$ by a word u_1 . We call this the $R7$ -process at u_0 . If $(u_1, \mathcal{L})$ is $H^\Lambda(7)$ -redundant the $R7$ -process can be repeated at u_1 , replacing u_1 with u_2 . If this can be repeated n times we obtain a sequence $u_0, \dots, u_n$ and we say that the $R7$ -process can be repeated n times at u_0 . Suppose that there exists an integer K (dependent only on (a, f) and (b, g)) such that the $R7$ -process can be repeated only K times, at $(a, f)(b, g)$. Then, provided this holds at all occurrences of $H^\Lambda(7)$ -redundancy, the $R7$ -process can be repeated at any further occurrences of $H^\Lambda(7)$ -redundancy to form an H^Λ -irredundant, positive resolution of W .

If (a, f) is a degenerate exponential H -letter then $f \in \mathbb{Z}$ and the $R7$ -process can be repeated at most f times at $(a, f)(b, g)$. A similar statement holds if (b, g) is degenerate. Suppose then that (a, f) and (b, g) are proper exponential H -letters. Let $l(a) = m$, $l(b) = n$ and assume first that $\mathcal{L}$ implies that $f \equiv 0 \pmod{m}$. Suppose also that the $R7$ -process can be repeated $m+n$ times at $(a, f)(b, g)$. Then, for all r with $1 \leq r \leq m+n$, we have $a^{-1} \wedge r = b \wedge r$. This implies in particular that $m = 1$ if and only if $n = 1$. If $m = n$ then W has $H^\Lambda(6)$ -redundancy, so we may assume $m \neq n$ and $m, n > 1$. Let $r = m+n$ and $u = a^{-1} \wedge r = b \wedge r$. Then u has periods m and n so, from [16], u has period $e = \gcd(m, n)$. As $e|m$ and $e|n$, both a and b have period e . Thus W also has $H^\Lambda(3)$ -redundancy, a contradiction. Hence, when $\mathcal{L}$ implies $f \equiv 0 \pmod{m}$, the $R7$ -process cannot continue for more than $m+n-1$ iterations at $(a, f)(b, g)$.

Now suppose that $\mathcal{L}$ implies $f \equiv k \pmod{m}$, with $0 \leq k \leq m-1$. After k iterations of the $R7$ -process $(a, f)(b, g)$ is replaced by $(a, f-k)(b', g-k)$, for some cyclic permutation b' of b . As $\mathcal{L}$ implies $f-k \equiv 0 \pmod{m}$, it follows that at most $m+n-1$ further iterations are possible. Hence the $R7$ -process cannot be repeated more than $m+n+k-1 < 2m+n-1$ times at $(a, f)(b, g)$.

Hence in all cases there is an integer K computable from $(a, f)(b, g)$ such that the $R7$ -process can be repeated only K times at $(a, f)(b, g)$. Thus, we can effectively form an H^Λ -irredundant, positive resolution of W . As in the proof of Lemma 4.3 we may assume that W and this resolution of W are both in standard form. This completes the proof of the Lemma.

Let $u \in F(H^\Lambda)$ constrained by a system of parameters $\mathcal{L}$. We say $(u, \mathcal{L})$ is *cyclically H^Λ -redundant* if $(u', \mathcal{L})$ is H^Λ -redundant, for some cyclic permutation u' of u . If $(u, \mathcal{L})$ is H^Λ -irredundant and $(u', \mathcal{L})$ is cyclically H^Λ -redundant for some cyclic permutation of u we say $(u, \mathcal{L})$ is *strictly cyclically H^Λ -redundant*. In this case if $u = p_1 \cdots p_n$, with $p_i \in (H^\Lambda)^{\pm 1}$ then the word $p_n p_1$ must be $H^\Lambda(i)$ -redundant, for $i = 5, 6$ or 7 .

The equation $(\mathbf{w} = 1, \beta, \mathcal{L})$ is *cyclically H^Λ -redundant* if $\beta(d)$ is cyclically H^Λ -redundant for some $d \in L_D(\mathbf{w})$ and *cyclically H^Λ -irredundant* otherwise.

Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be H^Λ -irredundant. If $d \in L_D(\mathbf{w})$ and $\beta(d)$ is cyclically H^Λ -redundant then we have $\beta(d) = (b, g)p(a, f)$, where $(b, g), (a, f) \in H^\Lambda$, $p \in F(H^\Lambda)$, and $(a, f)(b, g)$ has $H^\Lambda(i)$ -redundancy, with $i = 5, 6$ or 7 . Given this

situation we define a set $R'(W, i) = \{(\mathbf{w} = 1, \beta_i, \mathcal{L})\}$, where $\beta_i(d') = \beta(d')$, for $d' \neq d$ and $\beta_i(d)$ is dependent on i as follows.

- $i=5$: $\beta_5(d) = p(c, h)$, where (c, h) is as defined in $R(W, 5)$.
- $i=6$: $\beta_6(d) = p(a, f + \varepsilon g)$, where (a, f) , (b, g) and ε are as described in the definition of $H^\Lambda(6)$ -redundancy.
- $i=7$: Let $(a, f)(b, g)$ satisfy the conditions for $H^\Lambda(7)$ -redundancy. In the case where (a, f) and (b, g) are both proper exponential H -letters then let $\beta_7(d) = (b_2 \cdots b_n b_1, g - 1)p(a, f - 1)$. If (a, f) is a proper exponential H -letter but (b, g) is not then $\beta_7(d) = (\tau(b, g - 1), g - 1)p(a, f - 1)$. If (b, g) is a proper exponential H -letter but (a, f) is not then $\beta_7(d) = (b_2 \cdots b_n b_1, g - 1)p(\iota(a, f - 1), f - 1)$.

Lemma 4.6. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation. Then there exists an algorithm which, given W , outputs a cyclically H^Λ -irredundant, positive resolution of W in standard form.*

Proof. Given Lemma 4.5 we may assume W is H^Λ -irredundant and positive. As in the proof of Lemma 4.5 we may use the sets $R'(W, i)$ to remove occurrences of H^Λ -redundancy in $p(a, f)(b, g)$, at each stage replacing W by a set with elements which are H^Λ -irredundant, positive and such that either the sum of H^Λ -length, H -length and exponential length is smaller or there are fewer occurrences of cyclic H^Λ -redundancy than at the previous stage. It therefore suffices to show that each of the sets $R'(W, i)$ is a resolution of W .

Suppose we have an occurrence of cyclic H^Λ -redundancy at $\beta(d)$, as described above, and suppose d occurs in w_i . Let $w_i = u_0 x^{-1} d x u_1$ and $\beta(d) = (b, g)p(a, f)$. Suppose (ϕ, α) is a solution to W . Define $\phi' : D \cup X \rightarrow H$ by $\phi'(v) = \phi(v)$, if $v \in D \cup X \setminus \{x, d\}$. Define $\phi'(d) = \hat{\alpha}\beta_i(d)$. If $i = 5$ or 6 define $\phi'(x) = (b \wedge \alpha(g))^{-1}\phi(x)$. If $i = 7$ define $\phi'(x) = b_1^{-1}\phi(x)$. Then clearly $\phi'|_{L_D(\mathbf{w})} = \hat{\alpha}\beta_i$ and it is easy to check that $\phi'(w_i) = \phi(w_i)$. Hence (ϕ', α) is a solution to the unique element W_i of $R'(W, i)$. Conversely, if (ϕ', α) is a solution to W_i , for some i , then define $\phi : F(D \cup X) \rightarrow H$ by $\phi(v) = \phi'(v)$, for $v \in D \cup X \setminus \{x, d\}$, $\phi(d) = \hat{\alpha}\beta(d)$ and $\phi(x)$ using the same rules as above, relating ϕ and ϕ' . Then (ϕ, α) is a solution to W . Hence $R'(W, i)$ is a resolution of W in each case. As in the proof of Lemma 4.3 we may assume that W and these resolutions of W are in standard form.

Let a be a reduced word in H and $\mathbf{s}$ a set of words of H . Suppose that a has a reduced subword u such that uv is a cyclic permutation of an element of $\mathbf{s}^{\pm 1}$, for some $v \in H$, with $l(v) < l(u)$. Then we say that a is *relator-reducible* with respect to $(H, \mathbf{s})$. In this case, if $a = a_0 u a_1$, where $l(a) = l(a_0) + l(u) + l(a_1)$ and $a_i \in H$, we say that the reduced word representing $a_0 v^{-1} a_1 \in H$ is an *elementary relator-reduction* of a , with respect to $(H, \mathbf{s})$, *replacing* u by v^{-1} . If a is not relator-reducible it is said to be *relator-reduced*, with respect to $(H, \mathbf{s})$.

Let N be the normal closure of $\mathbf{s} \subset H$. If the word problem is solvable in H then given $a \in H$ we may determine whether or not a is relator-reducible and if so perform an elementary relator-reduction. This decreases the length of a and so we may perform further elementary relator-reductions until a relator-reduced word $a' \in H$, with $aN = a'N$, is obtained. Then a' is called a *relator-reduction* of a , with respect to $(H, \mathbf{s})$. A degenerate exponential H -letter $(a, f) \in H^\Lambda$ is said to be *relator-reducible*, with respect to $(H, \mathbf{s})$, if a is relator-reducible, with respect to

$(H, \mathbf{s})$. A non-degenerate exponential H -letter $(a, f) \in H^\Lambda$ is said to be *relator-reducible* if some cyclic permutation of a is relator-reducible. A exponential H -letter which is not relator-reducible is called *relator-reduced*. Let $\mathbf{w} = \{w_1, \dots, w_k\}$ be a system of quadratic words. The equation $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$ and basis $(a_1, \dots, a_k)$ is said to be *relator-reducible* if an element $(a, f) \in H_{*, a_i}^\Lambda$, which is relator-reducible with respect to $\mathbf{s}_{a_j}$, occurs in $\beta(d)$ for some $d \in L_D(w_i)$. Otherwise W is said to be *relator-reduced*.

Lemma 4.7. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained, quadratic exponential equation. Then there exists an algorithm which, given W , outputs a relator-reduced, cyclically H^Λ -irredundant, positive resolution of W in standard form.*

Proof. Given Lemma 4.6 we may assume W is cyclically H^Λ -irredundant and positive. Suppose a relator-reducible H_{*, a_i} -letter (a, f) occurs in $\beta(d)$, for some $d \in L_D(w_i)$, with $\beta(d) = p_0(a, f)p_1$. Assume first that (a, f) is degenerate. Define an equation $W' = (\mathbf{w} = 1, \beta', \mathcal{L})$ as follows. Let $\beta'(d') = \beta(d')$, for all $d' \in L_D(\mathbf{w}) \setminus \{d\}$ and $\beta'(d) = p_0(a', l(a'))p_1$, where a' is a relator-reduction of a with respect to $(H_{*, a_i}, \mathbf{s}_{a_i})$. Clearly solutions of W and W' coincide and W' has smaller H -length than W . We may now replace W' by a positive, cyclically H^Λ -irredundant resolution of W' , observing that none of the operations involved increases H -length. This process may be repeated until a positive, cyclically H^Λ -irredundant resolution of W is obtained, with no relator-reducible degenerate exponential H -letters.

Assume then that all relator-reducible exponential H -letters occurring in W are proper and that (a, f) is relator-reducible and occurs in $\beta(d)$, with $d \in L_D(w_i)$ and that $\beta(d) = p_0(a, f)p_1$. Define $\text{crr}(W)$ to be the sum of $l_H(a, f)$ over all occurrences of relator-reducible proper exponential H -letters occurring in W .

Suppose first that a is relator-reducible and has relator-reduction a_0 . Let $a_0 = b^{-1}a_1b$, where a_1 is cyclically reduced and b is reduced. Suppose $\mathcal{L}$ implies that $f \equiv k \pmod{l(a)}$, with $0 \leq k < l(a)$. Let $\lambda \in \Lambda$ such that λ does not occur in W and let $c = \iota(a, k)$. Suppose c has relator-reduction c_0 . Define equations $W_i = (\mathbf{w} = 1, \beta_i, \mathcal{L}_i)$, for $i = 0$ and 1 , as follows. Let $\beta_i(d') = \beta(d')$, for all $d' \neq d$, let $\beta_0(d) = p_0(c_0, l(c_0))p_1$ and let $\beta_1(d) = p_0(b^{-1}, l(b))(a_1, \lambda l(a_1))(bc_0, l(bc_0))p_1$. Set $\mathcal{L}_0 = \mathcal{L} \cup \{f = k\}$ and $\mathcal{L}_1 = \mathcal{L} \cup \{g = 0, \lambda > 0\}$, where $g = f - k - \lambda l(a)$.

The pair (ϕ, α) is a solution to W_0 if and only (ϕ, α) is a solution to W with $\alpha(f) = k$. Suppose (ϕ, α) is a solution to W . If $\alpha(f) = ql(a) + k$, then we may assume that $q \geq 0$, since W is positive, and in the light of the previous remark we assume that $q > 0$. Given that $q > 0$, define a retraction $\alpha' : M \rightarrow \mathbb{Z}$ by $\alpha'(\mu) = \alpha(\mu)$, for all $\mu \in \Lambda \setminus \{\lambda\}$, and $\alpha'(\lambda) = q$. We have $\hat{\alpha}'\beta_1(d) = \hat{\alpha}(p_0)b^{-1}a_1^qbc_0\hat{\alpha}(p_1)$, so $\hat{\alpha}'\beta_1(d)N_{a_i} = \hat{\alpha}\beta(d)N_{a_i}$. Thus, if $\phi' : F(L_D(\mathbf{w})) \rightarrow H$ is defined by $\phi'(a) = \phi(a)$, for $a \neq d$ and $\phi'(d) = \hat{\alpha}'\beta_1(d)$ we have $\phi'|_{L_D(\mathbf{w})} = \hat{\alpha}'\beta_1$ and $\phi'(w_j)N_{a_j} = \phi(w_j)N_{a_j} = N_{a_j}$, for all j . Hence (ϕ', α') is a solution to W_1 .

Conversely, given a solution (ϕ', α') to W_1 we have, since $\mathcal{L}_1$ implies that $g = 0$, that $l(a)\alpha'(\lambda) = \alpha'(f) - k = ql(a)$ so $\hat{\alpha}'\beta_1(d)N_{a_i} = \hat{\alpha}'\beta(d)N_{a_i}$. Therefore, if we define $\phi : F(L(\mathbf{w})) \rightarrow H$ by $\phi|_{L(\mathbf{w}) \setminus \{d\}} = \phi'|_{L(\mathbf{w}) \setminus \{d\}}$ and $\phi(d) = \hat{\alpha}'\beta(d)$ it follows, since $\phi(w_j)N_{a_j} = \phi'(w_j)N_{a_j}$, for all w_j , that (ϕ, α') is a solution to W . Hence $R = \{W_0, W_1\}$ may be effectively constructed, given W , and is a resolution of W . Note that although a_1 is relator-reduced it may have a cyclic permutation which is relator-reducible. However $l(a_1) < l(a)$ so $\text{crr}(W_1) < \text{crr}(W)$, in any case.

Now suppose $\beta(d) = p_0(a, f)p_1$, $\mathcal{L}$ implies that $f \equiv k \pmod{l(a)}$, a is relator-reduced but that $a = a'a''$ and the cyclic permutation $a''a'$ of a is relator-reducible. Let $a''a'$ have relator-reduction a_0 and let $a_0 = b^{-1}a_1b$, where a_1 is cyclically reduced. Define the equations $W_i = (\mathbf{w} = 1, \beta_i, \mathcal{L}_i)$, for $i = 0, 2$ and 3 as follows. Let W_0 be as above and, for $i = 2$ and 3 , let $\beta_i|_{L_D(\mathbf{w}) \setminus \{d\}} = \beta|_{L_D(\mathbf{w}) \setminus \{d\}}$. Define

$$\beta_2(d) = p_0(a_2, l(a_2))p_1,$$

where a_2 is the relator-reduction of $a \wedge (l(a) + k)$. Define

$$\beta_3(d) = p_0(d_0, l(d_0))(a_1, \lambda l(a_1))(d_1, l(d_1))p_1,$$

where $\lambda \in \Lambda$ but λ does not occur in W , d_0 is the relator-reduction of $a'b^{-1}$ and d_1 is the relator-reduction of $ba''\iota(a, k)$. Let $\mathcal{L}_2 = \mathcal{L} \cup \{f - l(a) + k = 0\}$ and $\mathcal{L}_3 = \mathcal{L} \cup \{g = 0, \lambda > 0\}$, where $g = f - \lambda l(a) - (k + l(a))$. Let R' be the set consisting of those of W_0, W_2 and W_3 that are defined. Then as in the case where a is reducible it follows that R' is a resolution of W and again $\text{crr}(W_i) < \text{crr}(W)$, for $i = 0, 2, 3$.

Note that, as W is positive, elements of R or R' are positive. We may now replace each element of R or R' by a cyclically H^Λ -irredundant, positive resolution, observing that this process introduces no new occurrences of relator-reducibility at proper exponential H -letters. After removal of any relator-reducibility which may have been introduced at degenerate exponential H -letters we obtain a resolution of W every element of which is cyclically H^Λ -irredundant, positive and has no relator-reducible degenerate exponential H -letters. Furthermore elements W' of this resolution satisfy $\text{crr}(W') < \text{crr}(W)$ since this is true of R and R' . Continuing this process we obtain, eventually, the required resolution (when $\text{crr}(W') = 0$). As in the proof of Lemma 4.3 we may assume that W and this resolution are both in standard form.

As before, let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation with basis $(a_1, \dots, a_k)$. Let s be a cyclic permutation of an element of $\mathbf{s}_{a_i}^{\pm 1}$ such that $s = s_0^m$, where s_0 is cyclically reduced, $m \geq 1$ and $s_0 \in H_{*, a_i}$. Suppose a proper exponential letter $(s_0, f) \in H_{*, a_i}^\Lambda$ occurs in the i th component of W : that is there exists $d \in L_D(w_i)$ such that $\beta(d) = p_0(s_0, f)p_1$, for some $p_0, p_1 \in F(H_{*, a_i}^\Lambda)$. Then we say that (s_0, f) is a *relator-unconstrained* exponential H -letter occurring in $\beta(d)$, with respect to $(H_{*, a_i}, \mathbf{s}_{a_i})$. The equation W is called *relator-unconstrained* if a relator-unconstrained exponential H -letter occurs in $\beta(d)$, for some $d \in L_D(\mathbf{w})$ and *relator-constrained* otherwise.

Lemma 4.8. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained, quadratic exponential equation. Then there exists an algorithm which, given W , outputs a relator-constrained, relator-reduced, cyclically H^Λ -irredundant, positive resolution of W in standard form.*

Proof. Given Lemma 4.7 we may assume W is relator-reduced, cyclically H^Λ -irredundant and positive. Suppose that a relator-unconstrained exponential H -letter (s_0, f) occurs in W , say $\beta(d) = p_0(s_0, f)p_1$, for some $d \in L_D(w_u)$, where $s = s_0^m$ is a cyclic permutation of an element of $\mathbf{s}_{a_u}^{\pm 1}$, $m \geq 1$, $1 \leq u \leq k$.

Let $\{j_1, \dots, j_p\}$ be the set of integers such that $0 < j_i < l(s)/2$ and $\mathcal{L}_i = \mathcal{L} \cup \{f \equiv -j_i \pmod{l(s)}\}$ is consistent, for $i = 1, \dots, p$. Let $\{j_{p+1}, \dots, j_{p+q}\}$ be the set of integers such that $0 \leq j_i \leq l(s)/2$ and $\mathcal{L}_i = \mathcal{L} \cup \{f \equiv j_i \pmod{l(s)}\}$

is consistent, for $i = p + 1, \dots, p + q$. Let $T = L_D(\mathbf{w})$ and, for $i = 1, \dots, p$, define $\beta_i : L_D(\mathbf{w}) \rightarrow F(H^\Lambda)$ by $\beta_i|_{T \setminus \{d\}} = \beta|_{T \setminus \{d\}}$ and $\beta_i(d) = p_0(s_0^{-1} \wedge j_i, j_i)p_1$. For $i = p + 1, \dots, p + q$, define $\beta_i : F(H^\Lambda) \rightarrow H$ by $\beta_i|_{T \setminus \{d\}} = \beta|_{T \setminus \{d\}}$ and $\beta_i(d) = p_0(s_0 \wedge j_i, j_i)p_1$. Let $W_i = (\mathbf{w} = 1, \beta_i, \mathcal{L}_i)$, for $i = 1, \dots, p + q$ and $R = \{W_1, \dots, W_{p+q}\}$.

For all i it follows that W_i is a positive, constrained, quadratic exponential equation with smaller exponential length than W . To see that R is a resolution of W , suppose first that (ϕ, α) is a solution to W . Let $\alpha(f) = bl(s) + c$, where $0 \leq c < l(s)$, $b, c \in \mathbb{Z}$. There is either an i such that $c = j_i$, with $p + 1 \leq i \leq p + q$, or there is an i such that $c = l(s) - j_i$, with $1 \leq i \leq q$. In either case α is a solution to $\mathcal{L}_i$. Fix such an i and define $\phi' : F(L(\mathbf{w})) \rightarrow H$ by $\phi'|_{L(\mathbf{w}) \setminus \{d\}} = \phi|_{L(\mathbf{w}) \setminus \{d\}}$ and $\phi'(d) = \hat{\alpha}\beta_i(d)$. Assume first that $p + 1 \leq i \leq p + q$ so that $0 \leq j_i \leq l(s)/2$. Then

$$\begin{aligned} \phi'(d) &= \hat{\alpha}(p_0)(s_0 \wedge j_i)\hat{\alpha}(p_1) \\ &= \hat{\alpha}(p_0)s^{-b}(s_0 \wedge \alpha(f))\hat{\alpha}(p_1). \end{aligned}$$

Thus $\phi'(d)N_{a_u} = \phi(d)N_{a_u}$ and (ϕ', α) is a solution to W_i . Similarly, if $1 \leq i \leq q$, so that $0 < j_i < l(s)/2$, then

$$\begin{aligned} \phi'(d) &= \hat{\alpha}(p_0)(s_0^{-1} \wedge j_i)\hat{\alpha}(p_1) \\ &= \hat{\alpha}(p_0)s^{-b-1}s^{b+1}(s_0^{-1} \wedge (l(s) - c))\hat{\alpha}(p_1) \\ &= \hat{\alpha}(p_0)s^{-b-1}(s_0 \wedge (l(s)b + c))\hat{\alpha}(p_1) \\ &= \hat{\alpha}(p_0)s^{-b-1}(s_0 \wedge \alpha(f))\hat{\alpha}(p_1). \end{aligned}$$

Thus $\phi'(d)N_{a_u} = \phi(d)N_{a_u}$ and (ϕ', α) is a solution to W_i .

Conversely, if (ϕ', α') is a solution to $W_i \in R$ then α' is a solution to $\mathcal{L}$, as $\mathcal{L} \subset \mathcal{L}_i$. Define $\phi : F(L(\mathbf{w})) \rightarrow H$ by $\phi|_{L(\mathbf{w}) \setminus \{d\}} = \phi'|_{L(\mathbf{w}) \setminus \{d\}}$ and $\phi(d) = \hat{\alpha}'\beta(d) = \hat{\alpha}'(p_0)(s_0 \wedge \alpha'(f))\hat{\alpha}'(p_1)$. If $1 + p \leq i \leq p + q$ then $\alpha'(f) = bl(s) + j_i$, for some $b \in \mathbb{Z}$, so as above it follows that $\phi'(d)N_{a_u} = \phi(d)N_{a_u}$ and that (ϕ, α') is a solution to W . Similarly, if $1 \leq i \leq q$ then $\alpha'(f) = bl(s) - j_i$, for some $b \in \mathbb{Z}$, so

$$\begin{aligned} \phi'(d) &= \hat{\alpha}'(p_0)(s_0^{-1} \wedge j_i)\hat{\alpha}'(p_1) \\ &= \hat{\alpha}'(p_0)s^{-b}(s_0 \wedge \alpha'(f))\hat{\alpha}'(p_1), \end{aligned}$$

and (ϕ, α') is a solution to W .

Therefore R is a resolution of W . Using Lemma 4.7, we may replace R by a resolution which is relator-reduced, cyclically H^Λ -irredundant, positive and in standard form. None of the operations involved increase exponential length and so repetition of this process gives the required resolution.

Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation. Then W is called *singular* if $\beta(d) = 1 \in F(H^\Lambda)$, for some $d \in L_D(\mathbf{w})$ and *non-singular* otherwise. The set $S(W) = \{d \in L_D(\mathbf{w}) : \beta(d) = 1\}$ is called the set of *singularities* of W .

Lemma 4.9. *Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained, quadratic exponential equation. Then there exists an algorithm which, given W , outputs a non-singular, relator-constrained, relator-reduced, cyclically H^Λ -irredundant, positive resolution of W in standard form.*

Proof. We may assume that W is in standard form so that $\mathbf{w} = \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p})$, where $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ is some positive 3-partition of $(n, t, p) \in \mathbb{Z}^3$. With the above notation assume $S(W) = \{d_{i_1}, \dots, d_{i_s}\}$, where $0 < i_1 < \dots < i_s$. Define $i_0 = 0$, $i_{s+1} = n + 2t + p + 1$ and $T = \{i_1, \dots, i_s\}$ and define a map

$$\sigma : \{1, \dots, n + 2t + p\} \setminus T \longrightarrow \{1, \dots, n + 2t + p - s\},$$

by

$$\sigma(i) = i - j, \text{ when } i_j < i < i_{j+1}.$$

Suppose $\mathbf{w} = w_1, \dots, w_k$, with $w_j = q(\xi_j, n_j, t_j, p_j)$, for $j = 1, \dots, k$. Let $S_j = T \cap \{\xi_j + 1, \dots, \xi_j + n_j\}$, for $j = 1, \dots, k$. Form a new system of equations $W' = (\mathbf{w}' = 1, \beta', \mathcal{L})$ from W as follows. If $n_j - |S_j| = t_j = p_j = 0$ then delete w_j from the system. Otherwise replace w_j with a new equation w'_j formed by

- (i) deleting $x_i^{-1}d_i x_i$, from w_j when $i \in S_j$,
- (ii) replacing $x_i^{-1}d_i x_i$ with $x_{\sigma(i)}^{-1}d_{\sigma(i)}x_{\sigma(i)}$, when $i \notin S_j$,
- (iii) replacing $[x_i, x_{i+t_j}]$ with $[x_{\sigma(i)}, x_{\sigma(i+t_j)}]$, when $\xi_j + n_j + 1 \leq i \leq \xi_j + n_j + t_j$,
- (iv) replacing x_i^2 with $x_{\sigma(i)}^2$, when $\xi_j + n_j + 2t_j \leq i \leq \xi_j + n_j + 2t_j + p_j$.

Define $\beta' : L_D(\mathbf{w}') \longrightarrow F(H^\Lambda)$ by

$$\beta'(d_i) = \beta(d_{\sigma^{-1}(i)}).$$

Now suppose that (ϕ, α) is a solution to W . Define

$$\phi'(a_i) = \phi(a_{\sigma^{-1}(i)}),$$

for $a_i = x_i \in L_X(\mathbf{w})$ and $a_i = d_i \in L_D(\mathbf{w})$. By construction (ϕ', α) is then a solution to W' . Conversely, let (ϕ', α) be a solution to W' . For $i \notin T$, define $\phi(a_i) = \phi'(a_{\sigma(i)})$, where $a_i = x_i \in L_X(\mathbf{w})$ or $a_i = d_i \in L_D(\mathbf{w})$. Define $\phi(d_i) = \phi(x_i) = 1$, for $i \in T$. Then (ϕ, α) is a solution to W . Hence $\{W'\}$ is a resolution of W .

Given Lemma 4.8 we may assume that W , and so also W' , is relator-constrained, relator-reduced, cyclically H^Λ -irredundant and positive.

The process of forming the resolution of Lemma 4.9 is called *reduction of singularities*.

A parameter system $\mathcal{L}$ is said to be *normalised* if $\mathcal{L} = \mathcal{C} \dot{\cup} \mathcal{E} \dot{\cup} \mathcal{I}$, where:

- P1:** $\mathcal{C}$ is a finite set of congruences of the form $f \equiv 0 \pmod{m}$, where $f \in M$ and $m > 0$, $m \in \mathbb{Z}$;
- P2:** $\mathcal{E}$ is a finite set of diophantine equations of the form $g = 0$, $g \in M$;
- P3:** $\mathcal{I}$ is a finite set of diophantine inequalities of the form $h \geq 0$, for $h \in M$;
- P4:** if $c \in \mathbb{Z}$ and c occurs in a congruence of $\mathcal{C}$ with modulus m then $0 \leq c < m$.

Henceforward we shall assume that if $(\mathbf{w} = 1, \beta, \mathcal{L})$ is a constrained quadratic exponential equation, then $\mathcal{L}$ is normalised.

A resolution of a constrained, quadratic exponential equation which is in standard form, cyclically H^Λ -irredundant, positive, relator-reduced, relator-constrained, non-singular and normalised is called *special*.

Theorem 4.10. *Let $W = (\mathbf{w} = 1, \gamma, \mathcal{L})$ be a constrained, quadratic exponential equation with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$. Then there exists an algorithm which, given W outputs a special resolution of W .*

Now let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition and $\mathbf{z} \in (\bigcup_{k \geq 1} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$. If $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ is special we call $\mathbf{z}$ a *special* element of $(\bigcup_{k \geq 1} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$.

Lemma 4.11. *If $\mathbf{z}$ is special then the homogeneous equation*

$$Q_{\mathcal{H}} = Q(\bar{\mathbf{z}}, \mathcal{H}, \mathbf{n}, \mathbf{t}, \mathbf{p})$$

associated to $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ is special.

5. DECISION PROBLEMS

Let I be a recursive set and, for all $i \in I$, let C_i be a recursive set and R_i a recursively enumerable subset of $F_i = F(C_i)$. Let H_i be the group with presentation $\langle C_i | R_i \rangle$, for all $i \in I$. Let $K = I$ and set $X_k = \{k\}$, for all $k \in K$. Then $F_{*,i} = F_i$, for all $i \in I$.

The *quadratic exponential equation problem* or *QE-problem* in the indexed set $(\langle C_i | R_i \rangle : i \in I)$ is that of determining whether or not, given a constrained quadratic equation $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ with environment $((\{i\})_{i \in I}; (F_i)_{i \in I}; (H_i)_{i \in I})^\Lambda$, the equation W has a solution and, if so, of finding one. In fact, given that the word problem is solvable in H_i , for all i , once the existence of a solution to W has been established then a solution (ϕ, α) may be found by enumeration of all possible values of $\phi(x_j)$ and $\alpha(\lambda_j)$, for x_j and λ_j occurring in W . However the algorithms described here for the QE-problem will determine whether a solution exists by constructing one. A similar comment applies to all the other decision problems considered herein.

Since we wish to consider the extension of solvability from indexed families of groups such as $(H_i)_{i \in I}$ to free-products and one-relator products of such groups we reformulate this problem in terms of free-products. Take I , C_i , R_i , F_i and H_i to be defined as at the beginning of this section. As in Section 3 let K be a recursive set and $X_k \subset I$ be a recursive subset of I , for all $k \in K$. Define $H_{*,k}$ as in (3.1), let $\mathbf{s}_k$ be a recursive subset of $H_{*,k}$, all elements of which have length at least 2, and define N_k and G_k as in (3.2). The groups $H_{*,k}$ and G_k are equipped with natural presentations as defined in Section 3.

The QE-problem in the indexed set $(G_k : k \in K)$ is to determine, given a constrained quadratic equation W with environment

$$((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda,$$

whether or not W has a solution and, if so, to find one. The *special quadratic exponential equation problem* or *SQE-problem* in $(G_k : k \in K)$ is defined in the same way as the QE-problem with the additional condition that W is special.

We say that the QE-problem in $(G_k : k \in K)$ is *solvable* if there is an algorithm which, given input such a constrained quadratic exponential equation outputs a solution if one exists and otherwise indicates the absence of such a solution. *Solvability* of the SQE-problem in $(G_k : k \in K)$ is defined analogously.

Now let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition. In the notation above the *exponential genus problem* at $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$, or *GE-h, n, t, p-problem*, in $(G_k : k \in K)$ is that of determining whether or not, given a consistent parameter system $\mathcal{L}$ and an n -tuple $\mathbf{z}$ of elements of $(\bigcup_{k \geq 1} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$, the triple $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}\text{-genus}(\mathbf{z})$ and if so of finding a solution to $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. The *special exponential genus problem* at $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$, or *SGE-h, n, t, p-problem*, in $(G_k : k \in K)$ is defined in the same way as the exponential genus problem with the additional constraint that $\mathbf{z}$ must be special. Note that these are decision problems for presentations of the groups

G_k given by some fixed environment. In particular the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in $((C_i | R_i), i \in I)$ is to determine whether or not, given an n -tuple $\mathbf{z}$ of elements of $(\dot{\cup}_{i \in I} F_i^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$, the triple $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}\text{-genus}(\mathbf{z})$ and if so to find a solution to $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$, an equation with environment $((\{i\})_{i \in I}; (F_i)_{i \in I}; (H_i)_{i \in I})^\Lambda$.

The $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in $(G_k : k \in K)$ is *solvable* if there exists an algorithm which, given a consistent parameter system $\mathcal{L}$ and an n -tuple $\mathbf{z}$ of elements of $(\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$, decides whether or not $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}\text{-genus}(\mathbf{z})$ and if so outputs a solution to $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. The $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is *solvable* if the obvious analogous conditions are satisfied.

Example 5.1. Let $I = K = X_1 = \{1\}$, let $C_1 = \{c_1, \dots, c_n\}$ and let $R_1 = \emptyset$. Then $H_1 = F_1$, the free group of rank n . The $\mathcal{QE}$ -problem in F_1 is to determine, given a quadratic exponential equation with environment $(X_1; F_1; F_1)^\Lambda$, whether or not the equation has solution and if so of finding one. This problem is shown to be solvable by Comerford and Edmunds in [5].

Example 5.2. In [23] Lipschutz and Miller describe a family of decision problems several of which may be given in terms of quadratic exponential equations. We repeat the descriptions of [23] here for such cases. Given any two of these problems Lipschutz and Miller construct a group presentation in which the first problem is solvable and the second is not.

- (i) The *order problem* is to determine the order of a given group element. With the notation of Example 3.7 we see (loc. cit. (i)) that if the $\mathcal{GE}((1), (1), (0), (0))$ -problem is solvable in G_1 then we can determine whether or not the element $h_1 \in H_{*,1}$ has finite order in G_1 . If so then we obtain a non-zero integer n such that $h_1^n = 1$ in G_1 . Define equations

$$W_j = ((w) = 1, \beta_j, \mathcal{L}_0), \text{ for } j = 1, \dots, n,$$

where $w = d_1$, $\beta_j(d_1) = (h_1^j, l(h_1^j))$ and $\mathcal{L}_0 = \emptyset$. The order of h_1 is the smallest j such that W_j has a solution. The W_j 's are instances of the $\mathcal{GE}((0), (1), (0), (0))$ -problem in G_1 and so are also instances of the $\mathcal{GE}((0), (1), (0), (0))$ -problem. Hence if the $\mathcal{GE}((0), (1), (0), (0))$ -problem is solvable in G_1 we may solve the order problem in G_1 .

- (ii) The *power problem* is to determine, given two elements of a group, whether or not the first is a power of the second. Again Example 3.7(i) shows that we can solve this problem in G_1 if the $\mathcal{GE}((0), (1), (0), (0))$ -problem is solvable in G_1 .
- (iii) The *generalised power problem* is to determine, given two elements of a group, whether or not a power of the first is a power of the second. Example 3.7(ii) shows that we can solve this problem in G_1 if the $\mathcal{GE}((2), (1), (0), (0))$ -problem is solvable in G_1 .

Example 5.3. Continuing the development of the previous example, the *power conjugacy problem* is to determine, given two elements of a group, whether or not a power of the first is conjugate to a power of the second. From Example 3.7(iii) it follows that that we can solve this problem in G_1 if the $\mathcal{GE}((2), (2), (0), (0))$ -problem is solvable in G_1 .

Now let $\mathcal{F}$ be a collection of recursive group presentations (elements of which we'll refer to as groups). The $\mathcal{QE}$ -problem in $\mathcal{F}$ is the union of the $\mathcal{QE}$ -problem over all environments $((\{i\})_{i \in I}; F(C_i)_{i \in I}; (H_i)_{i \in I})^\Lambda$, where I is a recursive set, $((C_i :$

$R_i))_{i \in I}$ is a family of presentations indexed by I , such that $\langle C_i : R_i \rangle$ is an element of $\mathcal{F}$ and a presentation for H_i , for all $i \in I$. The $\mathcal{SQE}$ -problem, the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem, the $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem, the $\mathcal{GE}$ -problem and the $\mathcal{GE}$ -problem in $\mathcal{F}$ are all defined analogously.

Given the collection $\mathcal{F}$ define $\mathcal{F}^*$ to be the collection of groups of the form $*_{i \in I} H_i$, where I is a recursive set and H_i belongs to $\mathcal{F}$, for all $i \in I$. Each element of $\mathcal{F}^*$ is given by its natural presentation. Note that every element of $\mathcal{F}$ belongs to $\mathcal{F}^*$. Next define $\mathcal{F}^*_\sim$ to be the collection consisting of all groups of $\mathcal{F}^*$ together with groups of the form H/N , where $H = *_{i \in I} H_i$ is an element of $\mathcal{F}^*$, as above, with I a set of size at least 2, and N is the normal closure of a recursively enumerable subset of elements, of length at least 2, of H . Again each element of $\mathcal{F}^*_\sim$ is given by its natural presentation.

Given a subcollection $\mathcal{C}$ of groups of $\mathcal{F}^*_\sim$ the $\mathcal{QE}$ -problem in $\mathcal{C}$ can therefore be described as follows. Let I and K be a recursive sets such that $\langle C_i | R_i \rangle$ is a presentation of a group H_i of $\mathcal{F}$, for all $i \in I$. Let X_k be a recursive subset of I and $H_{*,k}$ be defined by (3.1), for each $k \in K$. Let $\mathbf{s}_k$ be a recursively enumerable subset of elements, of length at least 2, of $H_{*,k}$ and let N_k and G_k be defined by (3.2), for each $k \in K$. Finally assume that these sets have been chosen so that G_k belongs to $\mathcal{C}$, for all $k \in K$. The $\mathcal{QE}$ -problem in $\mathcal{C}$ is to determine, given such data and a constrained quadratic exponential equation W with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$, whether or not W has a solution and, if so, to find one. Similarly the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in $\mathcal{C}$ is to determine, given such data, a consistent parameter system $\mathcal{L}$ and an element $\mathbf{z} \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ whether or not $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}\text{-genus}(\mathbf{z})$ and, if so, to find a solution to $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. The $\mathcal{SQE}$ and $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problems in $\mathcal{C}$ have similar descriptions, differing only in that W and $\mathbf{z}$ must be special. The $\mathcal{GE}$ and $\mathcal{SGE}$ -problems also have similar descriptions.

Theorem 5.4. *Let $\mathcal{F}$ be a collection of groups with solvable word problem and let $\mathcal{C}$ be a subcollection of $\mathcal{F}^*_\sim$. The $\mathcal{QE}$ -problem in $\mathcal{C}$ is solvable if and only if the $\mathcal{SQE}$ -problem in $\mathcal{C}$ is solvable.*

Proof. If the $\mathcal{QE}$ -problem is solvable then the $\mathcal{SQE}$ -problem is obviously also solvable. Conversely suppose the $\mathcal{SQE}$ -problem in $\mathcal{C}$ is solvable. Let $W = (\mathbf{w} = 1, \beta, \mathcal{L})$ be a constrained quadratic exponential equation. Using the algorithm of Theorem 4.10 we obtain a special resolution R of W . As the $\mathcal{SQE}$ -problem is solvable we may determine, for each element $Y \in R$ whether or not a solution to Y exists. If there is such a solution then, by definition of resolution, we have an algorithm which outputs a solution of W . Otherwise W has no solution.

Lemma 5.5. *Let $\mathcal{F}$ be a collection of groups and let $\mathcal{C}$ be a subcollection of $\mathcal{F}^*_\sim$. Assume all the groups in $\mathcal{F}$ have solvable word problem and let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition of (h, n, t, p) . Then the following are equivalent.*

- (i) *The $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in $\mathcal{C}$ is solvable.*
- (ii) *The $\mathcal{SGE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem in $\mathcal{C}$ is solvable, for all positive 4-partitions $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ of (h, n', t, p) such that $n' \in \mathbb{Z}$ and $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.*
- (iii) *The $\mathcal{SGE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem in $\mathcal{C}$ is solvable, for all positive 4-partitions $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq_g (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.*

Proof. For the duration of the proof choose recursive sets I and K such that $\langle C_i | R_i \rangle$ is a presentation of a group H_i of $\mathcal{F}$, for all $i \in I$. Choose X_k to be a recursive subset of I and let $H_{*,k}$ be defined by (3.1), for each $k \in K$. Choose a recursively enumerable set $\mathbf{s}_k$ of $H_{*,k}$ and let N_k and G_k be defined by (3.2), for each $k \in K$. Finally assume that these sets have been chosen so that G_k belongs to $\mathcal{C}$, for all $k \in K$.

We show first that (i) implies (iii). Suppose that the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in $\mathcal{C}$ is solvable. Then the $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is obviously also solvable. Assume that $\mathbf{n} = (n_1, \dots, n_k)$ is a partition of an integer n and define $\mu_j = \mu_j(\mathbf{n})$, for $j = 1, \dots, k+1$, as in (3.9). Let $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ be a positive 4-partition with $\mathbf{h}' = (h'_1, \dots, h'_l)$, $\mathbf{n}' = (n'_1, \dots, n'_l)$, $\mathbf{t}' = (t'_1, \dots, t'_l)$, $\mathbf{p}' = (p'_1, \dots, p'_l)$ and $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq_g (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Then there exist integers $j_1, \dots, j_l$ such that $1 \leq j_1 < \dots < j_l \leq k$ with $h'_i \leq h_{j_i}$, $n'_i \leq n_{j_i}$, $t'_i \leq t_{j_i}$, and $p'_i \leq p_{j_i}$, for $i = 1, \dots, l$. Define $\mu'_j = \mu_j(\mathbf{n}')$, for $j = 1, \dots, l+1$. Let $\mathcal{L}$ be a system of parameters and let $\mathbf{z}' = (z'_1, \dots, z'_{n'})$ be an element of $(\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}', \mathbf{h}')$ with basis $(a'_1, \dots, a'_l)$. Define an element $\mathbf{z} = (z_1, \dots, z_n)$ of $(\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ as follows. For s such that $\mu_{j_i} + 1 \leq s \leq \mu_{j_i} + n'_i$ define $z_s = z'_{s'}$, where $s' = s - \mu_{j_i} + \mu'_i$. For all other $s \in \{1, \dots, n\}$ define $z_s = (1, 0) \in H^\Lambda$. Then $\mathbf{z}$ has basis $(a_1, \dots, a_k)$, where $a_{j_i} = a'_i$, and $a_j = 1$, if $j \notin \{j_1, \dots, j_l\}$.

Recalling (2.1) and (2.2) let $\xi_j = \xi_j(\mathbf{n}, \mathbf{t}, \mathbf{p})$ for $j = 1, \dots, k$, $\mathbf{q} = \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p})$, $\xi'_j = \xi'_j(\mathbf{n}', \mathbf{t}', \mathbf{p}')$ for $j = 1, \dots, l$ and $\mathbf{q}' = \mathbf{q}(\mathbf{n}', \mathbf{t}', \mathbf{p}')$. Recalling (3.10) let $\beta = \beta(\mathbf{z}, \mathbf{n}, \mathbf{t}, \mathbf{p})$, $\beta' = \beta(\mathbf{z}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$, $Q = Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p}) = (\mathbf{q} = 1, \beta, \mathcal{L})$ and $Q' = Q(\mathbf{z}', \mathcal{L}, \mathbf{n}', \mathbf{t}', \mathbf{p}') = (\mathbf{q}' = 1, \beta', \mathcal{L})$. Given a homomorphism $\phi : F(L(\mathbf{q})) \rightarrow H$ we may construct a homomorphism $\phi' : F(L(\mathbf{q}')) \rightarrow H$ as follows. For each $i \in \{1, \dots, l\}$ and s' with $1 + \xi'_i \leq s' \leq n'_i + \xi'_i$, define

$$\phi'(x_{s'}) = \phi(x_s) \text{ and } \phi'(d_{s'}) = \phi(d_s),$$

where $s = s' - \xi'_i + \xi_{j_i}$. For s' with $n'_i + \xi'_i < s' \leq n'_i + 2t'_i + p'_i + \xi'_i$, define

$$\phi'(x_{s'}) = \phi(x_s),$$

where $s = s' - \xi'_i - n'_i + \xi_{j_i} + n_{j_i}$. If (ϕ, α) is a solution to Q then (ϕ', α) is a solution to Q' .

Similarly, given a homomorphism $\phi' : F(L(\mathbf{q}')) \rightarrow H$ we may construct a homomorphism $\phi'' : F(L(\mathbf{q})) \rightarrow H$ as follows. For each $j_i \in \{j_1, \dots, j_l\}$ and s with $1 + \xi_{j_i} \leq s \leq n'_{j_i} + \xi_{j_i}$, define

$$\phi''(x_s) = \phi'(x_{s'}) \text{ and } \phi''(d_s) = \phi'(d_{s'}),$$

where $s = s' - \xi'_i + \xi_{j_i}$. For each $j_i \in \{j_1, \dots, j_l\}$ and s with $n_{j_i} + \xi_{j_i} < s \leq n_{j_i} + 2t_{j_i} + p_{j_i} + \xi_{j_i}$, define

$$\phi''(x_s) = \phi'(x_{s'}),$$

where $s = s' - \xi'_i - n'_i + \xi_{j_i} + n_{j_i}$. For all other i , in an appropriate range, define $\phi(d_i) = 1$ and $\phi(x_i) = 1$. If (ϕ', α) is a solution to Q then (ϕ'', α) is a solution to Q' . Therefore solutions of Q correspond to solutions of Q' . The algorithm for the solution of the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem can thus be used to solve the $\mathcal{SGE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem. Therefore (i) implies (iii).

Suppose now that (ii) holds, so the $\mathcal{SGE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable for all all positive 4-partitions $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ of (h, n', t, p) such that $n' \in \mathbb{Z}$ and $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Given $\mathbf{z} \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$, where $\mathcal{L}$ is some parameter system, the equation $Q = Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ is in standard form. From Lemma 4.10 it follows

that there is an algorithm which outputs a special resolution R of Q . Furthermore the only operation involved in producing R which changes any of the n_i , t_i , p_i or the length k of the partition, is reduction of singularities. Reduction of singularities never reduces t_i , p_i , t or p and never increases n_i or k . As none of the operations involved in producing R increases the exponential length of any of the equations involved it follows that if $(\mathbf{q}(\mathbf{n}', \mathbf{t}', \mathbf{p}') = 1, \mathcal{L}', \beta')$ is an element of R with boundary labels list $(z'_1, \dots, z'_{n'})$ then there is a partition $\mathbf{h}'$ of h such that $\mathbf{z}' \in (\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}', \mathbf{h}')$, $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \leq (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ and $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ is a partition of (h, n', t, p) , for some $n' \in \mathbb{Z}$. The algorithm for the $\mathcal{SGE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem may therefore be used to determine whether this element of R has a solution and if so to find one. From the definition of resolution it now follows that an algorithm for the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem exists. Therefore (ii) implies (i). Finally, it follows directly from the definition of $\leq_g$ that (iii) implies (ii).

We define the $\mathcal{GE}$ -problem and $\mathcal{SGE}$ -problem in $\mathcal{C}$ to be the unions of the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problems and the $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problems, respectively, over all positive 4-partitions $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ of length k , for all $k > 0$.

Lemma 5.6. *Let $\mathcal{F}$ be a collection of groups and let $\mathcal{C}$ be a subcollection of $\mathcal{F}_\sim^*$. Assume all the groups in $\mathcal{F}$ have solvable word problem. Then the following are equivalent.*

- (i) *The $\mathcal{QE}$ -problem in $\mathcal{C}$ is solvable.*
- (ii) *The $\mathcal{GE}$ -problem in $\mathcal{C}$ is solvable.*
- (iii) *The $\mathcal{SGE}$ -problem in $\mathcal{C}$ is solvable.*

Proof. That assertions (ii) and (iii) are equivalent follows from Lemma 5.5. Suppose the $\mathcal{QE}$ -problem is solvable. In the notation of the proof of Lemma 5.5 Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition of (h, n, t, p) , $\mathcal{L}$ a parameter system and $\mathbf{z}$ an element of $(\dot{\cup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$. The algorithm for the $\mathcal{QE}$ -problem may then be used to determine a solution to the equation $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. The $\mathcal{GE}$ -problem is therefore solvable.

Conversely suppose that the $\mathcal{GE}$ -problem is solvable. Using the notation of the proof of Lemma 5.5 again let $W = (\mathbf{w} = 1, \mathcal{L}, \beta)$ be a constrained quadratic exponential equation with environment $((X_k)_{k \in K}; (H_i)_{i \in I}; (G_k)_{k \in K})^\Lambda$. Using Theorem 4.2 we may assume that $\mathbf{w}$ is in standard form and so $\mathbf{w} = \mathbf{q}(\mathbf{n}, \mathbf{t}, \mathbf{p})$, for some $\mathbf{n}, \mathbf{t}$ and $\mathbf{p}$. Let $\mathbf{z}$ be a boundary labels list for $(\mathbf{w}, \beta)$ and let $Q = Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Then W has a solution if and only if $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}\text{-genus}(\mathbf{z})$. Therefore, if Q has exponential coordinates $\mathbf{h}$, the algorithm for the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem may be used to determine a solution to W . It follows that the $\mathcal{QE}$ -problem is solvable.

5.1. Decision problems in free products and one-relator products. We shall consider exponential genus problems for collections of groups consisting of free-products and one-relator products of groups in which such problems are already solvable.

Let $\mathcal{F}$ be a collection of group presentations and define $\mathcal{F}_\sim^*$ as in the previous section. Define the subcollection $\mathcal{C}_m$ of $\mathcal{F}_\sim^*$ to consist of $\mathcal{F}_\sim^*$ together with groups of the form H/N , where $H = *_{i \in I} H_i$, H_i belongs to $\mathcal{F}$, I is a recursive set of size at least 2 and N is the normal closure of an element r^d of H where r is cyclically reduced, not a proper power in H , $l(r) \geq 2$ and $d \geq m$. We call such groups d -torsion one-relator products with factors in $\mathcal{F}$. If H/N is a d -torsion group then

we may always write $H = A * B$, where A and B belong to $\mathcal{F}^*$ and the conditions on r still hold: that is H/N is a d -torsion group with exactly two factors in $\mathcal{F}^*$. Let s denote the element r^d of H . Then $l(s) = dl(r)$ is at least $2d$ and s is cyclically reduced as a word in H .

A d -torsion group $A * B/N$, where N is the normal closure of r^d , is said to be *exceptional of type $E(p, *, d)$* if $r = aUbU^{-1}$, with $a, b \in A \dot{\cup} B$ and $a^p = 1$ and of *type $E(p, q, d)$* if in addition $b^q = 1$. A d -torsion group may be exceptional in more than one way, involving different p, q, d : see [9] for more details. We define the collection of groups $\mathcal{C}_m^+$ to consist of $\mathcal{F}^*$ together with those groups of $\mathcal{C}_m \setminus \mathcal{F}^*$ which are not (in any way) exceptional of type $E(2, *, m)$.

Example 5.7. Let $\mathcal{F}$ be a collection of group presentations and consider the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in $\mathcal{F}$, where $\mathbf{h}$ is a partition of 0. Suppose

$$\mathbf{z} = (z_1, \dots, z_n) \in (\dot{\cup}_{i \in I} H_i^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h}),$$

for some recursive set I , groups $H_i \in \mathcal{F}$, and consistent parameter system $\mathcal{L}$. Since no parameters occur in $\mathbf{z}$, for all j , we may regard z_j an element of H_i , for some $i \in I$. The parameter system $\mathcal{L}$ has no bearing on the solutions to the equation associated to $\mathbf{z}$ and we can replace it by the empty set. We write *genus* instead of $\emptyset$ -genus. If $\mathbf{n} = (n_1, \dots, n_k)$, $\mathbf{t} = (t_1, \dots, t_k)$ and $\mathbf{p} = (p_1, \dots, p_k)$ then

$$(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \text{genus}(\mathbf{z}) \text{ if and only if } (n_i, t_i, p_i) \in \text{genus}(z_i), \text{ for } i = 1, \dots, k.$$

Therefore the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem reduces to the union of the

$$\mathcal{GE}((0), (n_i), (t_i), (p_i))\text{-problems.}$$

That is, we need not consider systems of equations of size greater than 1 if $\mathbf{h}$ is a partition of zero. When $\mathbf{h}$ is a partition of zero we refer to the $\mathcal{QE}$ -problem as the $\mathcal{Q}$ -problem. It is known (see [4], [6], [13], [14]) that the $\mathcal{Q}$ -problem is solvable in free groups and that if the $\mathcal{Q}$ -problem is solvable in $\mathcal{F}$ then the $\mathcal{Q}$ -problem is solvable in $\mathcal{F}^*$.

Assume now that n, t, p are non-negative integers such that $n + t + p > 0$ and that the $\mathcal{GE}((0), (n'), (t'), (p'))$ -problem is solvable in $\mathcal{F}$, for all n', t', p' such that

$$(5.1) \quad \begin{aligned} t' + p'/2 &\leq t + p/2 \text{ and} \\ n' &\leq n + 2(t - t') + (p - p'). \end{aligned}$$

Now let m be a fixed integer and let $\mathcal{C}$ be a collection of one-relator products with factors in $\mathcal{F}$ such that every element of $\mathcal{C}$ is d -torsion, for some $d \geq m$. It is known ([7], [8], [9]) that if (5.1) and one of the following conditions (i)–(iii) holds, for all groups G of $\mathcal{C}$, then the $\mathcal{GE}((0), (n), (t), (p))$ -problem is solvable in $\mathcal{C}$.

- (i) $m \geq 5$ and G is not in any way exceptional of type $E(2, 3, 5)$ or $E(2, 3, 6)$.
- (ii) $m \geq 4$ and no letter occurring in r has order 2.
- (iii) $m \geq 2$ and the groups of $\mathcal{F}$ are locally indicable.

The previous example motivates the following definition. Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition of length k of (h, n, t, p) , for some $h, n, t, p \in \mathbb{Z}$. Define $\mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ to be the set of positive 4-partitions $(\mathbf{h}', \mathbf{n}', \mathbf{p}', \mathbf{t}')$, of length k' , satisfying the conditions (i)–(iii) below. Assume $\mathbf{h}' = (h'_1, \dots, h'_{k'})$, $\mathbf{n}' = (n'_1, \dots, n'_{k'})$, $\mathbf{t}' = (t'_1, \dots, t'_{k'})$ and $\mathbf{p}' = (p'_1, \dots, p'_{k'})$.

- (i) There exists a partition $\mathbf{l} = (l_1, \dots, l_k)$ such that $0 \leq l_j \leq \max\{1, h_j\}$, for $j = 1, \dots, k$ and

$$k' = \sum_{j=1}^k l_j.$$

- (ii) Setting $q_0 = 0$ and $q_j = \sum_{s=1}^j l_s$, for $j = 1, \dots, k$, we have

$$\sum_{i=q_{j-1}+1}^{q_j} h'_i \leq h_j, \text{ for } j = 1, \dots, k.$$

- (iii) For $q_{j-1} + 1 \leq i \leq q_j$ and $1 \leq j \leq k$
 (a) $t'_i + p'_i/2 \leq t_j + p_j/2$ and
 (b) $n'_i \leq n_j + 2(t_j - t'_i) + (p_j - p'_i)$.

The main result proved here is the following.

Theorem 5.8. *Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition and let $\mathcal{F}$ be a collection of group presentations with solvable $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem, for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Then the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in the subcollection $\mathcal{C}_6^+$ of $\mathcal{F}^*$, for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.*

Most of the remainder of the paper is given to proving a restricted version of this theorem. Here we state the restricted version and show how Theorem 5.8 follows from it.

Theorem 5.9. *Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, let I be a recursive set and let $A, B, H_i, i \in I$, groups, with fixed presentations given, such that the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in the indexed set of groups $(A, B) \cup (H_i : i \in I)$, for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Let $H = A * B$ and let $s = r^m$, where $r \in H$ is cyclically reduced, not a proper power in H , $l(r) \geq 2$ and $m \geq 6$. Set $G = H$ or $G = H/N$, where N is the normal closure of s in H . In the latter case assume further that G is not in any way exceptional of type $E(2, *, m)$. Then the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in $(G, A, B) \cup (H_i : i \in I)$, for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.*

Proof of Theorem 5.8. Assume that Theorem 5.9 holds and set $\mathcal{N} = \mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Suppose that $\mathcal{C}$ is a subcollection of the collection $\mathcal{F}^*$ for which it has been shown that if the hypotheses of Theorem 5.8 hold then the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is solvable in $\mathcal{C}$. Given any $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}$ we have $\mathcal{N}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \subseteq \mathcal{N}$, so the hypotheses of Theorem 5.8 hold for $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ instead of $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Hence the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in $\mathcal{C}$. In particular it suffices to show that the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is solvable in $\mathcal{C}_6^+$.

Assume then that the hypotheses of Theorem 5.8 hold. First we shall prove the following. Let l be a positive integer and $(A_1, \dots, A_l, B_1, \dots, B_l) \cup (H_i : i \in I)$ be an indexed set of groups all of which belong to $\mathcal{F}$. For $i = 1, \dots, l$ let $H_{*,i} = A_i * B_i$ and let N_i be a subgroup of $H_{*,i}$ satisfying either

01. $N_i = 1 \in H_{*,i}$ or
 02. N_i is the normal closure of $r_i^{m_i}$ in $H_{*,i}$, where $r_i \in H_{*,i}$ and $m_i \geq 6$ is an integer such that $H_{*,i}/N_i$ is an m_i -torsion one-relator product which is not exceptional of type $E(2, *, m)$.

Set $G_i = H_{*,i}/N_i$, $i = 1, \dots, l$. Then the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in

$$(G_1, \dots, G_l, A_1, \dots, A_l, B_1, \dots, B_l) \cup (H_i : i \in I),$$

for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}$.

The case $l = 1$ follows from Theorem 5.9. Assume inductively that the result holds for some $l - 1 \geq 1$. Then the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in

$$(G_1, \dots, G_{l-1}, A_1, \dots, A_l, B_1, \dots, B_l) \cup (H_i : i \in I),$$

for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}$. Using Theorem 5.9 again the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in

$$(G_1, \dots, G_l, A_1, \dots, A_l, B_1, \dots, B_l) \cup (H_i : i \in I),$$

as required.

Now let $l_1, \dots, l_n$ be integers, $l_i \geq 2$ and let $A_{i,1}, \dots, A_{i,l_i}$ belong to $\mathcal{F}$, for $i = 1, \dots, n$. Define $H_{*,i} = * \{A_{i,j} : j = 1, \dots, l_i\}$ and let N_i be a subgroup of $H_{*,i}$ satisfying $\mathcal{O}1$ or $\mathcal{O}2$, for $i = 1, \dots, n$. We shall show that, if $G_i = H_{*,i}/N_i$, then the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in

$$(G_1, \dots, G_l, A_{1,1}, \dots, A_{n,l_n}) \cup (B_i : i \in I),$$

for any recursive set I and groups B_i of $\mathcal{F}$, and for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}$.

The case $l_i = 2$ for all i is covered by the argument above. Using induction on $\sum_{i=1}^n (l_i - 2)$ we assume that $l_i > 2$, for some i . Without loss of generality we may assume that $i = 1$ and set

$$H'_{*,1} = A_{1,2} * \dots * A_{1,l_1}.$$

The inductive hypothesis implies that the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in

$$(H'_{*,1}, G_2, \dots, G_n, A_{1,1}, \dots, A_{n,l_n}) \cup (B_i : i \in I),$$

for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}$. As $G_1 = (A_{1,1} * H'_{*,1})/N_1$, it follows from Theorem 5.9 that the $\mathcal{GE}(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}')$ -problem is solvable in

$$(G_1, H'_{*,1}, G_2, \dots, G_n, A_{1,1}, \dots, A_{n,l_n}) \cup (B_i : i \in I)$$

and so in particular in

$$(G_1, G_2, \dots, G_n, A_{1,1}, \dots, A_{n,l_n}) \cup (B_i : i \in I),$$

for all $(\mathbf{h}', \mathbf{n}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}$, as required.

Now let R be a recursive set and let $\mathcal{I}$ be a set of groups, indexed by R , all of which belong to $\mathcal{C}_6^+$. Then we can write $R = J \cup K$, where J and K are recursive sets, and there exists an element B_j of $\mathcal{F}$, for all $j \in J$, and G_k of $\mathcal{C}_6^+$ but not to $\mathcal{F}$, for all $k \in K$, such that $\mathcal{I} = (B_j)_{j \in J} \cup G_{k \in K}$. Thus there exist recursive sets I , $X_k \subseteq I$ and groups H_i , $H_{*,k}$ and N_k , such that $H_{*,k} = *_{i \in X_k} H_i$, with $k \in K$, $H_i \in \mathcal{F}$, $G_k = H_{*,k}/N_k$, where $H_{*,k}$, N_k and G_k satisfy $\mathcal{O}1$ and $\mathcal{O}2$. Suppose that

$$\mathbf{z} \in \left(\bigcup_{k \in K} H_{*,k}^\Lambda \dot{\cup} \bigcup_{j \in J} B_j^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h} \right).$$

The equation Q associated to $\mathbf{z}$ has environment

$$(5.2) \quad ((X_k)_{k \in K}, (\{j\})_{j \in J}; (H_i)_{i \in I}, (B_j)_{j \in J}; (G_k)_{k \in K}, (B_j)_{j \in J})^\Lambda$$

and we may assume, after reordering if necessary, that $\mathbf{z}$ and Q have basis $(k_1, \dots, k_l, j_1, \dots, j_{l'})$, where $k_i \in K$ and $j_i \in J$. From Lemma 3.1 a solution to Q with environment (5.2) determines a solution to Q with environment

$$(5.3) \quad \begin{aligned} & (X'_{k_1}, \dots, X'_{k_l}, \{j_1\} \dots, \{j_{l'}\}; \\ & (H_i)_{i \in I'}, B_{j_1}, \dots, B_{j_{l'}}; \\ & G'_{k_1}, \dots, G'_{k_l}, B_{j_1}, \dots, B_{j_{l'}})^\Lambda \end{aligned}$$

and the same basis, where $\text{supp}(Q) = (X'_{k_1}, \dots, X'_{k_l}, \{j_1\} \dots, \{j_{l'}\})$, $I' = X'_{k_1} \cup \dots \cup X'_{k_l}$ and G'_k is defined in (3.8). Conversely a solution to Q with environment (5.3) is a solution to Q with environment (5.2). By construction $G'_{k_i} = H'_{*,k_i}/N'_i$, where $H'_{*,i}$, N'_i and G'_i , satisfy $\mathcal{O}1$ and $\mathcal{O}2$. Hence $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}\text{-genus}(\mathbf{z})$ if and only if there exists a solution to Q with environment (5.3). We may determine whether or not Q has a solution with this environment if we can solve the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem in

$$(5.4) \quad (G'_{k_1}, \dots, G'_{k_l}) \cup (H_i : i \in I') \cup (B_{j_1}, \dots, B_{j_{l'}}).$$

Since N'_{k_s} is either trivial or the normal closure of a single element, it follows that X'_{k_s} is finite, for $s = 1, \dots, l$. Hence, from the previous case, the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is solvable in (5.4) and the Theorem follows.

Corollary 5.10. *Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition and let $\mathcal{F}$ be a collection of group presentations with solvable $\mathcal{QE}$ -problem. Then the $\mathcal{QE}$ -problem is solvable in the subcollection $\mathcal{C}_6^+$ of $\mathcal{F}_\sim^*$.*

Proof. This follows from Theorem 5.8 and Lemma 5.6.

Corollary 5.11. *Let $\mathcal{F}$ be the cyclic groups. The $\mathcal{QE}$ -problem is solvable in $\mathcal{C}_6^+$. That is in free-products and d -torsion one-relator products which belong to $\mathcal{C}_6^+$, with cyclic factors. In particular the $\mathcal{QE}$ -problem is solvable in recursively generated free groups.*

Proof. The $\mathcal{QE}$ -problem is solvable in $\mathcal{F}$ since each instance reduces to a system of linear Diophantine equations, congruences and inequalities. The result follows from Corollary 5.10.

Corollary 5.12. *Let A and B be groups such that the $\mathcal{GE}$ -problem is solvable in (A, B) . Let $G = A * B / N$ be a d -torsion one-relator product of A and B , for $d \geq 6$, which is not in any way special of type $E(2, *, d)$. Then the $\mathcal{GE}$ -problem is solvable in (G, A, B) . In particular, if $z_1, \dots, z_k$ are elements of $A * B$ and $\mathcal{L}$ is a consistent parameter system in $\{\lambda_1, \dots\}$ then there is an algorithm to decide whether or not there are integers $n_1, \dots, n_k$ and elements $x_i \in A * B$ such that*

$$(i) \quad \prod_{i=1}^k x_i^{-1} z_i^{n_i} x_i \prod_{i=k+1}^{n+t} [x_i, x_{i+t}] \prod_{i=k+2t+1}^{k+2t+p} x_i^2 = 1, \text{ in } G \text{ and}$$

(ii) *there is a solution to $\mathcal{L}$ with $\lambda_i = n_i$, $i = 1, \dots, k$.*

6. PICTURES

Let A and B be groups and $m \geq 2$ an integer and set $H = A * B$. Let X'_A and X'_B be CW-complexes with $\pi_1(X'_A, x_A) = A$ and $\pi_1(X'_B, x_B) = B$, for some $x_A \in X'_A$ and $x_B \in X'_B$. Let $O = A$ or $O = B$ and let e_O be a copy of the unit interval $[0, 1]$. Form the quotient space X_O of $X'_O \dot{\cup} e_O$ by identifying x_O and $1 \in e_O$. Form the quotient space Y of $X_A \dot{\cup} X_B$ by identifying $0 \in e_A$ and $0 \in e_B$ to a point p . Then $\pi_1(Y, p) = H$.

Fix a word $r \in H$ with $l(r) \geq 2$, such that r is not a proper power in H . Let $s = r^m$, $N(s) = \langle \langle s \rangle \rangle^H$ and $G = H/N(s)$. Suppose that $s = a_1 b_1 \cdots a_n b_n$, when written in normal form for the free product. Let Δ_s be a closed unit disk with oriented boundary divided into $2n$ intervals. Label the intervals consecutively $a_1, b_1, \dots, a_n, b_n$, so that $\partial\Delta_s$, read from a suitable starting point in the direction of orientation, is labelled s . Form the quotient space Z of $Y \dot{\cup} \Delta_s$ by identifying $\partial\Delta_s$ with a path in (Y, p) representing $s \in H$, according to the label of $\partial\Delta_s$. Then $\pi_1(Z, p) = G$. Alternatively, let $G = H$ and define $\Delta_s = p \in Y$, so that $Z = Y$ and $\pi_1(Z, p) = G$ again.

Let Σ be a compact surface and let $f : \Sigma \longrightarrow Z$ be a continuous map satisfying the following properties.

- S1. f is transverse to the midpoint c of Δ_s and the closure of $f^{-1}(\text{int}(\Delta_s))$ is a regular neighbourhood of $f^{-1}(c)$.
- S2. f is transverse to p on $\Sigma_0 = \Sigma \setminus f^{-1}(\text{int}(\Delta_s))$ and $f^{-1}(e_A \cup e_B)$ is a regular neighbourhood of $f^{-1}(p)$ in Σ_0 .

Let $\Gamma = f^{-1}(\Delta_s)$. Then (f, Γ) is a *sketch* on Σ over G (with respect to Z) which we refer to merely as Γ , when the meaning is clear. A connected component of the closure of $f^{-1}(\text{int}(\Delta_s))$ is called a *vertex* of Γ and the set of all vertices is denoted $\mathcal{V}(\Gamma)$ or just $\mathcal{V}$. Condition S1 implies that $\mathcal{V}$ consists of a collection of embedded disks in $\text{int}(\Sigma)$, each of which is a regular neighbourhood of a point of $f^{-1}(c)$. If $G = H$ then $\mathcal{V} = \emptyset$. A connected component of $f^{-1}(p)$ is called an *arc* of Γ and the set of all arcs is denoted $\mathcal{A} = \mathcal{A}(\Gamma)$. Condition S2 implies that $\mathcal{A}$ consists of a set of properly embedded 1-submanifolds of Σ_0 .

A connected component of $\partial v \setminus \{\cup e : e \in \mathcal{A}\}$, where $v \in \mathcal{V}$, is called a *vertex corner* of Γ . A component of $\partial\Sigma \setminus \Gamma$ is called a *boundary corner* of Γ . The closure of a connected component of $\Sigma \setminus \Gamma$ is called a *region* of Γ and the set of regions is denoted $\mathcal{R}(\Gamma)$ or $\mathcal{R}$. Every vertex and boundary corner belongs to a unique region of Γ .

Now suppose that Σ' is a connected component of Σ and define $\Gamma' = \Sigma' \cap \Gamma$. If Σ' is orientable we may choose consistent orientations of each region and vertex of Σ' . We call such a choice an *orientation* of Γ' . If β is a boundary component of Σ' then an orientation ζ of Γ' gives rise to a unique orientation $\zeta(\beta)$ of β . Similarly the orientation of regions induces an orientation on ∂v , which we denote $\zeta(\partial v)$, for all $v \in \mathcal{V}$. Thus $\zeta(\partial v)$ is the opposite of the orientation induced on ∂v by the orientation of v . If l is a sub-interval of a corner c of Γ then there is a unique orientation $\zeta(l)$ of l induced from the orientation of the boundary component (of Σ' or $v \in \mathcal{V}$) to which c belongs.

If Σ' is non-orientable then we choose an orientation for each connected component of $\partial\Sigma'$, for each connected component of $\partial\Delta$, for all regions Δ of Γ' , and for ∂v , for all vertices v of Γ' . To each vertex v we assign the orientation opposite




Figure 1: The orientation of the boundary is shown outside and the orientation of the vertex inside: $s = abcd$.

to that chosen for ∂v . Call the collection of these orientations ζ an *orientation* of Γ' . Let l be a sub-interval of a corner c of Γ' . Then c belongs to a boundary component β of Σ' or to the boundary β of a vertex. The orientation $\zeta(\beta)$ induces an orientation of l which we call $\zeta(l)$.

We now fix an orientation for Γ' for all connected components Σ' of Σ . We call these orientations an *orientation* of Γ and denote this by ζ . We refer to the orientation induced on y by ζ as $\zeta(y)$, whenever y is an appropriate subset of Σ .

Let (f, Γ) be a sketch on Σ with orientation ζ . We define a labelling function f^l from corners of Γ to H as follows. Let c be a corner of Γ with orientation $\zeta(c)$ and let q be the closed path $f(c)$ in Y . If c meets an arc of Γ then q is a directed path based at p . In this case we define the *label* $f^l(c)$ of c read in the direction $\zeta(c)$ to be the element of H represented by the based directed path q . If c does not meet an arc of Γ then we choose a base point x on c and regard q as a directed path based at $f(x)$. Since c meets no arc of Γ , the path q lies in $Y \setminus p$ and we may choose a path α from p to x such that $\text{int}(\alpha)$ lies in X_A or X_B . Then the directed path $\alpha q \alpha^{-1}$ represents an element g of $A \cup B$. The label of c , read in the direction of orientation $\zeta(c)$, is defined to be g . If c read in the direction $\zeta(c)$ has label w then the label of c read in the direction opposite to $\zeta(c)$ is w^{-1} .

Definition 6.1. The sketch Γ endowed with the orientation ζ and the labelling function f^l is called a *picture* on Σ over G .

A picture over $G = H$ may be regarded as a picture over $G = H/N(s)$, having no vertices. Conversely a picture over $G = H/N(s)$ with no vertices may be regarded as a picture over H . If v is a vertex of Γ and x is a point of $\partial v \cap \{\cup e : e \in \mathcal{A}\}$ then the *label* of v with *base point* x is the word of H obtained by reading off the labels of corners of v in the order they occur whilst travelling once round ∂v , from x in the direction $\zeta(\partial v)$: where the label of each corner c is read in the direction of orientation induced from that of $\zeta(\partial v)$, namely $\zeta(c)$. Since $f(\partial v) = \partial \Delta_s$ we have:

P1. the label of v (with any base point) is a cyclic permutation of s^ε , where $\varepsilon = \pm 1$.

The orientation ζ on ∂v induces an orientation on $f(\partial v) = \partial \Delta_s$. If this is the same as the fixed orientation of $\partial \Delta_s$ then we may take $\varepsilon = 1$ and we say that v is a *positive* vertex (with respect to ζ). Otherwise we take $\varepsilon = -1$ and say v is a *negative* vertex (with respect to ζ), as shown in Figure 1. Similarly, if β is a

connected component of $\partial\Sigma$ or a connected component $\partial\Delta$, for some region Δ of Γ , let $x \in \beta \cap \{\cup e : e \in \mathcal{A}(\Gamma)\}$ or, if β consists of a boundary corner meeting no arc of Γ , let x be the chosen base point of this corner. Then the *label* of β with *base point* x is the word obtained by reading once round β , from x in the direction of $\zeta(\beta)$. When reference is made to the label of such β it is to be assumed that a base point has been chosen. The *boundary labels* of a region Δ are the labels of the connected components of $\partial\Delta$.

As a consequence of properties *S1* and *S2* and the definitions above a picture Γ on Σ satisfies the following properties *P2*–*P4*.

- P2.* Let Δ be a region of Γ with boundary labels $d_1, \dots, d_n$. Then either $d_i \in A$, for $i = 1, \dots, n$, in which case Δ is called an *A-region* or $d_i \in B$, for $i = 1, \dots, n$, in which case Δ is called a *B-region*.
- P3.* Let $O = A$ or B and let Δ be an *O-region*, with boundary labels $d_1, \dots, d_n$, which is homeomorphic to the connected sum of an n -punctured sphere with t torii and p projective planes. Then there exist elements $w_1, \dots, w_n, x_1, \dots, x_t, y_1, \dots, y_t, z_1, \dots, z_p \in O$ such that

$$(6.1) \quad \prod_{i=1}^n w_i^{-1} d_i w_i \prod_{i=1}^t [x_i, y_i] \prod_{i=1}^p z_i^2 = 1,$$

in O .

- P4.* If a is an arc of Γ then a separates an *A-region* from a *B-region*.

Conversely properties *P1* to *P4* characterise pictures.

Lemma 6.2. *Let Σ be a compact surface and let Γ consist of*

- (i) *a finite collection $\mathcal{V}$ of disjoint closed disks in $\text{int}(\Sigma)$ (called vertices);*
- (ii) *a finite collection $\mathcal{A}$ of disjoint connected properly embedded 1-submanifolds of $\Sigma_0 = \Sigma \setminus \cup_{v \in \mathcal{V}} \text{int}(v)$ (called arcs);*
- (iii) *regions and corners and an orientation ζ defined as in the case of the sketch above;*
- (iv) *a labelling function f^l from corners of Γ to $A \cup B$, which respects orientation, and a collection of labels defined as for pictures above.*

*If Γ satisfies *P1* to *P4* then Γ is a picture on Σ over G .*

We define the *genus* of a compact connected surface Σ , with k boundary components, to be $\text{genus}(\Sigma) = (2 - \chi(\Sigma) - k)/2$. The *genus* of a disconnected compact surface is defined to be the sum of the genera of its connected components.

An arc of a picture, which is not a closed curve, both ends of which meet $\partial\Sigma$ is called an arc of *type I*. Let Δ be a region of Γ . Then we define

$$\begin{aligned} t(\Delta) &= \text{number of arcs of } \Gamma \text{ in } \partial\Delta, \\ \beta(\Delta) &= \text{number of components of } [\partial\Sigma \setminus \{\text{arcs of } \Gamma\}] \cap \partial\Delta \\ &= \text{number of boundary corners of } \Delta, \\ \rho(\Delta) &= \text{number of components of } [\Gamma \setminus \{\text{arcs of } \Gamma\}] \cap \partial\Delta \\ &= \text{the number of vertex corners of } \Delta, \\ \gamma(\Delta) &= \text{number of components } \partial\Delta \text{ meeting no arc of } \Gamma \\ \varepsilon(\Delta) &= \text{number of arcs of } \Gamma \text{ in } \partial\Delta \text{ meeting no vertex of } \Gamma \\ &= \text{the number of arcs which are closed curves or of type } I \text{ in } \partial\Delta. \end{aligned}$$

If $\chi(\Delta) = 1$ and $t(\Delta) = 2$ then we call Δ *collapsible*.

6.1. Boundary intervals. Let β be a connected component of $\partial\Sigma$, with orientation ζ , and let $b_1, \dots, b_n$ be n points occurring consecutively around β (read in the direction of $\zeta(\beta)$). We call $(b_1, \dots, b_n)$ a *boundary partition* of β . The closed intervals $[b_i, b_{i+1}]$ of β , for $i = 1, \dots, n$ (subscripts modulo n) are called *boundary intervals* of β . The boundary interval $[b_i, b_{i+1}]$ has *length* $||[b_i, b_{i+1}]||$ equal to the number of connected components of $[b_i, b_{i+1}] \setminus \Gamma$. (The b_i 's do not have to belong to $\Gamma \cap \beta$.)

Let $\mathcal{L}$ be a consistent system of parameters and $z = \prod_{i=1}^n (h_i, m_i) \in F(H^\Lambda)$. The boundary partition $(b_1, \dots, b_n)$ is said to $\mathcal{L}$ -*admit* z if (with subscripts modulo n)

- (i) $||[b_i, b_{i+1}]|| = 1$, for all i such that $l(h_i) = 1$,
- (ii) $||[b_i, b_{i+1}]|| = l(h_i \wedge m_i)$, for all i such that (h_i, m_i) is degenerate and
- (iii) $||[b_i, b_{i+1}]|| = q_i l(h_i) + k_i$, for some $k_i, q_i \in \mathbb{Z}$ such that $q_i \geq 0$ and $\mathcal{L}$ implies that $m_i \equiv k_i \pmod{l(h_i)}$, for all i such that $l(h_i) > 1$ and (h_i, m_i) is non-degenerate.

If α is a solution to $\mathcal{L}$, the boundary partition $(b_1, \dots, b_n)$ $\mathcal{L}$ -admits z and $[b_i, b_{i+1}]$ maps to a path in Y representing $\hat{\alpha}(h_i, m_i)$, then we say that $[b_i, b_{i+1}]$ has *label* $\hat{\alpha}(h_i, m_i)$ and *prime label* (h_i, m_i) , for $i = 1, \dots, n$. In this case the label of β is $\hat{\alpha}(z)$ and we say that β has *prime label* z with *partition* $(b_1, \dots, b_n)$.

Let $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 3-partition of (n, t, p) of length k , with $\mathbf{n} = (n_1, \dots, n_k)$ etc.. We shall say that a surface has *type* $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ if it is a disjoint union of compact surfaces Σ_i , $i = 1, \dots, k$, such that Σ_i has n_i boundary components, genus $t_i + p_i/2$ and is orientable if $p_i = 0$. In this case, if Γ is a picture on $\Sigma_1 \dot{\cup} \dots \dot{\cup} \Sigma_k$ we say that Γ has *partitioned boundary labels list* $(u_1, \dots, u_n)$ if the boundary labels of $\Gamma \cap \Sigma_i$ are $u_{\mu_i+1}, \dots, u_{\mu_i+n_i}$, for $i = 1, \dots, k$, where μ_i is defined as above.

Now with $H = A * B$ and $G = H/N(s)$ or $G = H$ as above, let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, $\mathbf{z}$ a special element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ and α a solution to $\mathcal{L}$. Suppose that Γ is a picture on a surface of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with partitioned boundary labels list $(\hat{\alpha}(z_1), \dots, \hat{\alpha}(z_n))$. Then we may choose a boundary partition $\mathbf{b}_i$ for the boundary component β_i labelled $\hat{\alpha}(z_i)$ such that $\mathbf{b}_i$ $\mathcal{L}$ -admits z_i . Then β_i has prime label z_i with partition $\mathbf{b}_i$, for all i .

Definition 6.3. Given such boundary partitions we set $\mathbf{b} = \mathbf{b}_1, \dots, \mathbf{b}_n$ and say that Γ has *boundary partition* $\mathbf{b}$ and *prime labels* $\mathbf{z}$ with *partition* $\mathbf{b}$. For ease of notation we write $\hat{\alpha}(\mathbf{z})$ for $(\hat{\alpha}(z_1), \dots, \hat{\alpha}(z_n))$.

Definition 6.4. Let a be a boundary arc and let x be an endpoint of a on $\partial\Sigma$. Then x meets 2 distinct boundary corners: denote the closures of these corners by c_1 and c_2 . If $\text{int}(c_1 \cup c_2)$ contains a point of a boundary partition then a is called an H^Λ -arc. A vertex incident to an H^Λ -arc is called an H^Λ -vertex. The sets of H^Λ -arcs and H^Λ -vertices of Γ are denoted by $\mathcal{A}_B$ and $\mathcal{V}_B$, respectively.

Definition 6.5. A boundary interval with prime label a minor H^Λ letter is called a *minor* boundary interval. A minor boundary interval with label of exponential length at least 1 is called a *partisan* boundary interval. The boundary components of Σ containing partisan boundary intervals are called *partisan* boundary components. A region of Γ containing a partisan boundary interval in its boundary is called a *partisan* region.

For future use we record the following result.





Figure 2:

Figure 3: $\sim r^{-m}$ denotes a cyclic permutation of r^{-m}

Lemma 6.6. *The number of H^Λ -vertices of Γ is at most $2W_1$ (see (3.15)).*

Proof. Traverse each boundary component of $\partial\Sigma$ once in the direction of orientation ζ counting H^Λ -vertices as they occur.

6.2. Bridge moves and vertex cancellation. Let u_1 and u_2 be vertices of a picture Γ , with orientation ζ , and let e be an arc with end-points x_1 and x_2 on u_1 and u_2 , respectively. Let N be a regular neighbourhood of $u_1 \cup u_2 \cup e$. If N is orientable let ζ_e be an orientation of N . If $\zeta_e(u_i) = \zeta(u_i)$, for both $i = 1$ and 2 , or $\zeta_e(u_i) \neq \zeta(u_i)$, for both $i = 1$ and 2 , then set $\delta = 1$. Otherwise set $\delta = -1$. If N is non-orientable set $\delta = 1$. The latter occurs if $u_1 = u_2$ and N is homeomorphic to a Möbius band (see Figure 2(a)). For $i = 1, 2$ set $\varepsilon_i = 1$ if u_i is positive and $\varepsilon_i = -1$ if u_i is negative.

Definition 6.7. If $\delta\varepsilon_1\varepsilon_2 = -1$ we say that u_1 and u_2 are *e-confluent*.

Thus u_1 and u_2 are *e-confluent* if f preserves orientation ζ_e on precisely one of ∂u_1 and ∂u_2 . Figure 3(a) shows a pair of *e-confluent* vertices, with $\varepsilon_1 = \delta = 1$ and $\varepsilon_2 = -1$. Figure 3(b) shows a pair of vertices which are not *e-confluent*, with $\varepsilon_2 = \delta = -1$ and $\varepsilon_1 = 1$.




Figure 4: Cancelling vertices

If $u_1 = u_2$ and a regular neighbourhood N of $u_1 \cup e$ is non-orientable then u_1 is e -confluent to itself, as in Figure 2(a). On the other hand if N is orientable, as in Figure 2(b), then u_1 is not e -confluent to itself.

In the above setting let w_i be the label of u_i with base point x_i (so w_i is obtained by reading in the direction opposite to $\zeta(u_i)$). If $w_1 w_2^\delta = 1$ in H then u_1 and u_2 are said to be vertices which *cancel* (see Figure 4). A picture with no pair of distinct vertices which cancel is said to be *reduced*.

Suppose that u_1 and u_2 cancel. If u_1 and u_2 are e -confluent then we may assume, without loss of generality, that $\varepsilon_1 = \delta = 1$ and $\varepsilon_2 = -1$, that $w_1 = r^m$ and w_2 is a cyclic permutation of r^{-m} , so w_2^{-1} is a cyclic permutation of r^m . Suppose that $r = r_0 r_1$ and $w_2^{-1} = (r_1 r_0)^m$. As u_1 and u_2 cancel and $\delta = 1$ we have

$$(r_0 r_1)^m = w_1 = w_2^{-1} = (r_1 r_0)^m.$$

It follows that r^m has periods $\lambda = \min\{|r_0|, |r_1|\}$ and l . From [16, Section 3, Proposition 1] it follows that r^m , and hence r , has period $d = \gcd(\lambda, l)$. By assumption r is not a proper power so this cannot happen unless $d = l$ and $\lambda = 0$, in which case $w_2^{-1} = r^m$.

Now suppose that u_1 and u_2 cancel but are not e -confluent. Without loss of generality we may assume that $\varepsilon_1 = \varepsilon_2 = \delta = 1$, that $w_1 = r^m$ and that $w_2 = r_1 r^{m-1} r_0$, where $r = r_0 r_1$. We have

$$(r_1^{-1} r_0^{-1})^m = w_1^{-1} = w_2 = r_1 r^{m-1} r_0,$$

so $r_1^2 = r_0^2 = 1$. It follows that r has a cyclic permutation $xUyU^{-1}$, where $x, y \in A \dot{\cup} B$, with $x^2 = y^2 = 1$. That is G is of type $E(2, 2, m)$.

Now suppose that a and b are distinct arcs of Γ and that γ is the image of an embedding of $[0, 1]$ in Σ with one end-point on a and the other on b such that $\gamma \cap \Gamma = \{a, b\}$. Then $f(\gamma)$ is a closed path in (X_O, p) , for $O = A$ or B . If $f(\gamma)$ is nullhomotopic, then after a homotopy (constant outside a regular neighbourhood of γ) we may assume that $f(\gamma) = p$. A further homotopy can be effected to make $f(\gamma) = x_O$, resulting in a new picture, as shown in Figure 5. We call the transformation from the original to the new picture a *bridge move*.

If u and v are a pair of distinct vertices which cancel then we can perform bridge moves on arcs incident u and v , as shown in Figure 6, to increase the number of arcs joining u to v . Repeating such bridge moves we obtain a picture in which all




Figure 5: A bridge move




Figure 6: Bridge moves on a pair of cancelling vertices

arcs incident to u are incident to v , in which case we call u, v and their incident arcs a *floating dipole*: see Figure 7. An arc c of Γ such that c is a simple closed curve bounding a disk D with $\text{int}(D) \cap \Gamma = \emptyset$ is called a *floating arc*.

If C is a simple closed arc in $\Sigma \setminus \Gamma$ and one component of $\Sigma \setminus C$ is a disk C_D such that $\Gamma \cap C_D$ is a floating dipole or floating arc then $f(C)$ is null-homotopic in Y . Performing a homotopy, constant outside $C_D \cup C$, we may replace f by a map such




Figure 7: A floating dipole

that $f(C_D) = x_O$, where $O = A$ or B . Therefore we may form a new picture on Σ by removing all arcs and vertices of $C_D \cap \Gamma$ from Γ . We call this operation *removal* of a floating dipole or arc. We consider the equivalence relation $\sim$ generated by the relations (i)–(iii) below on the set of pictures on a surface Σ , with n boundary components with fixed boundary labels $u_1, \dots, u_n \in H$. Suppose Γ and Γ' are such pictures on Σ over G . Then $\Gamma \sim \Gamma'$ if Γ' is obtained from Γ by

- (i) an isotopy of Σ ,
- (ii) a bridge move or
- (iii) removal of a floating dipole or floating arc.

Definition 6.8. A picture is *efficient* if it has fewest arcs in its equivalence class.

It follows that an efficient picture is reduced, since otherwise we can perform bridge moves on arcs incident to a pair of vertices which cancel until a floating dipole is produced. Removal of the floating dipole results in a picture with fewer arcs, contrary to efficiency of the original picture.

Let g and n be fixed integers and $u_1, \dots, u_n$ fixed elements of H . Consider the set T of all pictures over G on surfaces with n boundary components and genus at most g , with boundary labels $u_1, \dots, u_n$.

Definition 6.9. A picture in T is said to be *minimal* if no other picture in T has fewer arcs.

Minimal pictures are efficient and efficient pictures are reduced. In general it is not the case that reduced pictures are efficient or that efficient pictures are minimal.

Proposition 6.10 ([7]). *Let Γ be a minimal picture over G on a surface Σ . Then no arc of Γ is closed curve.*

Proof. Suppose that c is an arc of Γ and a closed curve. The closure of $\Sigma \setminus c$ has 2 boundary components c_1 and c_2 corresponding to c . Form a new surface Σ_1 by capping c_i with a disk, for $i = 1, 2$. Then $\Gamma \setminus c$ is a picture on Σ_1 , since $f(c) = p \in Z$. The genus of Σ_1 is no more than that of Σ and $\Gamma \setminus c$ has the same boundary labels as Γ , but fewer arcs. This contradicts the minimality of Γ .

Proposition 6.11 ([7]). *Let Γ be a minimal picture over G on a compact connected surface Σ . Then $\chi(\Delta) \geq \chi(\Sigma)$, for all regions Δ of Γ .*

Proof. Suppose that Δ is a region such that $\chi(\Delta) < \chi(\Sigma)$. Let $\Sigma_0 = \Sigma \setminus \text{int}(\Delta)$ and let $\Omega_0, \dots, \Omega_k$ be the connected components of Σ_0 . Then $\chi(\Delta) < \chi(\Sigma) = \chi(\Sigma_0) + \chi(\Delta) = \sum_{i=0}^k \chi(\Omega_i) + \chi(\Delta)$. Therefore $\chi(\Omega_i) > 0$, so Ω_i is a disk or a sphere, for some i . As Σ is connected this means that Ω_i is a disk with $\Omega_i \cap \Delta = C$, where C is homeomorphic to S^1 , for such i . Let Ω' be a regular neighbourhood of Ω_i and let $\Gamma' = \Gamma \cap \Omega'$. Then Ω' is a disk and the boundary of Ω' lies in Δ . Hence the boundary label of Γ' is an element u of $A \dot{\cup} B$ such that $u = 1$ in G . As the Freiheitssatz holds for G ([16]) it follows that $u = 1$ in A or B , as appropriate. Thus we may replace Γ' with the empty picture on Ω' without altering Σ or the boundary labels of Γ . This contradicts the minimality of Γ .

6.3. Parallel arcs and the graph of a picture. Given two arcs a_1 and a_2 of a picture Γ such that either

- (i) a_1 and a_2 bound an annulus S with $\text{int}(S) \cap \Gamma = \emptyset$ or
- (ii) there are corners c_1 and c_2 of Γ such that a_1, a_2, c_1 and c_2 bound a disk Δ with $\text{int}(\Delta) \cap \Gamma = \emptyset$,

then a_1 and a_2 are said to be *parallel* arcs. There is an equivalence relation on the set of arcs of Γ generated by the parallel relation on arcs. The equivalence class of an arc a is called an *edge* of Γ , denoted $\bar{a}$. If an edge consists of d parallel arcs we say that it has *width* d . We form the *graph* $\bar{\Gamma}$ of Γ on Σ as follows. Collapse each vertex of G to a point to give a vertex of G . Collapse each class of arcs $\bar{a}$ onto a : the resulting arcs are the edges of $\bar{\Gamma}$, except for any that are closed loops (meeting no vertex of Γ) and these are discarded. In addition there is one vertex on $\partial\Sigma$ wherever an edge of $\bar{\Gamma}$ meets $\partial\Sigma$. If e^+ is one end of an edge e of $\bar{\Gamma}$ then the end points of arcs of Γ which are identified to form e^+ comprise an *end* of the class of arcs corresponding to e . If $\bar{a}$ is a class of arcs with at least one end on the boundary then we say that $\bar{a}$ is a *boundary* class. Otherwise $\bar{a}$ is an *interior* class. An arc of Γ with both ends on $\partial\Sigma$ is called an *arc of type I*. An edge of $\bar{\Gamma}$ which is the image of a class of arcs of type I is called an *edge of type I*. We define the *degree* of a vertex v of Γ to be the degree of the corresponding vertex in the graph $\bar{\Gamma}$, that is $\deg(v) = \text{number of classes of arc incident to } v$.

If a is an arc of an edge e and a is incident to vertices u and v then u and v are *a-confluent* if and only if they are *b-confluent* for all arcs b of e . In this case we say that u and v are *e-confluent*.

The following facts are established in [16]. A word $x_1 \cdots x_n$ is said to have *period* λ if $x_{i+\lambda} = x_i$, for all i such that $1 \leq i \leq n - \lambda$.

Lemma 6.12. *Assume G is not of type $E(2, *, m)$. Let vertices u_1 and u_2 of a picture Γ over G be joined by an edge e of width k . Assume either that Γ is reduced and u_1 and u_2 are distinct or that Γ is minimal. Then $k \leq l - 2$.*

Proof. The ends of e determine subwords s_1 and s_2 of the boundary labels w_1 and w_2 of u_1 and u_2 , respectively, with $l(s_1) = l(s_2) = k - 1$. Suppose first that u_1 and u_2 are *e-confluent*. Without loss of generality we may assume that $\varepsilon_1 = \delta = 1$ and $\varepsilon_2 = -1$ (with the notation of Definition 6.7). Then s_1 is a cyclic subword of r^m , s_2 is a cyclic subword of r^{-m} and $s_2^{-1} = s_1$ (see Figure 8). Therefore r^m has







Figure 8:


 Figure 9: N after cutting along arc ab

identical subwords s_1 and s_2^{-1} . We may assume that s_1 is an initial subword of r^m . Then r^m begins $r_0 s_2^{-1}$, where r_0 is an initial subword of r of length $l(r_0) = \lambda \geq 0$. As r is cyclically reduced in $A * B$ it follows that λ is even. We may also assume that $\lambda \leq l/2$ (otherwise reverse the roles of s_1 and s_2). If $k \geq l - 1$ this implies that the union of s_1 and s_2 is an initial subword of r^m of period λ . If $\lambda > 0$ then $\lambda \geq 2$ so $l(s_2) + \lambda \geq l$ and then it follows that r^m has periods λ and l . Hence, from [16, Section 3, Proposition 1], r^m and also r have period $d = \gcd(\lambda, l)$. This cannot occur as r is not a proper power. Therefore $\lambda = 0$ and s_1 and s_2^{-1} coincide as subwords of r^m . In this case, if $u_1 \neq u_2$ then u_1 and u_2 cancel. As Γ is reduced this is a contradiction.

On the other hand, if $\lambda = 0$, $u_1 = u_2$ and u_1 is e -confluent to itself then a regular neighbourhood N of $u_1 \cup e$ is homeomorphic to a Möbius band. In this case we can perform bridge moves on arcs incident to u_1 and outside e to form a class containing all arcs incident to u_1 . We illustrate this diagrammatically: suppose N is as shown in Figure 9. Cutting along c and pasting together both copies of a and b we obtain the diagram of N shown in Figure 10. Clearly the number of arcs incident to the upper and lower halves of the vertex in this diagram are equal and so we may perform bridge moves as claimed. The result is shown in Figure 11. Since the dotted boundary of this diagram has trivial label we may remove u_1 and all its incident arcs from N , leaving a picture on Σ with the same boundary labels as Γ but fewer arcs. This contradicts minimality of Γ . Thus if u_1 and u_2 are e -confluent then $k \leq l - 2$.

Figure 10: N after cutting along c and pasting a and b Figure 11: N after performing bridge moves

Now suppose that u_1 and u_2 are not e -confluent and that $k \geq l-1$. Without loss of generality we may assume that $\varepsilon_1 = \varepsilon_2 = \delta$ and that s_1 is an initial subword of r^m . Then s_2 is a cyclic subword of r^m and s_1 is identically equal to s_2^{-1} , as a word in $A * B$. Again we may assume that r^m begins $r_0 s_2 = U$, with $l(r_0) = \lambda \leq l/2$. If s_1 has even length then s_1 and s_2 begin with letters of different groups, so λ




Figure 12:

must be odd. Similarly if $l(s_1)$ is odd then λ must be even. As $k \geq l - 1$ we have $l(s_i) \geq l - 2$ and $l(U) = l(s_2) + \lambda \geq l - 1$. We may write $s_1 = r_0 t$, where $s_2 = t r_1$, since $\lambda \leq l - 2 \leq l(s_i)$. Hence $s_1 = s_2^{-1}$ implies $t = t^{-1}$ and $r_0 = r_1^{-1}$. Hence $U = r_0 s_2 = r_0 t r_0^{-1}$, where $t = t^{-1}$, so t is of odd length and its middle letter has order 2. Therefore $U = V y V^{-1}$, for some V . If $l(U) = l - 1$ then $r = V y V^{-1} x$ and G is of type $E(2, *, m)$, a contradiction. Suppose $l(U) \geq l$. We have $r^m = V y V^{-1} W$ and we may write $V = V_0 x U_0$, where $x \in A \cup B$, $l(U_0 y U_0^{-1}) = l - 1$ and $l(V_0) \geq 0$. Hence r^m has a cyclic permutation $x U_0 y U_0^{-1} x^{-1} V_0^{-1} W V_0$ which must have period l . Again a contradiction arises as $x = x^{-1}$ and G is of type $E(2, 2, m)$.

Definition 6.13. A picture Γ on a surface Σ is called *minimalistic* if the following conditions hold.

- (i) No arc of Γ is a closed curve.
- (ii) $\chi(\Delta) \geq \chi(\Sigma_\Delta)$, whenever Δ is a region of Γ and Σ_Δ is the connected component of Σ containing Δ .
- (iii) No interior edge of Γ has width more than l . If G is not of type $E(2, *, m)$ then no interior edge has width more than $l - 2$.

From Proposition 6.10, Proposition 6.11 and Lemma 6.12, minimal pictures are minimalistic. The converse does not hold in general.

6.4. Routes and distance. If u and v are vertices incident to a region Δ satisfying $\rho(\Delta) = \beta(\Delta) = 2$, $\chi(\Delta) = 1$, $t(\Delta) = 4$ and $\varepsilon(\Delta) = 0$, as shown in Figure 12(a), then u and v are said to be *paired*. If e is a class of arcs having both end points on $\partial\Sigma$ and $e \cap \partial\Delta \neq \emptyset$, where Δ is a region satisfying $\rho(\Delta) = 1$, $\beta(\Delta) = 2$, $\chi(\Delta) = 1$, $t(\Delta) = 3$ and $\varepsilon(\Delta) = 0$, as shown in Figure 12(b), then u and e are said to be *paired*.

A *route* in Γ , from x_1 to x_n , is a sequence $x_1, \dots, x_n$, where x_i is a vertex or a class of arcs such that the set $\{x_i, x_{i+1}\}$ consists either of

- (i) two vertices which are joined by an arc or
- (ii) a vertex and a boundary class of arcs paired to that vertex or
- (iii) two vertices which are paired,

for $i = 1, \dots, n$. Let $l_1 = \{i : x_i \text{ is a boundary class of arcs}\}$ and $l_2 = \{i : x_i \text{ and } x_{i+1} \text{ are both vertices}\}$. The *length* of the route is then $|l_1| + |l_2|$. A *route* in $\bar{\Gamma}$ is defined to be the image of a route in Γ under the quotient map $\Gamma \longrightarrow \bar{\Gamma}$. A route is *simple* if $x_i \neq x_j$, for all $i \neq j$.

If S is a subset of $\mathcal{V}$ then a vertex v is said to be *distance* d from S if the length of a shortest route from v to a vertex of S is d . If there is no route from v to a vertex of S then the distance of v from S is defined to be ∞ . If e is a class of arcs that is paired to a vertex then the *distance* of e from S is $d + 1$, where d is the minimum of the distances from S of the vertices paired to e .

6.5. Pictures and equations. The correspondence between solutions of quadratic equations and pictures is given in the next three results.

Theorem 6.14. *Let $u_1, \dots, u_k$ be elements of H and let t, p be non-negative integers. Then there exist elements $w_1, \dots, w_{k+2t+p} \in H$ such that*

$$(6.2) \quad w = \prod_{i=1}^k w_i^{-1} u_i w_i \prod_{i=1}^t [w_{i+k}, w_{i+k+t}] \prod_{i=1}^p w_{i+k+2t}^2 = 1 \in G$$

if and only if there exists a picture Γ over G on a surface Σ , of genus $t + p/2$, orientable if $p = 0$, with k boundary components which have labels $u_1, \dots, u_k$ (with respect to some orientation of $\partial\Sigma$).

Proof. In the notation of Section 2, let $F = F(D \dot{\cup} X)$ and $q = q(0, k, t, p)$. Then $\Sigma(q)$ is the connected sum of a k -punctured sphere with t torii and p projective planes, and so has genus $t + p/2$ and is orientable if $p = 0$, and has boundary components $\beta_1, \dots, \beta_k$ with labels $d_1, \dots, d_k$, respectively.

Suppose a picture (f, Γ) with orientation ζ exists on a surface Σ of genus $t + p/2$, orientable if $p = 0$, with boundary components $\tilde{\beta}_1, \dots, \tilde{\beta}_k$ having labels $u_1, \dots, u_k$. Assume first that Σ is connected. Then Σ is homeomorphic to $\Sigma(q)$ and we may choose this homeomorphism $\theta : \Sigma(q) \longrightarrow \Sigma$ so that the boundary component β_i is mapped to $\tilde{\beta}_i$, so that the base point of β_i is mapped to a base point of $\tilde{\beta}_i$ and so that the orientation of $\tilde{\beta}_i$ is the same as that induced by $\theta(\beta_i)$. By construction $\Sigma(q)$ is a quotient of the disk D with boundary ∂D labelled q , so $f \circ \theta$ can be regarded as a map from D to Z . Then $f \circ \theta|_{\partial D}$ maps each labelled interval of ∂D to a path in Y representing an element of H . Let ϕ be the map from F to H induced by $f \circ \theta|_{\partial D}$ on the labelled intervals of ∂D . Then ϕ maps the boundary label q of D to an element of H representing the identity of G . Furthermore $\phi(d_i) = u_i$, for $i = 1, \dots, k$ and $\phi(x_i) = w_i \in H$, for $i = 1, \dots, k + 2t + p$, and it follows that $w = 1$ in G , as required.

Suppose now that Σ has connected components $\Sigma_1, \dots, \Sigma_n$, for some $n \geq 2$. We may assume that Σ_j is the connected sum of a k_j -punctured sphere with t_j torii and p_j projective planes where $k_j > 0$, $\sum_{j=1}^n k_j = k$, $\sum_{j=1}^n t_j = t$ and $\sum_{j=1}^n p_j = p$. Let the boundary labels of $\Gamma_i = \Sigma_i \cap \Gamma$ be $u_{i_1}, \dots, u_{i_{k_j}}$, where $i_1 < \dots < i_{k_j}$. The argument above implies the existence of elements $w_{ij}, x_{ij}, y_{ij}, z_{ij} \in H$ such that

$$\prod_{i=1}^{k_j} w_{ij}^{-1} u_{ij} w_{ij} \prod_{i=1}^{t_j} [x_{ij}, y_{ij}] \prod_{i=1}^{p_j} z_{ij}^2 = 1 \in G,$$

for $j = 1, \dots, n$. The equality (6.2) follows after some rearrangement of these equations.

Conversely, suppose that there exist w_i in H satisfying (6.2). Let F , q and $\Sigma(q)$ be defined as above, where $\Sigma(q)$ is a quotient of a disk D with boundary label q . Let $\hat{d}_i$ and $\hat{x}_i$ be the oriented sub-intervals of ∂D labelled d_i and x_i , respectively. For $g \in H$ let $\tilde{g}$ be a closed path in (Y, p) representing g . Define a map $f_\partial : \partial D \rightarrow Z$ by $f_\partial(\hat{x}_i) = \tilde{w}_i$ and $f_\partial(\hat{d}_i) = \tilde{u}_i$. As $w = 1$ in G the map f_∂ maps ∂D to a path which is nullhomotopic in Z and so extends to a map $f : D \rightarrow Z$. As f_∂ is identical on any two intervals of ∂D which are identified in forming $\Sigma(q)$, f factors through $\Sigma(q)$ and we may regard f as a map from $\Sigma(q)$ to Z . We may make f transverse to the mid-point c of Δ_s with a homotopy which fixes $\partial \Sigma(q)$. A further homotopy ensures that each component of the closure of $f^{-1}(\text{int}(\Delta_s))$ is a regular neighbourhood of $f^{-1}(c)$. Let $\Sigma_0 = \Sigma(q) \setminus f^{-1}(\text{int}(\Delta_s))$. Then f restricted to $\partial \Sigma_0$ is transverse to p and so there is a homotopy of f , relative to $\partial \Sigma_0$, to a map which is transverse to p on Σ_0 . Therefore we may assume that f gives rise to a sketch Γ on $\Sigma(q)$. Extend the orientation of the $\hat{d}_i$'s to an orientation of regions and vertices of $\Sigma(q)$. (This can be done consistently if $\Sigma(q)$ is orientable.) Let $f^l(d_i) = u_i$ and define f^l on vertex corners as in the definition of a picture. Then (f, Γ) is a picture on Σ over G with boundary labels $u_1, \dots, u_k$, as required.


Corollary 6.15. *Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, $\mathcal{L}$ a consistent system of parameters and $\mathbf{z}$ an element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$. Then the triple $(\mathbf{n}, \mathbf{t}, \mathbf{p}) \in \mathcal{L}$ -genus($\mathbf{z}$) if and only if there exists a solution α to $\mathcal{L}$ and a minimalistic picture Γ on a surface of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with partitioned boundary labels list $(\hat{\alpha}(z_1), \dots, \hat{\alpha}(z_n))$.*

Proof. Given a solution (ϕ, α) to $Q = Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ we have $\phi(q(\xi_j, n_j, t_j, p_j)) = 1$ in G , for $j = 1, \dots, k$. It follows, from Theorem 6.14, that there exists a picture Γ_j over G on a surface Σ_j of genus $t_j + p_j/2$, with n_j boundary components labelled $\phi(d_{\xi_j+1}), \dots, \phi(d_{\xi_j+n_j})$. If Γ_j exists then there also exists a minimalistic picture satisfying the same conditions, so we may assume that Γ_j is minimalistic. As (ϕ, α) is a solution to Q , $\phi(d_{\xi_j+i}) = \hat{\alpha}\beta(d_{\xi_j+i}) = \hat{\alpha}(z_{\mu_j+i})$, for all i, j . The picture $\Gamma = \Gamma_1 \cup \dots \cup \Gamma_k$ on $\Sigma_1 \dot{\cup} \dots \dot{\cup} \Sigma_k$ therefore satisfies the given requirements.

Conversely, given a solution α to $\mathcal{L}$ and a picture Γ on a surface Σ as described, suppose that Σ has connected components $\Sigma_1, \dots, \Sigma_k$, where Σ_j has genus $t_j + p_j/2$ and n_j boundary components labelled $\hat{\alpha}(z_{\mu_j+1}), \dots, \hat{\alpha}(z_{\mu_j+n_j})$. Let $\Gamma_j = \Gamma \cap \Sigma_j$ and $q_j = q(\xi_j, n_j, t_j, p_j)$ and define $\beta_j : F(L_D(q_j)) \rightarrow H$ by $\beta_j(d_{\xi_j+i}) = \hat{\alpha}(z_{\mu_j+i})$, for $j = 1, \dots, k$ and $i = 1, \dots, n_j$. Then, from Theorem 6.14, Γ_j gives rise to a solution ϕ_j to $(q_j = 1, \beta_j)$. Define $\phi : F(L(\mathbf{q})) \rightarrow H$ by $\phi(x_i) = \phi_j(x_i)$, for $\xi_j + 1 \leq i \leq \xi_j + n_j + 2t_j + p_j$ and $\phi(d_{\xi_j+i}) = \hat{\alpha}(z_{\mu_j+i})$, for $1 \leq i \leq n_j$. Then (ϕ, α) is a solution to Q .

Definition 6.16. Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, $\mathcal{L}$ a consistent system of parameters and $\mathbf{z}$ an element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$. Let α be a solution to $\mathcal{L}$ and Γ a picture on a surface of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with partitioned boundary labels list $(\hat{\alpha}(z_1), \dots, \hat{\alpha}(z_n))$. Then we say that (α, Γ) *corresponds* to the solution (ϕ, α) of $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ constructed in Corollary 6.15. The pair (α, Γ) is called *exponent-minimal* if it corresponds to an exponent-minimal solution (ϕ, α) to $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$.

When there is no ambiguity we refer to Γ , rather than (α, Γ) , as exponent-minimal. Observe that if an exponent-minimal picture with prime labels $\mathbf{z}$ exists then a reduced (or efficient or minimal) exponent-minimal picture with the same prime labels also exists, because all pictures in an equivalence class and in T have the same boundary labels.

Figure 13: A picture of type square-1 with base $D(2, 3, 1, 2)$

7. CORRIDORS

Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, $\mathcal{L}$ a consistent system of parameters, $\mathbf{z}$ a special element of $(H^A, \mathcal{L}, \mathbf{n}, \mathbf{h})$ and α a solution to $\mathcal{L}$. From now until further notice we assume that Γ is a picture on a surface of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with prime labels $\mathbf{z}$, labelled by $\hat{\alpha}(\mathbf{z})$ and with boundary partition $\mathbf{b} = (b_1, \dots, b_{W_1})$.

Let D be the disk $[a, a+1] \times [0, 1] \subset \mathbb{R}^2$, where $a \geq 0$ is an integer, and let x_0, y_0, x_1 and y_1 be non-negative integers with $x_i > 0$, $i = 1, 2$. Define a set $D(x_0, x_1, y_0, y_1)$ of *distinguished* points of D to be

- (i) $(a, i/(y_0 + 1))$, for $i = 1, \dots, y_0$;
- (ii) $(a + 1, i/(y_1 + 1))$, for $i = 1, \dots, y_1$;
- (iii) $(a + (i/(x_0 + 1)), 0)$, for $i = 1, \dots, x_0$ and
- (iv) $(a + (i/(x_1 + 1)), 1)$, for $i = 1, \dots, x_1$.

A picture Γ_D over G on D is said to have *type square-0* (with *base* $D(1, 1, 0, 0)$) if Γ_D has no vertices and one arc $\{1/2\} \times [0, 1]$. A picture Γ_D on D is said to have *type square-1* with *base* $D(x_0, x_1, y_0, y_1)$ if x_0, x_1, y_0 and y_1 are non-negative integers such that $x_0, x_1 > 0$, $x_0 + x_1 + y_0 + y_1 = ml$, Γ_D has one vertex v and Γ_D has ml arcs each beginning at a point of ∂v and ending at one of the distinguished points of $D(x_0, x_1, y_0, y_1)$. A picture of type square-1 with base $D(2, 3, 1, 2)$ is shown in Figure 13.

A picture Γ_D over G on D is said to have *type square-2* with *base* $D(x_0, x_1, y_0, y_1)$ if the following conditions hold.

- (i) x_0, x_1, y_0 and y_1 are positive integers such that $y_0, y_1 \geq 3$ and $x_0 + x_1 + y_0 + y_1 \leq 2ml - 1$.
- (ii) Γ_D has 2 vertices v_0 and v_1 and $ml + (x_0 + x_1 + y_0 + y_1)/2$ arcs. Γ_D has one non-empty class of arcs joining v_0 to v_1 and Γ_D has x_i arcs joining v_i to distinguished points of D on $[a, a+1] \times \{i\}$, for $i = 1, 2$.
- (iii) There are integers r_i , $i = 0, 1$ such that $0 < r_i < y_i$ and Γ_D has r_i arcs joining v_i to distinguished points of $\{a+i\} \times [0, 1]$ and $y_i - r_i$ arcs joining v_{1-i} to distinguished points of $\{a+i\} \times [0, 1]$.

A picture of type square-2 with base $D(4, 5, 7, 6)$, $r_0 = 3$ and $r_1 = 2$ is shown in Figure 14. A picture of type square- j is called a *square-picture* of type j , where $j = 0, 1$ or 2 . An arc e of a square-picture meeting $[a, a+1] \times \{i\}$, for $i = 0$ or 1 is




 Figure 14: A picture of type square-2 with base $D(4, 5, 7, 6)$


Figure 15:

called a *lateral* arc of the picture. An arc e of a square-picture of type 2 incident to both vertices is called an *internal-lateral* arc.

A square-picture (f_D, Γ_D) comes equipped with a labelling function f_D^l , from corners of Γ_D to $A \dot{\cup} B$. Since we wish to map square-pictures into arbitrary pictures in such a way that the intervals $([a + (k/(x_i + 1)), a + ((k + 1)/(x_i + 1))] \times \{i\})$ correspond to boundary corners of Γ we now refine this labelling function. We redefine the label of the corner $c_0 = (\{a\} \times [0, 1/(y_0 + 1)]) \cup ([a, a + (1/(x_0 + 1))] \times \{0\})$ setting $f_D^l([a, a + (1/(x_0 + 1))] \times \{0\}) = 1$ and $f_D^l(\{a\} \times [0, 1/(y_0 + 1)]) =$ the original label of c_0 . Similarly, we redefine the labelling function on the corner $c_1 = (\{a\} \times [y_0/(y_0 + 1), 1]) \cup ([a, a + (1/(x_1 + 1))] \times \{1\})$ so that the subinterval $[a, a + (1/(x_1 + 1))] \times \{1\}$ has trivial label and the defining property of boundary labels of regions of pictures is preserved (see Figure 15). We say such a labelling function on Γ_D is *left-trivial*. If Γ_D is of type 0 all labels are trivial so any labelling function is left-trivial. If Γ_D is not of type 0 then, since a regular neighbourhood of an arc of Γ_D maps to a contractible subset of Z , we may achieve a left-trivial labelling by a homotopy of f_D fixing all of D except for the components of $D \setminus \Gamma_D$ meeting c_0 and c_1 . We shall henceforward assume that square-pictures have left trivial labelling functions.




Figure 16:




Figure 17:

If D is the disk $[a, b] \times [0, 1]$, where $a, b \in \mathbb{Z}$, $0 \leq a < b$, we define D_c to be the disk $[c, c+1] \times [0, 1]$, for $c \in \mathbb{Z}$, $a \leq c < b$. We define a *sum* of pictures of *type 1* to be a picture Γ_D on the disk $D = [a, b] \times [0, 1]$, where

- (i) $b - a \geq 1$,
- (ii) $\Gamma_D \cap D_c$ is a picture of type square-0 or square-1, for $a \leq c \leq b-1$, and there is at least one c in this range such that $\Gamma_D \cap D_c$ is a picture of type square-1 and
- (iii) if $\Gamma_D \cap D_c$ has base $D_c[x_0, x_1, y_0, y_1]$ then $\Gamma_D \cap D_{c+1}$ has base $D_{c+1}[x'_0, x'_1, y_1, y'_1]$, for $a \leq c \leq b-2$.

For $j = 0$ or 2 we define a sum of pictures of *type j* similarly except that in condition (ii) all pictures must be of type square- j and if $j = 2$ then the following condition must also hold.

- (iv) If $b - a > 1$ then Γ_D has the form shown in either Figure 16 or 17 (where all the edges shown contain at least one arc).

Given a sum of square-pictures Γ_D on $D = [a, b] \times [0, 1]$ we set $\Gamma_D(c) = \Gamma_D \cap D_c$, for $a \leq c \leq b-1$. A sum of square-pictures on $[a, b] \times [0, 1]$ is said to have *width* $b - a$. The lateral and internal-lateral arcs of the square-pictures making up a sum of square-pictures are called *lateral* and *internal-lateral* arcs, respectively, of the sum.




Figure 18:

Let $D = [a, a+1] \times [0, 1]$ and let Γ_D be a square-picture of type j with base $D(x_0, x_1, y_0, y_1)$ on D . Suppose $j > 0$. If $j = 1$ let v_0 be the vertex of Γ_D . If $j = 2$ let v_i be the vertex of Γ_D incident to $[a, a+1] \times \{i\}$, for $i = 0, 1$. Let a_0 be the arc joining v_0 to $(a+1/(x_0+1), 0)$. Travelling clockwise around ∂v_0 , starting from the endpoint of a_0 , let a_1 be the first arc encountered which does not meet $\{a\} \times [0, 1]$. If $j = 1$ then a_1 connects v_0 to $(a+1/(x_1+1), 1)$. If $j = 2$ then a_1 connects v_0 to v_1 . In the latter case, travelling clockwise around ∂v_1 , starting from a_1 , let a_2 be the first arc encountered which does not meet $\{a\} \times [0, 1]$: so a_2 joins v_1 to $(a+1/(x_1+1), 1)$. The label of v_i , read from an appropriate starting point in a clockwise direction, is $r^{\delta_i m}$, where $\delta_i = \pm 1$, for $0 \leq i \leq j-1$. The label u_i on the subinterval of ∂v_i , read clockwise, from a_i to a_{i+1} is then a subword of $r^{\delta_i m}$, for $0 \leq i \leq j-1$. Let o_i , ρ_i and σ_i be non-negative integers such that $\rho_i < m$, $o_i < l(r)$, $\sigma_i < l(r)$ and $u_i = \tau(r^{\delta_i}, o_i)(r^{\delta_i})^{\rho_i} \iota(r^{\delta_i}, \sigma_i)$, for $0 \leq i \leq j-1$. In cases $j = 0$ or 1 define $\delta_i = o_i = \rho_i = \sigma_i = 0$, for $j \leq i \leq 1$.

Definition 7.1. The *left interior marking* of Γ_D is the sequence

$$\delta_0, o_0, \rho_0, \sigma_0, \delta_1, o_1, \rho_1, \sigma_1.$$

For example suppose that, in Figure 18, $r = ab$, $m = 4$ and the label on v_0 read clockwise from a_0 is $(ab)^4$. Then Γ_D has left interior marking $1, 0, 1, 0, 0, 0, 0, 0$. Again, in Figure 19 suppose $r = ab$, $m = 7$, the label of v_0 read from a_0 is $(ba)^7$ and the label of v_1 read from a_1 is $(a^{-1}b^{-1})^7$. Then Γ_D has left interior marking $1, 1, 1, 1, -1, 1, 2, 0$.

Let $D = [a, b] \times [0, 1]$ and suppose that there exists a map $\mu : D \rightarrow \Sigma$ and boundary intervals P_0 and P_1 of Γ satisfying the following properties.

- (i) μ is injective on $D \setminus (\{0, 1\} \times [0, 1])$;
- (ii) either μ is injective on $\{0, 1\} \times [0, 1]$ or $\mu(a, t) = \mu(b, t)$ or $\mu(a, t) = \mu(b, 1-t)$, for all $t \in [0, 1]$;
- (iii) $\partial \Sigma \cap \mu(D) = \mu([a, b] \times \{0, 1\})$ and $\mu([a, b] \times \{i\}) \subset P_i$, for $i = 0, 1$;
- (iv) $\mu(\{c\} \times [0, 1])$ is transverse to Γ and meets no vertex of Γ , for $c \in \mathbb{Z}$ with $a \leq c \leq b$.

Then we call μ a *binding map* (on D) between P_0 and P_1 . Now suppose that P_i is not a partisan boundary interval, for $i = 0, 1$, and that no arc of Γ contained in $\mu(D)$ is an H^Λ -arc. Let (f_D, Γ_D) be a sum of square-pictures of type j on D , with




Figure 19:

left-trivial labelling function f_D^l , and let $\mu(D) = B$. Let c_0, c_1 be the left-hand corners of D defined above and in addition let $c_2 = (\{b\} \times [0, 1/(y_1 + 1)]) \cup ([b - 1 + (x_0/(x_0 + 1)), b] \times \{0\})$ and $c_3 = (\{b\} \times [y_1/(y_1 + 1), 1]) \cup ([b - 1 + (x_1/(x_1 + 1)), b] \times \{1\})$. Recall that f is the map from Σ to Z described in the definition of picture.

Assume that $\mu(\Gamma_D) = \Gamma \cap B$ and

- (i) $f^l(\mu(c)) = f_D^l(c)$, for all corners c of Γ_D except c_0, c_1, c_2 and c_3 ,
- (ii) $f(\mu([a, a + (1/(x_i + 1))] \times \{i\}))$ is nullhomotopic, for $i = 0, 1$, and
- (iii) $d_i = \mu([b - 1 + (x_i/(x_i + 1)), b] \times \{i\}) \subset c'_i$, where c'_i is a boundary corner of Γ and $f(c'_i \setminus d_i)$ is nullhomotopic, for $i = 0, 1$.

Then $C = (D, \Gamma_D, \mu)$ is called a j -corridor of width $b - a$ with binding map μ binding P_0 and P_1 . In this case $\Gamma_B = \Gamma \cap \mu(D)$ is a picture on B and we call B the image of C . The images $\mu(x)$ of arcs x of the picture Γ_D are called arcs of C . If $\mu(a, t) = \mu(b, t)$ or $\mu(a, t) = \mu(b, 1 - t)$, for all $t \in [0, 1]$, then all arcs of C are arcs of Γ . Otherwise, given an arc x of Γ_D , the arc $\mu(x)$ of C is an arc of Γ if and only if x does not meet $\{a, b\} \times [0, 1]$. Nonetheless it is convenient to refer to $\mu(x)$ as an arc of C , for all arcs x of Γ_D . A corridor of width 1 is called a *compartment*. The left interior marking of rank s of the corridor C is defined to be the left interior marking of $\Gamma_D(s)$, for all s such that $a \leq s \leq b - 1$.

If e is a lateral arc, or internal-lateral arc, of Γ_D then $\mu(e)$ is called a *lateral arc*, or *internal-lateral arc*, of the corridor (D, Γ_D, μ) . If $C = (D, \Gamma_D, \mu)$ and $C' = (D', \Gamma'_D, \mu')$ are corridors with identical sets of lateral arcs then we say that C and C' are *equivalent* and this is an equivalence relation on the set of corridors. If C and C' are equivalent then μ' is homotopic to a map μ'' such that $\mu''(D') = \mu(D)$ and (D', Γ'_D, μ'') is a corridor equivalent to C . If $C' = (D', \Gamma'_D, \mu')$ is a corridor such that $\{\text{lateral arcs of } C'\} \subseteq \{\text{lateral arcs of } C\}$ then C' is called a *sub-corridor* of C . If $a \leq c < d \leq b$ and $D' = [c, d] \times [0, 1]$ then the map $\mu' = \mu|_{D'}$ is called the binding map *induced* from μ on D' and the corridor (D', Γ'_D, μ') is called the sub-corridor *induced* from (D, Γ_D, μ) , denoted $C[c, d]$ or $C[c]$, if $c + 1 = d$. If C is corridor and is not a sub-corridor of any corridor of greater width then we say that C is a *maximal* corridor. Obviously two corridors are equivalent if and only if each one is a sub-corridor of the other. If $C = (D, \Gamma_D, \mu)$ and $C' = (D', \Gamma_{D'}, \mu')$ are corridors, with images $B = \mu(D)$ and $B' = \mu'(D')$, we call $B \cap B'$ the *intersection* of C and C' , denoted $C \cap C'$, and if $B \cap B' \neq \emptyset$ we shall say that C and C' *overlap*.




Figure 20:

Definition 7.2. Choose one representative from each equivalence class of maximal j -corridors of Γ , where $j = 0$ or 1 . These corridors and the sub-corridors induced from them are called *designated* corridors of Γ (of type 0 and 1).

(We defer the definition of designated 2-corridor until later.) We may always choose the maximal designated corridor (D, Γ_D, μ) such that $D = [0, c] \times [0, 1]$, for some integer c . In the absence of any explicit statement to the contrary it is to be assumed that all corridors are designated corridors.

If (D, Γ_D, μ) is a maximal 0-corridor then Γ_D consists of arcs from a class of arcs of type I . If two distinct designated maximal 0-corridors overlap then we may choose different representatives of the two equivalence classes of corridors (see Figure 20) which do not overlap. We assume then that maximal designated 0-corridors do not overlap. If two distinct designated maximal 1-corridors overlap but the intersection contains no arc or vertex then, as in the type 0 case, we designate different representatives which do not overlap. We may assume then that the intersection contains an arc or vertex. If the intersection contains a vertex then it contains the lateral arcs incident to this vertex. After performing a homotopy if necessary we have two maximal corridors whose intersection contains a compartment. Therefore we may extend each corridor to contain all compartments of the other. Since the original corridors are maximal they must be equivalent, contradicting uniqueness of representatives. If the intersection contains an arc but no vertex then this arc must be the unique lateral arc of a compartment of type 0. The same argument applies in this case. Hence we may assume that maximal designated 1-corridors do not overlap.

Maximal 2-corridors require further analysis. To begin with we shall, from now on, call the 2-corridors defined above *simple* 2-corridors. For ease of reference we shall also refer to corridors of type 0 and 1 as *simple*: so simple j -corridors and j -corridors are the same thing, for $j = 0, 1$. Let $C = (D, \Gamma_D, \mu)$ be a simple 2-corridor and let $(B, \Gamma_B) = (\mu(D), \Gamma \cap \mu(D))$. Then the vertices of Γ_B together with those arcs of Γ whose closure is contained in B map to a subgraph $\bar{C}$ of $\bar{\Gamma}$ on Σ , which we call the *graph* of C . The images of lateral and internal-lateral arcs of C are called *internal* and *internal-lateral* edges, respectively, of $\bar{C}$. Let $C' = (D', \Gamma_{D'}, \mu')$ be a




Figure 21:

simple 2-corridor. Then C is a sub-corridor of C' if and only if $\bar{C} \subset \bar{C}'$. Moreover, if $\mu(D) \cap \mu'(D')$ contains an arc a then every arc equivalent to a also belongs to $\mu(D) \cap \mu'(D')$. Hence $\mu(D) \cap \mu'(D')$ contains an arc if and only if $\bar{C} \cap \bar{C}'$ contains an edge. Suppose now that C and C' both bind boundary intervals P_0 and P_1 . If $\bar{C} \subset \bar{C}'$ and both C and C' have width 1 then clearly C and C' are equivalent. Suppose next that $\bar{C} \subset \bar{C}'$, the width of C is 1 and the width of C' is at least 2. Assume further that the graph $\bar{C}$ meets P_i at a point x_i , let e_i be the edge of $\bar{C}$ incident to x_i and let u_i be the vertex of $\bar{C}$ incident to e_i , for $i = 0, 1$. (See Figure 21.) As $\bar{C} \subset \bar{C}'$ the vertices u_i also belong to $\bar{C}'$. There are two possibilities.

- (i) The internal-lateral edge of $\bar{C}$ joining u_0 and u_1 is an internal-lateral edge of $\bar{C}'$. In this case C is equivalent to a compartment of C' .
- (ii) The internal-lateral edge of $\bar{C}$ joining u_0 and u_1 is not an internal-lateral edge of $\bar{C}'$. Therefore either u_0 or u_1 is joined to a vertex of $\bar{C}'$ by an internal-lateral edge of $\bar{C}'$ which is not an internal-lateral edge of $\bar{C}$. Suppose that u_0 is joined to vertex v_1 of $\bar{C}'$ by an internal-lateral edge of $\bar{C}'$ which is the image of a class of arcs in a compartment $C'[j]$ of C' . Then u_1 is a vertex of compartment $C'[j-1]$ or $C'[j+1]$ of C' (see Figure 22). In this case C is not equivalent to any compartment of C' . The case u_1 joined to a vertex of $\bar{C}'$ by an internal-lateral edge of $\bar{C}'$ is similar.

In general if $\bar{C} \subset \bar{C}'$ then without loss of generality we may assume that $D = [0, c] \times [0, 1]$ and $D' = [0, c'] \times [0, 1]$, for some integers c and c' . If the compartment $C[0]$ is equivalent to a compartment of C' (as in case (i) above) then C is equivalent to a sub-corridor of C' . If, on the other hand, $C[0]$ is not equivalent to a compartment of C' (as in case (ii) above) then no compartment of C is equivalent to a compartment of C' . (This follows by induction starting with the case (ii) above.) In this case C is a sub-corridor of C' but equivalent to no sub-corridor of C' and we say that C is *diagonal* to C' . By definition, if C has width at least 2 and is diagonal to C' then C' has a sub-corridor which is diagonal to C . For example, consider Figure 23, which shows the graphs of two corridors $C \subset C'$: the edges of $\bar{C}$ are shown solid, whilst those of $\bar{C}' \setminus \bar{C}$ are dashed. The internal-lateral edges of $\bar{C}$ are shown diagonal whilst those of $\bar{C}'$ are vertical. Thus the left hand compartment of C meets the boundary at a and d whilst the right hand compartment meets it at c and e . The sub-corridor of C' between b and c is diagonal to C . We say that




Figure 22:




Figure 23:

simple 2-corridors C_0 and C_1 , both binding boundary intervals P_0 and P_1 and both of width c , are *diagonally-equivalent* if there is a sub-corridor of C_i of width $c-1$ which is diagonal to C_{1-i} , for $i = 0, 1$.

If C_2 is a corridor of width 2 then there is a unique equivalence class of compartments diagonal to C_2 . This is a consequence of condition (iv) for sums of pictures of type 2. It follows that if C_0 is a corridor diagonally-equivalent to a corridor C_1 and if C_1 is diagonally-equivalent to a corridor C_2 then C_0 and C_2 are equivalent. In particular if C_0 and C_1 are corridors diagonally-equivalent to a corridor C_3 then C_0 and C_1 are equivalent.

Lemma 7.3. *Let C and C' be maximal simple 2-corridors.*

- (i) *If a simple sub-corridor of C is equivalent to a sub-corridor of C' then C and C' are equivalent.*
- (ii) *If a simple sub-corridor of C is diagonal to C' then C and C' are diagonally-equivalent.*

Proof.

- (i) If some sub-corridor of C is equivalent to a sub-corridor of C' then there is a corridor C'' containing all compartments of C and C' . Then both C

and C' are sub-corridors of C'' and, as C and C' are maximal, C'' is a sub-corridor of them both. Hence C and C' are equivalent to C'' .

- (ii) Assume that some sub-corridor C_s of C is diagonal to C' . Choose $p < q$ such that $C_s = C[p, q]$ is maximal in this respect and write $C_s = (D_s, \Gamma_s, \mu_s)$. As C_s is maximal among sub-corridors of C diagonal to C' , either there is a vertex of $\mu(\Gamma_D(p-1))$ which is not a vertex of $\mu'(\Gamma_{D'})$ or $p = 0$. Similarly, either there is a vertex of $\mu(\Gamma_D(q))$ which is not a vertex of $\mu'(\Gamma_{D'})$ or $q = c$. If both $p = 0$ and $q = c$ then C is a sub-corridor of C' and the width of C' is necessarily greater than that of C , contrary to maximality of C . Thus either $p > 0$ or $q < c$. Assume first that $p > 0$ and that there is a vertex of $\mu(\Gamma_D(p-1))$ which is not a vertex of $\mu'(\Gamma_{D'})$. Since the vertices of $\mu(\Gamma_D(p))$ belong to different compartments of C' , it follows that one of the vertices of $\mu(\Gamma_D(p-1))$ belongs to a compartment of C' (and is joined to a vertex of $\mu(\Gamma_D(p))$ by an internal-lateral arc of C'). Hence one vertex of $\mu(\Gamma_D(p-1))$ is contained in $\mu'(\Gamma_{D'})$. If $p \geq 2$ it is therefore possible to extend C' to a wider corridor by adjoining one vertex of $\mu(\Gamma_D(p-1))$ and one vertex of $\mu(\Gamma_D(p-2))$. This is contrary to the maximality of C' and so if $p > 0$ then $p = 1$. A similar argument shows that if $q < c$ then $q = c - 1$. Now if $p = 1$ and $q = c - 1$ then it follows that C' is a sub-corridor of C , so C and C' are equivalent, contrary to hypothesis. Hence either $p = 1, q = c$ or $p = 0, q = c - 1$. It follows that the width of C_s is one less than that of C and that the compartment of C not in C_s has precisely one vertex in C' . As C' has a sub-corridor diagonal to C the same holds on reversing the roles of C and C' . Thus C and C' are diagonally-equivalent.

From the proof of the second part of the above Lemma it follows that if C and C' are diagonally equivalent then $\bar{C}$ contains exactly one vertex and one lateral edge not in $\bar{C}$, and vice-versa.

Given that $C = (D, \Gamma_D, \mu)$ and $C' = (D', \Gamma_{D'}, \mu')$ are diagonally-equivalent simple 2-corridors we define the sum $C + C'$ to be $(D \cup D', \Gamma_D \cup \Gamma_{D'}, \mu \cup \mu')$ and call this an *extended* 2-corridor, *extending* C and C' . The set of *lateral* arcs of an extended 2-corridor $C + C'$ is the union $\{\text{lateral arcs of } C\} \cup \{\text{lateral arcs of } C'\}$. We define the set of *sub-corridors* of the extended 2-corridor $E = C + C'$ to be the union of $\{E\}$ with the sets of sub-corridors of C and C' . (We shall not need to consider extended corridors as sub-corridors of other extended corridors.) If $E = C + C'$ is an extended 2-corridor then we define the *graph* of E to be $\bar{E} = \bar{C} \cup \bar{C}'$ and the *image* of E to be $B \cup B'$, where B and B' are the images of C and C' , respectively. By the remark following the Lemma above $\bar{E}$ has one more lateral edge than both $\bar{C}$ and $\bar{C}'$.

We define the set of 2-corridors to be the union of the set of all simple 2-corridors and the set of all extended 2-corridors. We extend the equivalence relation defined on simple 2-corridors by defining 2-corridors C and C' to be *equivalent* if they have the same lateral arcs. As graphs of simple 2-corridors have an even number of lateral edges, graphs of extended 2-corridors have an odd number of lateral edges and so no simple 2-corridor can be equivalent to an extended 2-corridor. A 2-corridor C is *maximal* in the set of 2-corridors if C is equivalent to C' , for all 2-corridors C' such that $\{\text{lateral edges of } C\} \subset \{\text{lateral edges of } C'\}$. If E and

E' are 2-corridors with images B and B' , respectively, then we say that E and E' have *intersection* $B \cap B'$ and that they *overlap* if $B \cap B' \neq \emptyset$.

Lemma 7.4. *Let $E = C_0 + C_1$ be an extended 2-corridor where C_0 and C_1 are diagonally-equivalent simple 2-corridors. Then the following hold.*

- (i) *If E is maximal as a 2-corridor then C_0 and C_1 are maximal simple 2-corridors.*
- (ii) *if $E = C_0 + C'_1$ then C_1 is equivalent to C'_1 .*

Proof. Let C_0 and C_1 have width c .

- (i) Let C'_i be a maximal simple 2-corridor containing C_i , $i = 0, 1$. Then C_0 is a sub-corridor of C'_0 diagonal to C'_1 . From Lemma 7.3(ii), C'_0 and C'_1 are diagonally-equivalent. Hence we may form the sum $C'_0 + C'_1$. As E is maximal it is equivalent to $C'_0 + C'_1$. If e is a lateral edge of $\bar{C}'_0 \setminus \bar{C}_0$ this implies that e is a lateral edge of $\bar{C}'_1 \setminus \bar{C}_1$. As C_0 and C_1 are diagonally equivalent there is only one such edge. However, if $\bar{C}'_0$ is wider than $\bar{C}_0$ then it must contain at least 2 more lateral edges than $\bar{C}_0$. Hence C'_0 and C_0 are equivalent. A similar argument applies to C_1 .
- (ii) There is a unique vertex v of $\bar{E}$ which belongs to $\bar{C}_1$ but not $\bar{C}_0$. Therefore v is the unique vertex of $\bar{E}$ which belongs to $\bar{C}'_1$ but not $\bar{C}_0$. Hence C_1 and C'_1 must have the same vertices which implies that they are equivalent.

Lemma 7.5. *Let E and E' be maximal 2-corridors.*

- (i) *If a simple sub-corridor of E is equivalent to a simple sub-corridor of E' or*
- (ii) *a simple sub-corridor of E is diagonal to a simple sub-corridor of E'*

then E and E' are equivalent 2-corridors.

Proof. If E and E' are simple 2-corridors then, from Lemma 7.3, (i) follows immediately and (ii) is not possible. If E is a simple 2-corridor and E' an extended 2-corridor then it follows from Lemma 7.4 and Lemma 7.3 with either (i) or (ii), that E' is equivalent to $E + C'$, where C' is a simple sub-corridor of E' , contrary to maximality of E . If E' is a simple 2-corridor and E an extended 2-corridor then again a contradiction arises. Hence we assume that both E and E' are extended 2-corridors, say $E = C_0 + C_1$ and $E' = C'_0 + C'_1$.

- (i) If a simple sub-corridor of E is equivalent to a simple sub-corridor of E' then, for some $i, j \in \{0, 1\}$ a sub-corridor of C_i is equivalent to a sub-corridor of C'_j . From Lemmas 7.4(i) and 7.3(i) it follows that C_i is equivalent to C'_j . Thus $E = C_i + C_{1-i}$ is equivalent to $C'_j + C_{1-i}$. Hence C_{1-i} is diagonally-equivalent to C'_j which is diagonally-equivalent to C'_{1-j} . It follows that C_{1-i} and C'_{1-j} are equivalent and therefore so are E and E' .
- (ii) If a simple sub-corridor of E is diagonal to a simple sub-corridor of E' then, for some $i, j \in \{0, 1\}$ a sub-corridor of C_i is diagonal to C'_j . From Lemma 7.4(i) and Lemma 7.3(ii) it follows that C_i and C'_j are diagonally-equivalent. As C_i is diagonally-equivalent to C_{1-i} it follows that C'_j is equivalent to C_{1-i} . As in the previous case it follows that E and E' are equivalent.

Lemma 7.6. *Let $E = C_0 + C_1$ and $E' = C'_0 + C'_1$ be maximal extended 2-corridors. Then either E and E' are equivalent or $\bar{E} \cap \bar{E}' \subset \bar{C}_i \cap \bar{C}'_j$, for some $i, j \in \{0, 1\}$.*

Proof. If $\bar{E} \cap \bar{E}' = \emptyset$ then the result holds trivially. Suppose then that E and E' are not equivalent and that $\bar{E} \cap \bar{E}' \neq \emptyset$. Note that this implies that the images of E and E' are disks (and not annuli or Möbius bands). There is a unique vertex u_0 of $\bar{C}_0$ not in $\bar{C}_1$ and a unique vertex u_1 of $\bar{C}_1$ not in $\bar{C}_0$. Similarly, there is a unique vertex v_0 of $\bar{C}'_0$ not in $\bar{C}'_1$ and a unique vertex v_1 of $\bar{C}'_1$ not in $\bar{C}'_0$. Fix $j = 0$ or 1 : if the vertex v_j is not in $\bar{E}$ then $\bar{E} \cap \bar{E}' \subset \bar{E} \cap \bar{C}'_{1-j}$. Now if u_i is not in $\bar{C}'_{1-j}$, for $i = 0, 1$, then $\bar{E} \cap \bar{E}' \subset \bar{C}_{1-i} \cap \bar{C}'_{1-j}$. On the other hand if $u_i \in \bar{C}'_{1-j}$, for $i = 0$ and 1 , then $\bar{C}'_{1-j}$ contains $\bar{E}$, contrary to maximality of E . Finally suppose that v_j belongs to $\bar{E}$, for $j = 0$ and 1 . As $\bar{E}$ and $\bar{E}'$ are both graphs of extended 2-corridors on a disk this implies that $\bar{E}'$ is a subgraph of $\bar{E}$. As E and E' are not equivalent it follows that $\bar{E}'$ is a proper subgraph of $\bar{E}$ and so E' is not maximal, a contradiction.

Definition 7.7. Choose one representative E from each equivalence class of maximal 2-corridors of Γ . If E is the representative of an equivalence class of maximal extended 2-corridors then we call E a *designated* extended 2-corridor. Corresponding to each designated extended 2-corridor E choose an ordered pair (C, C') of diagonally-equivalent simple 2-corridors such that $E = C + C'$ and say that E has *foundation* C and *extension* C' . If C is the chosen representative of an equivalence class of maximal simple 2-corridors or C is the foundation of a designated extended 2-corridor then we call C and all its induced sub-corridors *designated* simple 2-corridors. We say that a 2-corridor is *designated* if it is either a designated extended 2-corridor or a designated simple 2-corridor.

We assume from now on that all 2-corridors are designated. In particular if C_s is a proper sub-corridor of the extended 2-corridor E with foundation C then, by definition, C_s is a sub-corridor induced from C . (Note that if E is a designated extended 2-corridor with foundation C and extension C' then C is a designated simple 2-corridor but C' is not.)

Suppose now that $E = C_0 + C_1$ and $E' = C'_0 + C'_1$ are maximal designated 2-corridors, with foundations C_0 and C'_0 , respectively. Assume further that E and E' are not equivalent. If E and E' overlap then we may assume that the images of E and E' are not annuli or Möbius bands and, as in the case of simple 2-corridors, that the graphs $\bar{E}$ and $\bar{E}'$ have at least an edge in common. Furthermore, from Lemma 7.5, no sub-corridor of E is equivalent or diagonal to a sub-corridor of E' and, from Lemma 7.6, $\bar{E} \cap \bar{E}' \subset \bar{C}_i \cap \bar{C}'_j$, for some $i, j \in \{0, 1\}$. It follows that no compartment of C_i is equivalent or diagonal to a sub-corridor of C'_j , and vice-versa. Therefore $\bar{C}_i \cap \bar{C}'_j$ contains a unique edge. Hence the intersection of E and E' contains one vertex and one class of lateral arcs (corresponding to the edge of $\bar{C}_i \cap \bar{C}'_j$). Therefore the union of the images of E and E' form a configuration of the form shown in Figure 24: in which case we say E and E' form a *keel*.

We collect together our conclusions on the intersection of corridors in the following Lemma.

Lemma 7.8. *If C and C' are maximal designated j -corridors, $j = 0, 1$ or 2 , then either C and C' do not overlap or $j = 2$ and C and C' form a keel.*

From now on all corridors are assumed to be designated.




Figure 24: The unique edge of the intersection is shown as a dashed line

7.1. Z-cancellation. Let $C = (D, \Gamma_D, \mu)$ be a j -corridor, where $j = 0, 1$ or 2 and $D = [0, c] \times [0, 1]$, for some integer c . Recall that Γ has boundary partition $\mathbf{b} = (b_1, \dots, b_{W_1})$ and suppose that C binds boundary intervals $P_0 = b_{p_0}$ and $P_1 = b_{p_1}$, where $1 \leq p_1 \leq W_1$, that P_i has prime label (h_i, f_i) and that $\alpha(f_i) = m_i$: so the label of P_i is $h_i \wedge m_i$. Taking the point $\mu((0, i))$ to precede $\mu((c, i))$ gives an orientation ζ_i of P_i . If ζ_i coincides with the fixed orientation $\zeta(P_i)$ of P_i set $\varepsilon_i = 1$ and if not set $\varepsilon_i = -1$. Fix an integer s with $0 \leq s \leq c - 1$ and assume that $\Gamma_D(s)$ has base $D_s[x_0, x_1, y_0, y_1]$. The label w_i of the sub-interval of P_i , between the point $\mu((s + \frac{1}{x_i+1}, i))$ and the end of the interval P_i , read in the direction of ζ_i , therefore begins with a terminal segment of the word $h_i^{\varepsilon_i}$ of uniquely determined length r_i such that $0 \leq r_i < l(h_i)$. We define the *left boundary marking of rank s of C* to be the sequence

$$p_0, p_1, \varepsilon_0, \varepsilon_1, r_0, r_1.$$

The *left boundary marking of rank s of an extended 2-corridor E* is defined to be the left boundary marking of rank s of its foundation.

Now assume that C above is a maximal simple (designated) j -corridor. Let $0 < p < q < c$ and consider the sub-corridor $C[p, q]$ of C of width $p - q \leq c - 1$. Suppose that the left boundary marking $p_0, p_1, \varepsilon_0, \varepsilon_1, r_0, r_1$ of rank p of C is equal to the left boundary marking of rank q of C . Let $\Gamma_D(p)$ have base $D_p(x_0, x_1, y_0, y_1)$ and $\Gamma_D(q)$ have base $D_q(x'_0, x'_1, y'_0, y'_1)$ and let I_i be the sub-interval

$$I_i = \left[p + \frac{1}{x_i + 1}, q + \frac{1}{x'_i + 1} \right] \times \{i\}$$

of $[0, c] \times \{i\}$, for $i = 0, 1$ (see Figure 25). Then the label of $\mu(I_i)$ read in the direction ζ_i is a cyclic permutation of $(h^{\varepsilon_i})^{\alpha_i}$, for some positive integer α_i . Assume further that the left interior markings of $\Gamma_D(p)$ and $\Gamma_D(q)$ are equal (so in particular $y_0 = y'_0$). Under these circumstances we form a new picture Γ' from Γ as follows. Let $T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ be the translation defined by

$$(7.1) \quad T(x, y) = (x - (q - p), y), \text{ for } (x, y) \in \mathbb{R}^2.$$

Let S be a homeomorphism $S : D_p \longrightarrow [p, q + 1] \times [0, 1]$ such that S fixes

$$\{p\} \times [0, 1] \cup [p, p + \frac{1}{x_0 + 1}] \times \{0\} \cup [p, p + \frac{1}{x_1 + 1}] \times \{1\}$$




Figure 25:

and $S((p+1, t)) = (q+1, t)$, for all $t \in [0, 1]$. Define $D^l = [0, p] \times [0, 1]$ and $D^r = [p+1, c - (q-p)] \times [0, 1]$. Consider the triple $C' = (D', \Gamma_{D'}, \mu')$ where $D' = [0, c - (q-p)] \times [0, 1]$,

$$\Gamma_{D'}(i) = \begin{cases} \Gamma_D(i) & \text{if } 0 \leq i \leq p-1 \\ T(\Gamma_D(i + (q-p))) & \text{if } p \leq i \leq c - (q-p) - 1 \end{cases},$$

$\mu'|_{D^l} = \mu|_{D^l}$, $\mu'|_{D^r} = (\mu \circ T^{-1})|_{D^r}$ and $\mu'|_{D_p} = (\mu \circ S)|_{D_p}$. Then $\Gamma_{D'}$ is a sum of square-pictures with left-trivial labelling on the disk D' . Remove $\mu(\Gamma_D)$ from Γ and replace it with $\mu'(\Gamma_{D'})$. Under the given hypotheses this results in a picture Γ' over G on Σ . The triple C' is, by construction, a maximal simple corridor of Γ' . Furthermore $\mu([p, q+1] \times [0, 1]) = \mu'([p, p+1] \times [0, 1])$ and $\Gamma \cap (\Sigma \setminus \mu([p, q+1] \times [0, 1])) = \Gamma' \cap (\Sigma \setminus \mu'([p, p+1] \times [0, 1]))$. Hence all corridors of Γ , except C are also corridors of Γ' . We designate C' as the representative for its equivalence class of maximal corridors in G' .

Now assume that C is the foundation of a designated maximal extended 2-corridor E with extension C_e . For example the graphs $\bar{C}$ and $\bar{C}_e$ may be as shown in Figure 26, where the vertical edges $e_1, \dots, e_n$ are composed of internal-lateral arcs of compartments of C and the diagonal edges $d_1, \dots, d_n$ are composed of internal-lateral arcs of compartments of C_e . Thus u is the vertex of C not in C_e and v is the vertex of C_e not in C .

Cancellation from p to q along C results in corridors C' and C'_e which have graphs $\bar{C}'$ and $\bar{C}'_e$, as shown in Figure 27. The compartments of C_e containing a' and all those to its left become compartments of C'_e , as does the compartment of C_e containing b' and all those to its right. In addition C'_e has one compartment containing a and q' which replaces all the compartments of C_e between those containing a' and b' . Therefore the only compartment of C'_e which is not a compartment of C_e is




Figure 26:




Figure 27:

that containing vertices a and q' . Note that in C the left interior markings of the compartments containing p and q are equal. It follows that the left interior markings of the compartments of C_e and C'_e containing a are equal. From the graph it is clear that C' and C'_e are diagonal and, since C and C_e are maximal so are C' and C'_e . Thus $E' = C' + C'_e$ is a maximal extended 2-corridor. We choose E' as the designated representative of its class in Γ' and C' and C'_e as its foundation and extension, respectively.

The formation of Γ' from Γ along a corridor C in this way is called *Z-cancellation along C from p to q* . If C is of type j we refer to $Z(j)$ -cancellation along C , $j = 0, 1, 2$. We remark that if Γ' is formed from Γ by Z -cancellation then the left interior and left boundary markings of compartments of corridors of Γ' are left interior and left boundary markings of compartments of corridors of Γ . Also, the boundary partition $\mathbf{b}$ is unaffected by Z -cancellation, the prime labels of boundary intervals are the same in Γ' as in Γ and for $j \neq p_i$ the boundary interval b_j has the same label in Γ' as in Γ . With the above notation, the label of $P_i = b_{p_i}$ in Γ' is $h_i \wedge (m_i - l(h_i)\alpha_i)$, for $i = 0, 1$.

Theorem 7.9. *If C is a j -corridor, for $j = 0, 1$ or 2 and C has more than $M(j)$ arcs then Z -cancellation is possible along some sub-corridor of C . ($M(j)$ is defined in (3.18).)*

Proof. Assume first that C is a simple corridor of width c , $C = (D, \Gamma_D, \mu)$. Then Z -cancellation is possible if there are integers q_0, q_1 , with $0 < q_0 < q_1 < c$, such that the left interior marking of C of rank q_0 equals the left interior marking of C of rank q_1 and the boundary marking of C of rank q_0 equals the boundary marking of C of rank q_1 .

Let $p_0, p_1, \varepsilon_0, \varepsilon_1, r_0, r_1$ be the left boundary marking of rank q_0 and let (h_{p_i}, m_{p_i}) be the prime label of the boundary interval b_{p_i} , for $i = 0, 1$. Since p_i and ε_i are determined by C then, for all q_1 with $0 < q_1 < c$, the left boundary marking of rank q_1 is $p_0, p_1, \varepsilon_0, \varepsilon_1, s_0, s_1$, for some integers s_0 and s_1 . By definition $0 \leq r_i, s_i < l(h_i) \leq W_2(\mathbf{z})$. Hence there are at most W_2^2 distinct left boundary markings of C . If $j = 0$ the left interior markings of all ranks of C are identical and the number of arcs of C is c . If $c > W_2^2 + 1 = M(0)$ then from the above it follows that Z -cancellation along C is possible, as claimed.

If $j = 1$ then the left interior marking of rank s of C is $\delta, o, \rho, \sigma, 0, 0, 0, 0$, where $\delta \in \{-1, 0, 1\}$, $o, \sigma \in \{0, \dots, l(r) - 1\}$, $0 \leq \rho < m$ and $\delta = o = \rho = \sigma = 0$ if and only if $\Gamma_D(s)$ is of type square-0. Therefore there are at most $3ml(r)^2$ distinct left interior markings of C . It follows that if C has width $c > W_2^2 \cdot 3ml(r)^2 + 1$ then Z -cancellation along C is possible. A square-picture of type 0 has 1 arc while a square-picture of type 1 has $ml(r)$ arcs. Hence if C has more than $(W_2^2 \cdot 3ml(r)^2 + 1)ml(r) \leq M(1)$ arcs then Z -cancellation along C is possible.

If $j = 2$ then the left interior marking of rank s of C is $\delta_0, o_0, \rho_0, \sigma_0, \delta_1, o_1, \rho_1, \sigma_1$, where $\delta_i = \pm 1$, $o_i, \sigma_i \in \{0, \dots, l(r) - 1\}$ and $0 \leq \rho_i < m$. Therefore there are at most $4m^2l(r)^4$ distinct left interior markings of C . It follows that if C has width $c > W_2^2 \cdot 4m^2l(r)^4 + 1$ then Z -cancellation along C is possible. A square-picture of type 2 has less than $2ml(r)$ arcs so if C has more than $(W_2^2 \cdot 4m^2l(r)^4 + 1)ml(r)$ arcs then Z -cancellation along C is possible.

If C is not simple then $j = 2$ and C has foundation a simple 2-corridor C^f . Then C has at most $ml(r)$ more arcs than C^f . Thus if C has more than $(W_2^2 \cdot 4m^2l(r)^4 + 2)ml(r) = M(2)$ arcs then, from the above Z -cancellation is possible along C^f and therefore along C , as claimed.

From this Theorem it follows that a Z -cancellation along a j -corridor may be achieved by a sequence of Z -cancellations along sub-corridors of width at most $M(j)$.

Definition 7.10. A picture on which no Z -cancellation is possible is called *Z -reduced*.

Corollary 7.11. If C is a j -corridor of a Z -reduced picture then C has at most $M(j)$ arcs.

Lemma 7.12. Let Γ be a picture on a surface Σ of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with prime labels $\mathbf{z}$, labelled by $\hat{\alpha}(\mathbf{z})$ and with boundary partition $\mathbf{b} = (b_1, \dots, b_{W_1})$. Assume that Γ' is obtained from Γ by Z -cancellation. Then Γ' is reduced if and only if Γ is reduced. Furthermore Γ' is reduced and minimalistic if and only if Γ is reduced and minimalistic.

Proof. Suppose first that Γ is reduced. If Γ' has a pair of cancelling vertices u, v then these must be adjacent in Γ' but not in Γ . Hence u, v belong to consecutive compartments of some corridor $C = (D, \Gamma_D, \mu)$ of Γ' which results from Z -cancellation in Γ . It follows that C must have type $j = 1$ or 2 . As u and v cancel in Γ' they are the images of vertices which cancel in Γ_D ; so we restrict attention to Γ_D and assume u and v belong to Γ_D . Suppose u belongs to $\Gamma_D(p)$, v belongs to $\Gamma_D(p+1)$ and that D_p and D_{p+1} have bases $D_p(x_0, x_1, y_0, y_1)$ and $D_p(x'_0, x'_1, y_1, y'_1)$, respectively. If $j = 1$ then it must be possible to perform a bridge move on arcs joining u and v to $[p, p+1] \times \{i\}$, for $i = 0$ or 1 . Assume that this occurs for $i = 0$ and let a_0 and

a_1 be the arcs joining u to $q_0 = (p + 1 - \frac{x_0}{1+x_0}, 0)$ and v to $q_1 = (p + 1 + \frac{1}{1+x_0}, 0)$, respectively: then a bridge move is possible transforming a_0 and a_1 into arcs b_0 , joining u to v , and b_1 with end points q_0 and q_1 . However this means that the interval $[q_0, q_1]$ of ∂D has trivial label, contrary to the standing hypothesis that $\mathbf{z}$ is special. Hence if Γ_D is of type 1 no such cancellation is possible.

Now assume that $j = 2$. If u and v are both incident to $[p, p+2] \times \{i\}$ then, as in the case $j = 1$, no bridge move is possible on arcs joining u and v to ∂D . Hence u and v do not cancel. We assume therefore that u is incident to $[p, p+1] \times \{0\}$ and that v is incident to $[p+1, p+2] \times \{1\}$. However no bridge moves are possible now because both regions incident to the edge of $\bar{\Gamma}_D$ joining u to v are triangular: a bridge move on the sides of a triangle would result in a trivial label on the corner of the triangle opposite the edge joining u to v . Thus no cancellation is possible if $j = 2$. It follows that if Γ is reduced then Γ' is reduced. A similar argument shows that the converse also holds.

Now suppose that Γ is reduced and minimalistic. Then Γ' is reduced. Regions involved in Z -cancellation all have Euler characteristic 1 and both Γ and Γ' are pictures on the surface Σ , so the condition on Euler characteristic of regions is satisfied by Γ' . No arc of Γ' is a closed curve since no such arcs are created in the process of Z -cancellation. Suppose that Γ' has interior edge of width greater than $l - 2$. Since Γ' is reduced it follows from Lemma 6.12 that both ends of this edge are incident to the same vertex. No such edge results from the process of Z -cancellation so this edge also belongs to Γ , which cannot therefore be minimalistic. It follows that Γ' is reduced and minimalistic. The converse follows similarly.

7.2. Corridor-sections. Let Γ_D be a sum of square-pictures of type j on the disk $D = [0, c] \times [0, 1]$ and let w_i denote the label of $[0, c] \times \{i\}$, for $i = 0, 1$, $j = 0, 1$ or 2. Assume that the following conditions hold.

- (i) There exist integers $p_0, p_1, \varepsilon_0, \varepsilon_1, t_0$ and t_1 such that $1 \leq p_i \leq W_1$, the p_i th letter (h_i, f_i) of $\mathbf{z}$ is not a degenerate letter or a minor letter of H^Λ , $\varepsilon_i = \pm 1$, $0 \leq t_i < l(h_i)$ and $w_i = g_i^{\alpha_i}$, for some $\alpha_i > 0$, and $g_i = \tau(h_i^{\varepsilon_i}, t_i) \iota(h_i^{\varepsilon_i}, l(h_i) - t_i)$, for $i = 0, 1$.
- (ii) Γ_D has at most $M(j)$ arcs.
- (iii) Let D^e be the disk $[c, c+1] \times [0, 1]$. Let T_c be the translation $T(x, y) = (x + c, y)$, for $(x, y) \in \mathbb{R}^2$, and let $\Gamma_{D^e} = T_c(\Gamma_D(0))$, a picture of type square- j on D^e . Then $\Gamma_D \cup \Gamma_{D^e}$ is a sum of square-pictures on the disk $[0, c+1] \times [0, 1]$.

Then $(\Gamma_D, p_0, p_1, \varepsilon_0, \varepsilon_1, t_0, t_1)$ is called a *corridor-section* on Γ_D , of width c , binding the p_0 th and p_1 th letters of $\mathbf{z}$, and of *type j* if Γ_D is a sum of square-pictures of type j .

The corridor-section $(\Gamma_D, p_0, p_1, \varepsilon_0, \varepsilon_1, t_0, t_1)$ is said to have *left interior marking* equal to the left interior marking of Γ_D , *left boundary marking* the sequence

$$p_0, p_1, \varepsilon_0, \varepsilon_1, t_0, t_1$$

and *extent* the pair (α_0, α_1) .

Let C be a corridor section of width c . The conditions on a corridor-sections imply that if $0 < p < c - 1$ then a new corridor-section C_p may be formed by cutting C along $\{p\} \times [0, 1]$ and then identifying $(0, t)$ with (c, t) , for all $t \in [0, 1]$. We set $C = C_0$. The left interior and left boundary markings of C_p are called the *left interior* and *left boundary* markings, respectively, of C of *rank p* . Thus the left

interior marking and left boundary marking of C coincide with the left interior and boundary markings of rank 0 of C .

Let L denote the set of all corridor sections. As the number of arcs of a corridor section is bounded L is finite. Let L have $|L|$ elements indexed by the integers $1, \dots, |L|$ and denote these elements $L(1), \dots, L(|L|)$. For $i = 1, \dots, |L|$, define the quintuple

$$L_5(i) = (L(i), s(i), t(i), \alpha(i, 0), \alpha(i, 1))$$

where $s(i)$ and $t(i)$ are the left boundary marking and left interior marking, respectively, of $L(i)$ and $(\alpha(i, 0), \alpha(i, 1))$ is the extent of $L(i)$. Furthermore write

$$s(i) = p(i, 0), p(i, 1), \varepsilon(i, 0), \varepsilon(i, 1), r(i, 0), r(i, 1)$$

for the left boundary marking of $L(i)$. For fixed i with $1 \leq i \leq |L|$ and each integer p with $1 \leq p \leq W_1$ define

$$\zeta(p, i) = \begin{cases} 0, & \text{if } p \neq p(i, 0) \text{ and } p \neq p(i, 1) \\ \alpha(i, 0), & \text{if } p = p(i, 0) \text{ and } p \neq p(i, 1) \\ \alpha(i, 1), & \text{if } p \neq p(i, 0) \text{ and } p = p(i, 1) \\ \alpha(i, 0) + \alpha(i, 1), & \text{if } p = p(i, 0) = p(i, 1) \end{cases}.$$

The following lemma follows directly from the definitions above and the remarks preceding Theorem 7.9.

Lemma 7.13. *Let C be a j -corridor of Γ and let $C[p, q] = (D, \Gamma_D, \mu)$ be a sub-corridor (of width $q - p$) with at most $M(j)$ arcs. If Z -cancellation takes place along C from p to q and the left boundary marking of C of rank p is t then (Γ_D, t) is a corridor-section of width $q - p$ and type j . Let $(\Gamma_D, t) = L(i)$, for some i with $1 \leq i \leq |L|$, let Γ' be the picture formed by this Z -cancellation and let the boundary interval b_p of Γ have label $h_p \wedge m_p$, for $p = 1, \dots, W_1$. Then the boundary interval of b_p in Γ' has label*

$$h_p \wedge (m_p - l(h_p)\zeta(p, i)),$$

for $p = 1, \dots, W_1$.

The Z -cancellation of Lemma 7.13, along C from p to q , where the left boundary marking, of rank p , of C is t is called Z -cancellation of corridor-section (Γ_D, t) . As remarked (above Definition 7.10) any Z -cancellation may be effected by a sequence of Z -cancellations of corridor-sections.

7.3. Z -insertion. Let $C = (D, \Gamma_D, \mu)$ be a maximal simple j -corridor, with $D = [0, c] \times [0, 1]$, binding b_{p_0} and b_{p_1} , for p_i with $1 \leq p_i \leq W_1$. Let the left interior marking and left boundary marking of C , of rank p , be s and t , respectively. Let (Γ_Δ, t) be a corridor section with left interior marking s , where $\Delta = [0, q] \times [0, 1]$. Let $T_p : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $T_q : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the translations defined by $T_p(x, y) = (x + p, y)$ and $T_q(x, y) = (x + q, y)$, for $(x, y) \in \mathbb{R}^2$ and positive integers p and q . Let D_p have base $D_p(x_0, x_1, y_0, y_1)$ and Δ_0 have base $\Delta_0(x'_0, x'_1, y'_0, y'_1)$. Let S be a homeomorphism $S : [p, p + q + 1] \times [0, 1] \rightarrow [p, p + 1] \times [0, 1]$ such that S fixes $\{p\} \times [0, 1]$,

$$S([p, p + \frac{1}{x'_i + 1}] \times \{i\}) = [p, p + \frac{1}{x_i + 1}] \times \{i\},$$

for $i = 0, 1$, and $S((p + q + 1, t)) = (p + 1, t)$, for all $t \in [0, 1]$. Define $D^l = [0, p] \times [0, 1]$, $D^c = [p, p + q + 1] \times [0, 1]$ and $D^r = [p + q + 1, c + q] \times [0, 1]$.

Consider the triple $C_I = (D_I, \Gamma_{D_I}, \mu_I)$, where $D_I = [0, c+q] \times [0, 1]$,

$$\Gamma_{D_I}(i) = \begin{cases} \Gamma_D(i) & \text{for } i = 0, \dots, p-1 \\ T_p(\Gamma_\Delta(i-p)) & \text{for } i = p, \dots, p+q-1 \\ T_q(\Gamma_D(i-q)) & \text{for } i = p+q, \dots, c+q-1 \end{cases},$$

$$\mu_I|_{D^l} = \mu|_{D^l}, \mu_I|_{D^c} = (\mu \circ S)|_{D^c} \text{ and } \mu_I|_{D^r} = (\mu \circ T_q^{-1})|_{D^r}.$$

Then Γ_{D_I} is a sum of square-pictures with left-trivial labelling on the disk D_I . Remove $\mu(\Gamma_D)$ from Γ and replace it with $\mu_I(\Gamma_{D_I})$. This results on a picture Γ_I over G on Σ . The triple C_I is, by construction, a maximal corridor of Γ_I . Furthermore $\mu([p, p+1] \times [0, 1]) = \mu_I([p, p+q+1] \times [0, 1])$ and $\Gamma \cap (\Sigma \setminus \mu([p, p+1] \times [0, 1])) = \Gamma_I \cap (\Sigma \setminus \mu_I([p, p+q+1] \times [0, 1]))$. Hence all corridors of Γ , except C , are also corridors of Γ_I . We designate C_I as the representative for its equivalence class of maximal corridors in Γ_I . Moreover if C is the foundation of a designated maximal extended 2-corridor of Γ then C_I is a maximal simple sub-corridor of a maximal extended 2-corridor of Γ_I (as can be seen by reversing the argument for cancellation along extended 2-corridors). In this case C_I is chosen as the foundation of the corresponding maximal extended 2-corridor of Γ_I , which becomes a designated 2-corridor of Γ_I .

The formation of Γ_I from Γ in this way is called *Z-insertion of (Γ_Δ, t) into C at position p* . If C is of type j we refer to *Z(j)-insertion*, $j = 0, 1, 2$.

Lemma 7.14. *If Γ_I is formed from Γ by Z-insertion of (Γ_Δ, t) then the left interior and left boundary markings of compartments of corridors of Γ_I are left interior and left boundary markings of square-pictures of Γ_Δ or of compartments of corridors of Γ . Also, the boundary partition $\mathbf{b}$ is unaffected by Z-insertion and, given that the boundary interval b_p of Γ has label $h_p \wedge m_p$, for $p = 1, \dots, W_1$, then the boundary interval of b_p in Γ_I has label*

$$h_p \wedge (m_p + l(h_p)\zeta(p, i)),$$

for $p = 1, \dots, W_1$.

The operations of Z-cancellation and Z-insertion are inverse to each other in an obvious way. To be more precise, let Γ_I be formed from Γ by Z-insertion of (Γ_Δ, t) into C at position $p \geq 1$ and let the resulting corridor of Γ_I be C_I as above. Then Z-cancellation of corridor-section (Γ_Δ, t) along C_I from p to q is possible and results in Γ . A similar statement holds starting with Z-cancellation.

7.4. The Z-graph. Let L denote the set of corridor-sections of $\mathbf{z}$ (indexed by the integers $1, \dots, |L|$ as before) and define the *left marking*, of rank p , of $L(i) \in L$ to be $m(i, p) = s(i, p), t(i, p)$, where $s(i, p)$ and $t(i, p)$ are the left interior and boundary markings, respectively, of rank p of $L(i)$. Define the *left marking* of $L(i)$ to be $m(i) = m(i, 0)$, let

$$m_{L(i)} = \{m(i, p) : 0 \leq p < \text{width of } L(i)\}$$

and let

$$m_{\mathbf{z}} = \bigcup_{i=1}^{|L|} m_{L(i)}.$$

Let C be a corridor of the picture Γ and define the *left marking*, of rank p , of C to be s, t , where s and t are the left interior and boundary markings, respectively,

of rank p , of C . Define the *active* left markings of Γ to be those left markings of corridors of C which are elements of $m_{\mathbf{z}}$.

The Z -graph $G(\mathbf{z})$ of $\mathbf{z}$ is defined as follows. The vertices of $G(\mathbf{z})$ are the subsets of $m_{\mathbf{z}}$. There is one directed edge labelled $L(i)$ from vertex R to vertex T corresponding to each corridor-section $L(i)$ such that $L(i)$ has left marking in R and $T = R \cup m_{L(i)}$. Corresponding to the picture Γ is the vertex

$$v(\Gamma) = \{\text{active left markings of } \Gamma\}$$

of $G(\mathbf{z})$ which we call the *initial* vertex of Γ .

Now, for $i = 0, \dots, n$, let Γ_i be a picture, over G on Σ , such that Γ_{i+1} is obtained from Γ_i by Z -cancellation of a corridor-section C_i along a corridor of Γ_i . Assume that $\Gamma = \Gamma_n$. Then this sequence of Z -cancellations uniquely determines a path $S_n, \dots, S_0$ in $G(\mathbf{z})$, where $S_i \in V(G(\mathbf{z}))$, $S_n = v(\Gamma)$, $S_i = S_{i+1} \cup \{\text{left markings of } C_i\}$ and the edge of the path joining S_{i+1} to S_i is labelled C_i . Then Γ_i is obtained from Γ_{i+1} by Z -insertion of C_i . Hence the set of active left markings of Γ_i is the union of the left markings of C_i with the active left markings of Γ_{i+1} . It follows, by induction, that the set of active left markings of Γ_0 is S_0 .

Conversely, let $S_n, \dots, S_0$ be a directed path in $G(\mathbf{z})$, where $S_n = v(\Gamma)$ and the edge of the path joining S_{i+1} to S_i is labelled C_i . Then $S_i = S_{i+1} \cup \{\text{left markings of } C_i\}$. Let $\Gamma_n = \Gamma$ and let m be an integer with $n \geq m > 0$. Suppose inductively that we have constructed pictures Γ_i with active left markings S_i , for $i = n, \dots, m$ such that Γ_{i+1} is obtained from Γ_i by Z -cancellation of the corridor-section C_i , for $i = n-1, \dots, m$. As C_{m-1} is the label of the edge joining S_m to S_{m-1} it follows that $S_{m-1} = S_m \cup \{\text{left markings of } C_{m-1}\}$ and that the left marking of C_{m-1} is in $S_m = \{\text{active left markings of } \Gamma_m\}$. Hence we may form a picture Γ_{m-1} from Γ_m by Z -insertion of C_{m-1} . The active left markings of Γ_{m-1} are then the left markings of C_{m-1} together with the active left markings of Γ_m : that is S_{m-1} . Moreover as Γ_{m-1} is formed from Γ_m by Z -insertion of C_{m-1} it follows that Γ_m is formed from Γ_{m-1} by Z -cancellation of C_{m-1} . Hence the path $S_n, \dots, S_0$ determines a sequence of Z -cancellations

$$\Gamma_0 \longrightarrow \dots \longrightarrow \Gamma_n = \Gamma,$$

with Γ_{i+1} obtained from Γ_i by Z -cancellation of C_i and the set of active left markings of Γ_0 equal to S_0 .

We have therefore proved the following.

Lemma 7.15. *Let Γ and Γ' be pictures over G on Σ . Then there exists a sequence*

$$\Gamma' = \Gamma_0 \longrightarrow \dots \longrightarrow \Gamma_n = \Gamma,$$

with Γ_{i+1} obtained from Γ_i by Z -cancellation of the corridor-section C_i if and only if there exists a directed path

$$v(\Gamma) = S_n, \dots, S_0 = v(\Gamma')$$

in $G(\mathbf{z})$ such that the edge joining S_{i+1} to S_i is the corridor section C_i .

Definition 7.16. Let $p = S_n, \dots, S_0$ be a directed path in $G(\mathbf{z})$ such that the edge joining S_{i+1} to S_i is the corridor section C_i . Let

$$\Gamma_0 \longrightarrow \dots \longrightarrow \Gamma_n,$$

be a sequence of pictures over G on Σ with Γ_{i+1} obtained from Γ_i by Z -cancellation of the corridor-section C_i . If $v(\Gamma_n) = S_n$ then we say that Γ_0 is the picture obtained from Γ_n by Z -insertion *following* p . If $v(\Gamma_0) = S_0$ then we say that Γ_n is the picture obtained from Γ_0 by Z -cancellation *following* p .

Let Q denote the quadratic exponential equation $Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ and $Q_{\mathcal{H}} = Q(\bar{\mathbf{z}}, \mathcal{H}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ the homogeneous equation corresponding to $\mathbf{z}$. Let the q th letter of $\mathbf{z}$ be (h_q, f_q) , for $1 \leq q \leq W_1$.

Lemma 7.17. *Let $p = S_n, \dots, S_0$ be a directed path in $G(\mathbf{z})$ with initial vertex $v(\Gamma_n)$. Let $e_1, \dots, e_t$ be the edges of $G(\mathbf{z})$ occuring in p , let $L(s_i)$ be the label of e_i and let n_i be the number of occurrences of e_i in p , for $i = 1, \dots, t$. Let Γ' be the picture obtained from Γ_n by Z -insertion *following* p . Then the following hold.*

- (i) *Let α_n be a solution to $\mathcal{H}$ such that (α_n, Γ_n) corresponds to a solution to $Q_{\mathcal{H}}$ and write $m_q = \alpha_n(\lambda_q)$, for $1 \leq q \leq W_1$. The boundary interval b_q of Γ' has label*

$$h_q \wedge \left(m_q + l(h_q) \left(\sum_{i=1}^t n_i \zeta(q, s_i) \right) \right),$$

for $1 \leq q \leq W_1$.

- (ii) *Let α be a solution to $\mathcal{L}$ such that (α, Γ') corresponds to a solution to Q . Let $\alpha_n : M \rightarrow \mathbb{Z}$ be the retraction defined by setting*

$$\alpha_n(\lambda_q) = \alpha(f_q) - l(h_q) \left(\sum_{i=1}^t n_i \zeta(q, s_i) \right),$$

for all $q \in \{1, \dots, W_1\}$ such that the q th letter of $\mathbf{z}$ is a proper exponential H^Λ -letter, and $\alpha_n(\lambda) = 0$, for all other $\lambda \in \Lambda$. Then (α_n, Γ_n) corresponds to a solution of $Q_{\mathcal{H}}$.

Proof. The first part of the Lemma follows from Lemma 7.14. For the second part note that the boundary label of b_q in Γ_n is $h_q \wedge \alpha_n(\lambda_q)$, for all q (using Lemma 7.14 again). By definition of Z -cancellation therefore $\alpha_n(\lambda_q) > 0$. Also $\alpha_n(\lambda_q) \equiv \alpha(f_q) \pmod{l(h_q)}$. It follows that (α_n, Γ_n) corresponds to a solution to $Q_{\mathcal{H}}$.

A picture Γ such that there exists a solution α to $\mathcal{H}$ and (α, Γ) corresponds to a solution to $Q_{\mathcal{H}}$ will be called a *basic* picture for $(\mathbf{z}, \mathcal{L})$.

We call a path in a graph with no repeated vertices a *simple path* and a path with initial and terminal vertices identical but no other pair of vertices identical a *cycle*. A cycle of length 1 is called a *loop*. The *path-subgraph* of a path is the subgraph consisting of all edges and vertices of the path. The directed path p of $G(\mathbf{z})$ corresponding to a sequence of Z -cancellations contains no cycle of length greater than 1 but may contain loops. We call a directed path p in $G(\mathbf{z})$ a *Z -path* if p has no cycle of length greater than 1. The path-graph of a Z -path consists of the path-subgraph of a directed simple path together with a (possibly empty) collection of loops. Given a directed path p assign weight $w_p(e)$ to each edge of $G(\mathbf{z})$, where $w_p(e)$ is the number of times the edge appears in p . We call the edge weighting w_p the *weighting* of the path p . Then there are finitely many directed paths in $G(\mathbf{z})$ which have the same weighting as p . We call a weighting of the edges of $G(\mathbf{z})$ a *Z -weighting* if it equals the weighting of a Z -path. Given a Z -weighting

w of $G(\mathbf{z})$ the subgraph P consisting of all edges with positive weight (and their incident vertices) is the path-subgraph of a Z -path p , every edge of P has positive weight and if e has weight greater than 1 then e is a loop. Hence any path p' with weighting w is a Z -path with the same initial and terminal vertices as p . Given a sequence of Z -cancellations the corresponding path p gives rise to a Z -weighting of $G(\mathbf{z})$.

Lemma 7.18. *Let p_0 and p_1 be Z -paths in $G(\mathbf{z})$ with weightings w_{p_0} and w_{p_1} , respectively, and initial vertex $v = v(\Gamma_s)$, for some basic picture Γ_s . Let Γ_i be the picture obtained from Γ_s by Z -insertion following p_i , $i = 0, 1$. If $w_{p_0} = w_{p_1}$ then the boundary interval b_q has the same label in Γ_0 and Γ_1 , for $1 \leq q \leq W_1$.*

Proof. From Lemma 7.17 the boundary labels of Γ_i depend only on the number of occurrences of edges of P in p_0 and p_1 . As $w_{p_0} = w_{p_1}$ every edge of $G(\mathbf{z})$ occurs the same number of times in p_0 as in p_1 .

Let P be the path-subgraph of a Z -path in $G(\mathbf{z})$, let $e_1, \dots, e_t$ be the edges of P and let $L(s_i)$ be the label of e_i . Let Γ_s be a basic picture for $(\mathbf{z}, \mathcal{L})$ with prime labels $\bar{\mathbf{z}}$, labelled by $\hat{\alpha}_s(\bar{\mathbf{z}})$, for some solution α_s to $\mathcal{H}$. Let $\lambda'_1, \dots, \lambda'_t$ be elements of Λ not occurring in $\mathbf{z}$, $\bar{\mathbf{z}}$ or $\mathcal{L}$. Let (h_q, f_q) be the q th letter of $\mathbf{z}$: so the label of the boundary interval b_q of Γ_s is $h_q \wedge \alpha_s(\lambda_q)$, for $q \in \{1, \dots, W_1\}$. Define the system of parameters $\mathcal{L}(\Gamma_s, P)$ as follows.

$$f_q = \alpha_s(\lambda_q) + l(h_q) \left(\sum_{i=1}^t \lambda'_i \zeta(q, s_i) \right), \text{ for } 1 \leq q \leq W_1,$$

$$\lambda'_i = 1, \text{ if } e_i \text{ is not a loop,}$$

$$\lambda'_i \geq 1, \text{ for } i = 1, \dots, t.$$

Theorem 7.19. *Let P be the path-subgraph of a Z -path with initial vertex $v(\Gamma_s)$ and let α_s be a solution to $\mathcal{H}$ such that (α_s, Γ_s) corresponds to a solution of $Q_{\mathcal{H}}$. Then the following are equivalent.*

- (i) *There exists a solution α to $\mathcal{L}$ and a picture Γ such that (α, Γ) corresponds to a solution of Q and such that Γ_s is obtained from Γ by Z -cancellation, along some Z -path with path-subgraph P and initial vertex $v(\Gamma_s)$.*
- (ii) *There exists a solution α_0 to $\mathcal{L} \cup \mathcal{L}(\Gamma_s, P)$.*

Furthermore α and α_0 may be chosen so that $\alpha_0(\lambda) = \alpha(\lambda)$, for all λ occurring in $\mathbf{z}$ or $\mathcal{L}$.

Proof. Assume (i) holds and let Γ_s be obtained from Γ by Z -cancellation following the Z -path p , with path-subgraph P . Assume p has edges $e_1, \dots, e_t$, let $L(s_i)$ be the label of e_i and let n_i be the number of occurrences of edge e_i in p , $i = 1, \dots, t$. Then, using Theorem 7.17(i), the label of the boundary interval b_q of Γ is

$$(7.2) \quad h_q \wedge \alpha(f_q) = h_q \wedge \left(m_q + l(h_q) \left(\sum_{i=1}^t n_i \zeta(q, s_i) \right) \right),$$

where $m_q = \alpha_s(\lambda_q)$. Define a retraction $\alpha_0 : M \rightarrow \mathbb{Z}$ by $\alpha_0(\lambda) = \alpha(\lambda)$, for $\lambda \in \Lambda \setminus \{\lambda'_1, \dots, \lambda'_t\}$ and $\alpha_0(\lambda'_i) = n_i$, for $i = 1, \dots, t$. Then $\alpha_0(f_q) = \alpha(f_q)$, for

$1 \leq q \leq W_1$, since λ'_i does not occur in $\mathbf{z}$. Hence, from (7.2),

$$\alpha_0(f_q) = m_q + l(h_q) \left(\sum_{i=1}^t \alpha_0(\lambda'_i) \zeta(q, s_i) \right)$$

and, in the light of the above remarks about weightings of Z -paths,

$$\alpha_0(\lambda'_i) \geq 1,$$

for all i , with equality if e_i is not a loop. Hence α_0 is a solution of $\mathcal{L}(\Gamma_s, P)$. As $\alpha_0(\lambda) = \alpha(\lambda)$, when $\lambda \notin \{\lambda'_1, \dots, \lambda'_t\}$, it follows that α_0 is a solution to $\mathcal{L}$ and that the final statement of the Theorem holds.

Conversely, suppose there exists a solution α_0 to $\mathcal{L} \cup \mathcal{L}(\Gamma_s, P)$, where P is the path-subgraph of some Z -path in $G(\mathbf{z})$ with initial vertex $v(\Gamma_s)$. Let P have edges $e_1, \dots, e_t$ and let $L(s_i)$ be the corridor-section labelling e_i . Then assigning weight $n_i = \alpha_0(\lambda'_i)$ to edge e_i of P (and 0 to all other edges of $G(\mathbf{z})$) gives a Z -weighting w of $G(\mathbf{z})$. Let p be a Z -path with path-subgraph P and weighting w : so edge e_i occurs $w(e_i) = \alpha_0(\lambda'_i)$ times in p . Then p has initial vertex Γ_s . Let Γ be the picture obtained from Γ_s by Z -insertion following p and let $m_q = \alpha_s(\lambda_q)$, for $q = 1, \dots, W_1$. Then the label of the boundary interval b_q in Γ is


$$\begin{aligned} h_q \wedge \left(m_q + l(h_q) \left(\sum_{i=1}^t n_i \zeta(q, s_i) \right) \right) \\ = h_q \wedge \alpha_0(f_q), \end{aligned}$$

as α_0 is a solution to $\mathcal{L}(\Gamma_s, P)$. Thus α_0 is a solution to $\mathcal{L}$ and the boundary label b_q of Γ has label $\hat{\alpha}_0(h_q, f_q)$: that is (α_0, Γ) corresponds to a solution of Q .

8. ANGLE ASSIGNMENT

In the following sections we consider figures showing various sub-configurations of a picture Γ . By convention every region shown in these figures is planar. Solid lines incident to vertices represent edges of the graph $\bar{\Gamma}$. Furthermore a dotted line joining two vertices, u and v of a figure, represents either a single edge of $\bar{\Gamma}$ joining u and v or a sequence $a_0, c_1, a_1, \dots, c_n, a_n$, where a_i is an edge of $\bar{\Gamma}$, c_i is a boundary corner of Γ , a_i is incident to c_i and c_{i+1} , for $i = 1, \dots, n-1$, a_0 is incident to u and c_1 , a_n is incident to c_n and v and $n \geq 1$. If a vertex (or region) of a picture Γ appears as vertex (or region) x in some configuration Y we call it a vertex (or region) of type $Y(x)$ or just Y , if the meaning is clear. Not all arcs or edges incident to vertices are necessarily shown in the diagrams. In these configurations the orientation ζ on intervals of $\partial\Sigma$ is shown by an accompanying arrow: for example the orientation of β in configuration $C0^-$ of Figure 42 is from left to right.

Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition of length k , $\mathcal{L}$ a consistent system of parameters, $\mathbf{z}$ a special element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ and α a solution to $\mathcal{L}$. Let Γ be a picture over G , on a compact surface Σ of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with prime labels $\mathbf{z}$, labelled by $\hat{\alpha}(\mathbf{z})$ and with boundary partition $\mathbf{b} = \mathbf{b}_1, \dots, \mathbf{b}_k$. We assign angles to the corners of regions of Γ as follows. Let β be a connected component of $\partial\Sigma$ with boundary partition $\mathbf{b}_u = (b_{l+1}, \dots, b_{l+\eta})$, for some integers l and η , such that $[b_j, b_{j+1}]$ has prime label (h_j, m_j) , with $l(h_j) = d_j$, $j = l+1, \dots, l+\eta$ (where

Figure 28: The angle at c is $-\phi_i - \phi_{i+k-1}$

$l + \eta + 1 \equiv l + 1$). Define

$$\phi_j(\beta) = \frac{d_j(d_j - 1)(\alpha(m_j) - 1)\pi}{\alpha(m_j)^2}, \text{ for } j = l + 1, \dots, l + \eta,$$

noting that under the given hypotheses on Γ the integer $\alpha(m_j) > 0$, for all j . For notational convenience we also define $\phi_i(\beta) = \phi_{l+i}$, $i = 1, \dots, \eta$, given β , l and η as above. Repeat this for all boundary components of Σ so that ϕ_j is defined for $j = 1, \dots, W_1$. If $[b_j, b_{j+1}]$ is a partisan boundary interval then we have $\phi_j = 0$.

We first assign angles to boundary corners of Γ . Let Δ be a region of Γ and let c be a boundary corner of Δ contained in a boundary component β of $\partial\Sigma$. Assume that β has boundary partition $(b_{l+1}, \dots, b_{l+n})$ and that c meets exactly k boundary intervals in this partition. Then there exists a unique integer i such that $c \subseteq [b_i, b_{i+1}] \cup \dots \cup [b_{i+k-1}, b_{i+k}]$ (where subscripts are taken modulo n and lie between $l + 1$ and $l + n$). If $k = 1$ then assign angle $\phi(c) = -\phi_i$ to c . If $k > 1$ then assign angle $\phi(c) = -\phi_i - \phi_{i+k-1}$ to c . (See Figure 28).

We now assign angles to vertex corners. Let Δ be a region of Γ and v a vertex meeting $\partial\Delta$ on a corner c . If both arcs meeting v at c are incident only to vertices we say that c is a corner of *incidence* 0. If one arc meets the boundary $\partial\Sigma$ and the other arc meets only vertices we say c is an *incidence* 1 corner and if both arcs at c meet $\partial\Sigma$ we say c is an *incidence* 2 corner.

We consider two cases.

- (i) First suppose that $\chi(\Delta) = 1$, $\rho(\Delta) = 1$ and $\beta(\Delta) = 1$, (so $t(\Delta) = 2$ and Δ is collapsible), as shown in Figure 29. Assume that Δ meets the boundary component $\beta \subseteq \partial\Sigma$, where β has boundary partition $(b_{l+1}, \dots, b_{l+n})$ and that $\partial\Delta$ meets k boundary intervals in this partition. Then there exists a unique integer i such that $\partial\Delta \cap \partial\Sigma \subseteq [b_i, b_{i+1}] \cup \dots \cup [b_{i+k-1}, b_{i+k}]$ (where subscripts are taken modulo n and lie between $l + 1$ and $l + n$). If $k = 1$ then assign angle $\phi(c) = \phi_i$ to c . If $k > 1$ then assign angle $\phi(c) = \phi_i + \phi_{i+k-1}$ to c . Note that in the latter case $[b_j, b_{j+1}]$ is a partisan boundary interval, for $j = i + 1, \dots, i + k - 2$, and v is an H^Λ -vertex.
- (ii) Now consider all those vertex corners c which do not come into the first case above. If c is an incidence i corner we assign an angle of

$$\phi = \phi(c) = \left(\frac{\rho(\Delta) - 2\chi(\Delta)}{\rho(\Delta)} + \frac{E(\Delta)i}{2} \right) \pi$$




Figure 29:




Figure 30: Configuration AA

to c , where

$$E(\Delta) = \begin{cases} 1, & \text{if } \beta(\Delta) = 0; \\ \beta(\Delta)/(\beta(\Delta) - \varepsilon(\Delta)), & \text{if } \rho(\Delta) \geq \varepsilon(\Delta); \\ (2\beta(\Delta) - \varepsilon(\Delta) + \rho(\Delta))/2(\beta(\Delta) - \varepsilon(\Delta)), & \text{if } \rho(\Delta) < \varepsilon(\Delta). \end{cases}$$

Now we adjust angles on corners of vertices in configuration AA of Figure 30. (The arrows indicate the orientation of boundary components.) We adjust the angle on the corners of x^1 and x^2 in Δ_4^1 and Δ_4^2 , respectively, as well as those on u and v^i . We make the angle on x^i in Δ_4^i equal to $4\pi/6$ and add $\pi/9$ to the corner of v^i in Δ_4^i and $\pi/18$ to both the corner of u in Δ_4^1 and in Δ_4^2 .

Next angles on corners of vertices in configuration AB of Figure 31 are adjusted. We adjust the angle $\phi(c)$ from $8\pi/6$ to $13\pi/12$ and angles $\phi(c_i)$ from $5\pi/6$ to $11\pi/12$, for $i = 1, 2$.




Figure 31: Configuration AB

Note 8.1. (i) If $\rho(\Delta) \geq \varepsilon(\Delta)$ then $E(\Delta) \geq 1$. If $\rho(\Delta) < \varepsilon(\Delta)$ then

$$E(\Delta) = 1 + \left(\frac{\varepsilon(\Delta) + \rho(\Delta)}{2(\beta(\Delta) - \varepsilon(\Delta))} \right) > 1,$$

as $\beta \leq \rho + \varepsilon$ and $E(\Delta)$ is only defined for regions with $\rho(\Delta) \geq 1$.

- (ii) Suppose c is of incidence 1. Then if $\chi(\Delta) = 1$ it follows that $\rho(\Delta) \geq 2$. Hence, if c is of incidence 1, the angle at c is at least $\pi/2$ and is at least $3\pi/2$ if $\chi(\Delta) \leq 0$.
- (iii) Suppose c is of incidence 2 and not in case (i) of the definition of angle above. If $\chi(\Delta) \leq 0$ then $\phi \geq 2\pi$. Assume now that $\chi(\Delta) = 1$, so $\rho(\Delta) \geq 1$. If $\rho(\Delta) = 1$ then $\beta(\Delta) \geq 2$ and $\varepsilon(\Delta) = \beta(\Delta) - 1$ so $\phi \geq \pi$. If $\rho(\Delta) > 1$ then again $\phi \geq \pi$. If $\rho(\Delta) > 2$ then $\phi > \pi$.
- (iv) If Γ has exactly one vertex and Σ is a disk then every vertex corner falls into case (i) of the definition above. We call such a picture a *single vertex disk* (see Figure 33). As minor boundary intervals contribute angles of zero, the sum of angles on the vertex of a single vertex disk may be small: if all the boundary intervals are minor the angle sum will be zero.




Figure 32:




Figure 33: A single vertex disk


Lemma 8.2. *Let c be a vertex-corner of a non-collapsible region Δ and let ϕ be the angle assigned to c . If $\chi(\Delta) \leq 0$ then either*

- i) c is incidence 0 and $\phi \geq \pi$ or*
- ii) c is incidence 1 and $\phi \geq 3\pi/2$ or*
- iii) c is incidence 2 and $\phi \geq 2\pi$.*

Proof. The result is immediate, from the definition of angles and Note 8.1.

Lemma 8.3. *Let c be a vertex-corner of a non-collapsible region Δ and let $\phi = \phi(c)$. Suppose $\chi(\Delta) = 1$ and c is of incidence 0. Then $\rho(\Delta) \geq 3$ and one of the following holds.*

- (i) Δ is of type $AA(\Delta_4^i)$, c is a corner of a vertex of type $AA(u)$ and $\phi = 7\pi/18$ (see Figure 30).*

Figure 34: $\rho(\Delta) = 3$ and $\phi(c) = \pi/3$

(ii)

$$\phi = \left(\frac{\rho(\Delta) - 2}{\rho(\Delta)} \right) \pi$$

and either $\beta(\Delta) = \varepsilon(\Delta) = 0$ or $0 \leq \varepsilon(\Delta) \leq \beta(\Delta) - 1$. In particular if $\rho(\Delta) = 3, 4, 5$ or 6 then $\phi = \pi/3, \pi/2, 3\pi/5$ or $2\pi/3$, respectively and $\phi \geq 5\pi/7$ when $\rho(\Delta) \geq 7$. (See Figures 34(a), 34(b) and 34(c).)

Proof. We have $\rho(\Delta) \geq 3$, for otherwise the label on c is trivial or Δ is collapsible. Since (i) follows directly from the definition of angles we may assume that Δ is not of type $AA(\Delta_4^i)$. Hence $\phi \geq \pi/3$. Clearly $\varepsilon(\Delta) = 0$ if $\beta(\Delta) = 0$. Otherwise as c is incidence 0 and $\chi(\Delta) = 1$ it follows that $\varepsilon(\Delta) \leq \beta(\Delta) - 1$. The remaining statements follow directly from the definition of angles.

Lemma 8.4. *Let a vertex v have a corner c in a non-collapsible region Δ and let ϕ be the angle assigned to c . If $\chi(\Delta) = 1$ and c is incidence 1 then one of the following holds*

- (i) $\phi = \pi/2$, $\rho(\Delta) = 2$, $\beta(\Delta) = 1$ and $\varepsilon(\Delta) = 0$ (see Figure 35(a)).
- (ii) $\phi = 2\pi/3$, Δ is of type $AA(\Delta_4^i)$ and v is of type $AA(x^i)$ (see Figure 30).
- (iii) $\phi = 5\pi/6$, $\rho(\Delta) = 3$, $\beta(\Delta) = 1$ or 2 , $\varepsilon(\Delta) = 0$, Δ is not of type $AB(\Delta)$ and if Δ is of type $AC^+(\Delta_4)$ then v is of type $AC^+(u_1)$ and if Δ is of type $AC^-(\Delta_3)$ then v is of type $AC^-(u_2)$ (see Figures 35(b), 35(c), 31 and 32).
- (iv) $\phi = 11\pi/12$, Δ is of type $AB(\Delta)$ and c is a corner of type $AB(c_i)$, $i = 1$ or 2 (see Figure 31) or
- (v) $\phi = 17\pi/18$, Δ is of type $AA(\Delta_4^i)$ and v is of type $AA(v^i)$, $i = 1$ or 2 (see Figure 30).
- (vi) $\phi \geq \pi$.

Proof.




Figure 35:

As $\chi(\Delta) = 1$ we assume $\gamma(\Delta) = 0$ (or Γ is an empty picture on the disk). As c is incidence 1, $\beta(\Delta) > 0$ and $\rho(\Delta) \geq 2$. Therefore $\phi \geq E(\Delta)\pi/2 \geq \pi/2$. If $\phi = \pi/2$ then $\rho(\Delta)$ must equal 2 and $E(\Delta)$ must equal 1. Since $\beta(\Delta) > \varepsilon(\Delta)$ this occurs only if $\rho(\Delta) \geq \varepsilon(\Delta)$ and $\beta(\Delta) = \varepsilon(\Delta) + 1 = 1$ as required.

If $\phi > \pi/2$ then $\rho(\Delta) > 2$ or $E(\Delta) > 1$. If $\rho(\Delta) = 2$ then $\varepsilon(\Delta) = \beta(\Delta) - 1$. Hence if $2 = \rho(\Delta) \geq \varepsilon(\Delta)$ then $E(\Delta) > 1$ implies that $E(\Delta) = \beta(\Delta) = 2$ or 3. On the other hand if $\varepsilon(\Delta) > \rho(\Delta)$ then $E(\Delta) = (\beta(\Delta) + 3)/2$ and $\beta(\Delta) \geq 4$. In either case $\phi \geq E(\Delta)\pi/2 \geq \pi$.

Now suppose that $\rho(\Delta) > 2$. If Δ is of type $AA(\Delta_4^i)$ then either (ii) or (v) holds. If Δ is of type $AB(\Delta)$ then (iv) holds. If Δ is not of type $AB(\Delta)$ but Δ is of type $AC^+(\Delta_4)$ or type $AC^-(\Delta_3)$ then either (iii) or (vi) holds. Hence we may assume that Δ is not of type $AA(\Delta_4^i)$, $AC^+(\Delta_4)$, $AC^-(\Delta_3)$ or $AB(\Delta)$. Thus $\phi \geq (1/3 + E(\Delta)/2)\pi \geq 5\pi/6$. If $\rho(\Delta) \geq 4$ then $\phi \geq (1/2 + E(\Delta)/2)\pi \geq \pi$. Hence we assume $\rho(\Delta) = 3$. Now c is incidence 1, $\chi(\Delta) = 1$ and $\rho(\Delta) = 3$, so $\varepsilon(\Delta) = \beta(\Delta) - 1$ or $\beta(\Delta) - 2$. If $3 = \rho(\Delta) < \varepsilon(\Delta)$ then


$$E(\Delta) = 1 + \left(\frac{\varepsilon(\Delta) + 3}{2(\beta(\Delta) - \varepsilon(\Delta))} \right) \geq 1 + 7/4 > 2,$$

so $\phi > \pi$. Hence we may assume that $\varepsilon(\Delta) \leq \rho(\Delta)$ and $\rho(\Delta) = 3$.

Now $\phi \geq 5\pi/6$ with equality only if $E(\Delta) = 1$. We have $\beta(\Delta) = \varepsilon(\Delta) + 1$ or $\beta(\Delta) = \varepsilon(\Delta) + 2$ and $E(\Delta) = \beta(\Delta)$ or $E(\Delta) = \beta(\Delta)/2$, respectively. Hence $E(\Delta) = 1$ implies $\beta(\Delta) = 1$ and $\varepsilon(\Delta) = 0$ or $\beta(\Delta) = 2$ and $\varepsilon(\Delta) = 0$. These two possibilities give $\phi = 5\pi/6$ and configurations of Figures 35(b) and 35(c), respectively. Furthermore if $E(\Delta) > 1$ then $E(\Delta) \geq 3/2$ so $\phi \geq 13\pi/12 > \pi$.

Lemma 8.5. *Let a vertex v have a corner c in a non-collapsible region Δ and let ϕ be the angle assigned to c . If $\chi(\Delta) = 1$ and c is incidence 2 then one of the following holds.*

- (i) $\phi = \pi$, $\beta(\Delta) = 2$, $\rho(\Delta) = 1$ or 2 and $\varepsilon(\Delta) = \beta(\Delta) - \rho(\Delta)$ (see Figures 36(a) and 36(b)).
- (ii) $\phi = 13\pi/12$ and c is a corner of type $AB(c)$ (see Figure 31).

Figure 36: $\phi(c) = \pi$

- (iii) $\phi = 7\pi/6$, c is not a corner of type $AB(c)$ and either Δ is of type $AC^+(\Delta_4)$ and v is of type $AC^+(x)$ or Δ is of type $AC^-(\Delta_3)$ and v is of type $AC^-(y)$ (see Figures 32 and 31).
- (iv) $\phi \geq 4\pi/3$.

Proof. If $\rho(\Delta) = 1$ then as $\chi(\Delta) = 1$, $\beta(\Delta) \geq 2$ and $\varepsilon(\Delta) = \beta(\Delta) - 1$. Hence if $\beta(\Delta) = 2$ then $\varepsilon(\Delta) = 1$ and $\phi = \pi$, as shown in Figure 36(a). Otherwise $\beta(\Delta) \geq 3$ and $\rho(\Delta) = 1 < \varepsilon(\Delta)$ so

$$\phi = \left(-1 + \frac{\beta(\Delta) + 2}{2} \right) \pi \geq 3\pi/2.$$

Suppose then that $\rho(\Delta) \geq 2$. If $\rho(\Delta) > 2$ then $\phi \geq 4\pi/3$ unless either case (ii) or (iii) holds. Hence we may assume $\rho(\Delta) = 2$. In this case $\beta(\Delta) = \varepsilon(\Delta) + 2$. If $\rho(\Delta) \geq \varepsilon(\Delta)$ then

$$E(\Delta) = \frac{\beta(\Delta)}{2} \geq 1,$$

with equality only if $\beta(\Delta) = 2$ and $\varepsilon(\Delta) = 0$. If $2 = \rho(\Delta) < \varepsilon(\Delta)$ then

$$E(\Delta) = \frac{\beta(\Delta) + 2 + \rho(\Delta)}{4} = \frac{\beta(\Delta)}{4} + 1 \geq 2.$$

Hence $\phi = E(\Delta)\pi \geq \pi$ with equality only if $\rho(\Delta) = 2$, $\beta(\Delta) = 2$ and $\varepsilon(\Delta) = 0$, as shown in Figure 36(b). If $\phi > \pi$ and $\rho(\Delta) > 2$ then, from the above,

$$\phi \geq \frac{3\pi}{2} > \frac{4\pi}{3}.$$

9. CURVATURE

Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, $\mathcal{L}$ a consistent system of parameters, $\mathbf{z}$ a special element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ and α a solution to $\mathcal{L}$. Throughout this section Γ is assumed to be a reduced, minimalistic picture over G on a compact surface Σ of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with boundary partition $\mathbf{b} = (b_1, \dots, b_{W_1})$, prime labels $\mathbf{z}$ and labelled by $\hat{\alpha}(\mathbf{z})$. Moreover we assume that Γ is not a single vertex disk.

Given a vertex v we denote the sum of angles on all corners of v by $\sigma(v)$. Similarly the sum of angles over all vertex and boundary corners of a region Δ is denoted $\sigma(\Delta)$ and over all corners of a boundary component β of $\partial\Sigma$ by $\sigma(\beta)$.

We define the *curvature* of the vertex v of Γ to be

$$\kappa(v) = 2\pi - \sigma(v) ,$$

of the region Δ to be

$$\kappa(\Delta) = \sigma(\Delta) - t(\Delta)\pi + 2\pi\chi(\Delta)$$

and of the boundary component β of $\partial\Sigma$ to be

$$\kappa(\beta) = -\sigma(\beta) .$$

Lemma 9.1. *Let β be a boundary component of $\partial\Sigma$ with prime label $\prod_{i=1}^n (h_i, m_i)$. Then $\kappa(\beta) \leq \pi \sum_{i=1}^n l(h_i)(l(h_i) - 1)$.*

Proof. Let $l(h_i) = d_i$. We have $l(h_i \wedge \alpha(m_i)) = \alpha(m_i)$, since $\mathbf{z}$ is special. Suppose that β has boundary partition $b_1, \dots, b_n$. A boundary interval $I = [b_i, b_{i+1}]$ meets $[b_i, b_{i+1}]$ boundary corners. If I meets a boundary corner c then I contributes $-\phi_i(\beta)$ to the angle of c . Summing over all boundary intervals of β we see that

$$\sigma(\beta) = \sum_{i=1}^n -\phi_i(\beta) |[b_i, b_{i+1}]|.$$

As $\phi_i = 0$ whenever $[b_i, b_{i+1}]$ is a minor boundary interval we have

$$\sum_{i=1}^n -\phi_i(\beta) |[b_i, b_{i+1}]| = \sum_{i=1}^n -\phi_i(\beta) \alpha(m_i),$$

so

$$\kappa(\beta) = -\sigma(\beta) = \sum_{i=1}^n \alpha(m_i) \phi_i(\beta) \leq \pi \sum_{i=1}^n \frac{d_i(d_i - 1)}{\alpha(m_i)} \leq \pi \sum_{i=1}^n d_i(d_i - 1).$$

Lemma 9.2. *If Δ is a collapsible region of Γ then $\kappa(\Delta) = 0$ when $\rho(\Delta) = 1$ or 2 . When $\rho(\Delta) = 0$ then $\kappa(\Delta) \leq 0$, with equality only if all boundary intervals meeting Δ are minor. If Δ is a non-collapsible region of Γ then*

- (i) if $\beta(\Delta) = 0$ then $\kappa(\Delta) \leq 0$;
- (ii) if $\beta(\Delta) > 0$ and $\rho(\Delta) = 0$ then $\kappa(\Delta) \leq -\pi$;
- (iii) if $\beta(\Delta) > 0$ and $\rho(\Delta) > 0$ then

$$\kappa(\Delta) \leq \begin{cases} 0 & \text{if } \rho(\Delta) \geq \varepsilon(\Delta) \\ -\pi/2 & \text{if } \rho(\Delta) < \varepsilon(\Delta) \end{cases} .$$

Proof. The result follows immediately from the definitions for collapsible regions so we can assume that Δ is non-collapsible. First suppose that Δ is an interior region; that is $\beta(\Delta) = 0$. Then $\rho(\Delta) = t(\Delta)$ so, as Γ is minimalistic,

$$\kappa(\Delta) = \rho(\Delta) \left(\frac{\rho(\Delta) - 2\chi(\Delta)}{\rho(\Delta)} \right) \pi - \rho(\Delta)\pi + 2\pi\chi(\Delta) = 0.$$

Now suppose Δ is a boundary region ($\beta(\Delta) > 0$). Note that the angle adjustments made to corners of vertices of types $AC^\pm(v)$, $AC^+(x)$, $AC^-(y)$, $AB(v)$, $AB(u_i)$,




Figure 37:

$AA(x^i)$, $AA(v^i)$ and $AA(u)$ have no effect on curvature of regions so may be ignored in this context. If $\rho(\Delta) = 0$ then $\sigma(\Delta) \leq 0$ and so $\kappa(\Delta) \leq (-t(\Delta) + 2\chi(\Delta))\pi$. If $\rho(\Delta) = 0$ and $\chi(\Delta) \leq 0$ then $\kappa(\Delta) < -\pi$. If $\rho(\Delta) = 0$ and $\chi(\Delta) = 1$ then, as Γ has no trivial boundary labels, $t(\Delta) \geq 2$. As Δ is non-collapsible we have $t(\Delta) > 2$ and so $\kappa(\Delta) \leq -\pi$. Hence we may assume that $\beta(\Delta) > 0$, $\rho(\Delta) > 0$ and $t(\Delta) = \rho(\Delta) + \beta(\Delta)$. Let $\rho_i = |\{\text{corners of } \Delta \text{ of incidence } i\}|$, for $i = 0, 1, 2$. We have $\rho(\Delta) = \rho_0 + \rho_1 + \rho_2$ and

$$\sigma(\Delta) \leq \rho(\Delta) \left(\frac{\rho(\Delta) - 2\chi(\Delta)}{\rho(\Delta)} \right) \pi + (\rho_1/2 + \rho_2) E(\Delta) \pi,$$

since all angles on boundary corners are non-positive. Now $\rho_1/2 + \rho_2 = \beta(\Delta) - \varepsilon(\Delta)$ so

$$\sigma(\Delta) \leq (\rho(\Delta) - 2\chi(\Delta))\pi + (\beta(\Delta) - \varepsilon(\Delta))E(\Delta)\pi.$$

Thus $\kappa(\Delta) = \sigma(\Delta) - (\rho(\Delta) + \beta(\Delta))\pi + 2\chi(\Delta)\pi$

$$\leq (\beta(\Delta) - \varepsilon(\Delta))\pi E(\Delta) - \beta(\Delta)\pi.$$

If $\rho(\Delta) \geq \varepsilon(\Delta)$ then

$$E(\Delta) = \frac{\beta(\Delta)}{\beta(\Delta) - \varepsilon(\Delta)}$$

and so $\kappa(\Delta) \leq 0$. If $\rho(\Delta) < \varepsilon(\Delta)$ then

$$E(\Delta) = \frac{2\beta(\Delta) - \varepsilon(\Delta) + \rho(\Delta)}{2(\beta(\Delta) - \varepsilon(\Delta))}$$

so that

$$\kappa(\Delta) \leq \frac{(\rho(\Delta) - \varepsilon(\Delta))\pi}{2} \leq -\pi/2.$$

This completes the proof of Lemma 9.2.

Lemma 9.3. *Let v be a vertex incident to two (not necessarily distinct) boundary classes of arc σ_1 and σ_2 (see Figure 37). Then the sum of angles on corners of v between σ_1 and σ_2 is at least π .*

Proof. Suppose that σ_1 and σ_2 are not adjacent classes of arc at v_i : that is, there is at least one class of arcs separating them. The angle on v between a boundary class and any other class is at least $\pi/2$, from Lemmas 8.4 and 8.5. Hence, in this

case, σ_1 and σ_2 are separated by an angle of at least π . On the other hand if σ_1 and σ_2 are adjacent classes then, using Lemmas 8.2 and 8.5, the angle on v between them is less than π only if they meet v at the corner of a collapsible region. In this case σ_1 and σ_2 form 1 class and Γ is a single vertex disk, a contradiction. Hence in all cases the angle between σ_1 and σ_2 is at least π .

Corollary 9.4. *Let v be a boundary vertex. Then $\kappa(v) \leq \pi$. If v meets $\partial\Sigma$ in more than one class of arcs then $\kappa(v) \leq 0$. If v meets $\partial\Sigma$ in more than two classes of arcs then $\kappa(v) \leq -\pi$.*

Let τ be a boundary class of arcs and suppose that one end of τ is contained in a boundary interval $[b, c]$ of some boundary component β of $\partial\Sigma$. In this case we say that τ *meets* $\partial\Sigma$ in $[b, c]$.

Lemma 9.5. *Let v be a boundary vertex of Γ incident to a class of arcs τ which meets $\partial\Sigma$ in a boundary interval $[b, c]$ with prime label (w, f) . If the width of τ is more than $ml/2 + 1$ then w is a cyclic subword of $r^{\pm m}$, (w, f) is a proper exponential H -letter, the width of τ is at most $ml/2 + l - 1$, and $ml/2 - l + 3 \leq l(w) \leq ml/2$.*

Proof. The class τ identifies a subword s of $r^{\pm m}$ which is also a subword of $w \wedge \alpha(f)$. To see this fix a transverse orientation of τ . Starting at the first edge of τ in this orientation we can read round v or along β to the last edge of τ . In the former case we obtain a subword s of $r^{\pm m}$ and in the latter of $w \wedge \alpha(f)$. Note that $l(s) = |\tau| - 1$, where $|\tau|$ denotes the width of τ . Hence $\alpha(f) \geq l(s) \geq ml/2 + 1$. Let $d = l(w)$. If $d \geq ml/2 + 1$ then it follows that w has a cyclic subword s' , which is a cyclic subword of s , such that $l(s') \geq ml/2 + 1$ and s' is a cyclic subword of $r^{\pm m}$. Therefore (w, f) is not relator-reduced, contradicting the hypothesis that $\mathbf{z}$ is special. Hence $l(w) = d \leq ml/2$ and so $\alpha(f) > d$ and in particular (w, f) is a proper exponential H -letter.

Let $\delta = \gcd(l, d)$. First suppose that $d + l - \delta \leq l(s)$. Then since s has periods l and $d = l(w)$, it follows from [16, Section 5, Proposition 1], that s has period δ . This implies that $\delta = l$ so that $l|l(w)$ and w is a cyclic permutation of $r^{\pm t}$ for some $t > 0$, $t \in \mathbb{Z}$. As (w, f) is H^Λ -irredundant it follows that $t = 1$. However this means that (w, f) is relator-unconstrained, a contradiction.

Hence we may assume that $d + l - \delta > l(s)$. As $\delta \geq 1$ and $l(s) \geq ml/2 + 1$ this implies that $d + l - 1 > l(s) \geq ml/2 + 1$, so $d \geq ml/2 - l + 3$. Furthermore $l(s) < ml/2 + l - \delta$ so $|\tau| \leq ml/2 + l - 1$. This completes the proof of Lemma 9.5.

Proposition 9.6. *If v is a vertex of Γ meeting $\partial\Sigma$ in exactly 2 classes of arc then either $\kappa(v) \leq -\pi/12$ or v appears as the vertex a in a configuration of type I1, I2, I3, I4, I5 or I6 as shown in Figures 38(a), 38(b), 38(c), 38(d), 38(e) or 38(f) respectively, or their mirror images.*

Proof. The condition that $\kappa(v) > -\pi/12$ implies $\sigma(v) < 25\pi/12$ which in turn implies that the boundary classes incident at v are separated by angles less than $13\pi/12$, using Lemma 9.3. Let the boundary classes at v be τ_1 and τ_2 and consider the angle ϕ between τ_1 and τ_2 (in the direction of orientation of ∂v). Suppose v is incident to some class of arcs τ between τ_1 and τ_2 . Then it follows, from Lemmas 8.2 to 8.5, as $\phi < 13\pi/12$ that τ is the unique class of arcs between τ_1 and τ_2 and that the regions Δ_i incident to v , τ and τ_i , for $i = 1$ and 2 , are discs with $\rho(\Delta_i) = 2$, $\beta(\Delta_i) = 1$ and $\varepsilon(\Delta_i) = 0$. To see that v belongs to one of the given configurations




Figure 38:

it remains to check the case in which there is no class of arcs incident at v between τ_1 and τ_2 . In this case it follows, from Lemmas 8.2 to 8.5, that as $\phi < 13\pi/12$ the region Δ incident to v , τ_1 and τ_2 is a disc with $\beta(\Delta) = 2$ and $\rho(\Delta) = 1$ or 2 . Hence v must lie in one of the given configurations.

If v is a vertex with $\kappa(v) > -\pi/12$ and v is the vertex a in one of the configurations Ij we say v is an Ij -vertex or v is of type $Ij(a)$.

From now on we make the following assumption.


Assumption 9.7. *The integer m is at least 6 and G is not of type $E(2, *, m)$.*

Theorem 9.8. *Let v be an interior vertex of Γ . Then $\kappa(v) \leq -\pi/3$.*

Proof. Since Γ is minimalistic and G is not of type $E(2, *, m)$, no interior class of arcs can have width more than $l - 2$. It follows that v is incident to at least 7 classes of arcs. Therefore $\kappa(v) \leq -\pi/3$.

We now consider vertices meeting $\partial\Sigma$ in precisely one class of boundary arcs. Suppose v is such a vertex. Then we shall show that if v has curvature $k(v) > -\pi/24$ and v is not an H^Λ -vertex then v belongs to one of a finite number of configurations.

Let v be a vertex incident to exactly one boundary class of arcs. Then v is incident to $\deg(v) - 1$ interior classes of arc, (where $\deg(v)$ denotes the degree of v in the graph $\bar{\Gamma}$ associated to Γ). Each interior class of arcs incident to v is separated from its neighbour by an angle of at least $\pi/3$ so if $\deg(v) > 5$ we have $\kappa(v) < -\pi/3$. Hence we may assume $\deg(v) \leq 5$.


 Figure 39: Configuration $B4$

Lemma 9.9. *Let v be a vertex of Γ incident to at least 4 interior and precisely 1 boundary class of arcs. Then either $\kappa(v) \leq -\pi/9$ or v is of type $B4(a)$ as shown in Figure 39.*

Proof. If $\deg(v) \geq 6$ then $\kappa(v) \leq -\pi/3$ so we may assume $\deg(v) \leq 5$. Then, from Lemmas 8.3 and 8.4, $\kappa(v) \leq 0$ with equality only if v is of type $B4(a)$, as shown in Figure 39, and not of type $AA(u)$. If v is of type $AA(u)$ then $\kappa(v) = -\pi/9$. If v is not of the type $B4(a)$ then $\kappa(v) < 0$ and, from Lemmas 8.3 and 8.4 again, either one of the incidence 0 corners at v has angle $\phi \geq \pi/2$ or one of the incidence 1 corners has angle $\phi \geq 5\pi/6$. In either case $\kappa(v) \leq -\pi/6$.

Corollary 9.10. *Let v be a vertex of Γ meeting $\partial\Sigma$ in precisely one class of arc of width at most $ml/2 + 1$. Then either $\kappa(v) \leq -\pi/9$ or v is of type $B4(a)$ as shown in Figure 39.*

Proof. There are at least $ml/2 - 1$ non-boundary arcs incident at v . Since Γ is minimalistic, $ml/2 - 1 \geq 3l - 1$ and G is not of type $E(2, *, m)$ there are at least 4 non-boundary classes of arc incident at v . The result follows from Lemma 9.9.

Lemma 9.11. *Let u and v be adjacent vertices of Γ with incidence 1 corners on a region Δ such that $\rho(\Delta) = 2$, $\beta(\Delta) = 1$, $\varepsilon(\Delta) = 0$ and $\chi(\Delta) = 1$, as shown in Figure 40. Let μ and ν be the boundary classes of arcs incident to u and v , respectively, and meeting Δ and assume that $\partial\Sigma \cap \mu$, $\partial\Sigma \cap \nu$ and the boundary corner of Δ are all contained in a boundary interval $\beta = [b, c]$ with prime label (w, f) . Assume that $\kappa(v) > -\pi/2$ and that $|\nu| \geq ml/2 + 2$. Then $|\mu| \leq l$ and w is a cyclic subword of $r^{\pm m}$.*




Figure 40:

Proof. The class μ meets u in $|\mu|$ arcs. Reading round u in the direction determined by the transverse orientation of μ induced from the chosen orientation ζ of $\partial\Sigma$ these arcs identify a subword s of $r^{\varepsilon m}$, where $\varepsilon = \pm 1$. Similarly the class ν meets v and identifies a subword t of $r^{\varepsilon m}$, where $\varepsilon = \pm 1$. For the duration of this proof we call μ (ν) *positive* if $\varepsilon = 1$ and *negative* otherwise. We have $l(s) = |\mu| - 1$ and $l(t) = |\nu| - 1$. Assume that $|\nu| \geq ml/2 + 2$ and that $|\mu| \geq l + 1$.

We may assume, without loss of generality, that the orientation ζ of β is such that $\Delta \cap \beta$ is oriented from μ to v and that an occurrence of w in the label of $[b, c]$ begins in the boundary corner following the first arc of ν (in this orientation of β). Then as above μ identifies a cyclic subword x of $w \wedge \alpha(f)$ and ν identifies an initial subword y of $w \wedge \alpha(f)$. Furthermore in H we have $x = s$ and $y = t$. From Lemma 9.5 it follows that w is a cyclic subword of $r^{\pm m}$ and $2l + 3 \leq ml/2 - l + 3 \leq l(w) \leq ml/2$, so w is an initial subword of y . Let x' be the terminal segment of x of length l and let y' be the segment of y beginning in position $l(w) - l$ and of length l . Then both x' and y' are subsegments of w beginning in position $l(w) - l$ and ending in position $l(w) - 1$. Hence $x' = y'$, in H . As $x = s$ and $y = t$ both x' and y' are cyclic permutations of $r^{\pm 1}$.

We consider separately the cases in which μ is positive and negative, under the assumption that ν is positive. The case ν negative follows by symmetry. First suppose μ is positive so x' and y' are both cyclic permutations of r . As cyclic subwords of r we may assume the position of the first letters of x' and y' are $\lambda \leq l/2$ apart. (Otherwise interchange x' and y' .) The union of x' and y' then forms a cyclic subword z of r^m of length $2l - \lambda$, and of periods l and λ . Let $\delta = \gcd(l, \lambda)$. Then $l + \lambda - \delta \leq 3l/2 - 1 < 2l - \lambda = l(z)$. Hence from [16, Section 3, Proposition 1] the word z has period δ . As r is not a proper power it follows that $\delta = l$ and since $\delta | \lambda$ and $\lambda \leq l/2$ we have $\lambda = 0$. Thus x' and y' are identically positioned as cyclic subwords of r .

Now suppose that the last letter of w , as a word in $A * B$, is ω . Then the letter following y' in y , and therefore in r^m , is ω . Hence the letter following x' in r^m is ω and so the label on the corner of u in Δ is ω . Therefore the label on the corner of Δ meeting u , read with orientation induced by that on β , is ω^{-1} . As w ends in the letter ω the boundary component of Δ is labelled ω . Hence the corner of v in Δ has trivial label, a contradiction. Hence μ cannot be positive.

If μ is negative then x' is a cyclic permutation of r^{-1} and y' is a cyclic permutation of r . As $x' = y'$ in $A * B$ this means that G is of type $E(2, 2, m)$, contrary to the standing assumption 9.7. Therefore $|\mu| \leq l$.

10. CONFIGURATIONS C

Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition, $\mathcal{L}$ a consistent system of parameters, $\mathbf{z}$ a special element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ and α a solution to $\mathcal{L}$. As in Section 9, Γ is assumed, throughout this section, to be a reduced, minimalistic picture over G on a compact surface Σ of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with boundary partition $\mathbf{b} = (b_1, \dots, b_{W_1})$, prime labels $\mathbf{z}$ and labelled by $\hat{\alpha}(\mathbf{z})$. We are now in a position to prove the following.

Theorem 10.1. *Let v be a vertex of Γ which meets $\partial\Sigma$ in precisely one boundary class of arcs ν . Then either*

- (i) *v is distance at most 1 from an H^Λ -vertex or*
- (ii) *$\kappa(v) \leq -\pi/24$ or*
- (iii) *v is of type $B4(a)$ (see Figure 39) or*
- (iv) *v appears as the vertex v in configuration $C0$, $C1^\pm$, $C2^\pm$, $C3^\pm$, $C4$ or $CC^\pm$, $|\nu| > ml/2 + 1$, and in each of these configurations $|\mu_i| \leq l$ (see Figures 42, 43, 44, 45, 46 and 47).*

Proof. Assume that v satisfies none of the conditions (i)–(iii). As v is not an H^Λ -vertex it follows that ν meets $\partial\Sigma$ in a boundary interval $[b, c]$ which we shall assume has prime label (w, f) . Note that failure of condition (ii) implies that v is not of type $AC^\pm(v)$, $AB(u_i)$, $AA(u)$ or $AA(v^i)$. From Lemma 9.9 it follows that v is incident to at most 3 interior classes of arcs. From Corollary 9.10 it follows that $|\nu| > ml/2 + 1$ and from Lemma 9.5 it follows that (w, f) is a proper exponential H -letter, w is a subword of $r^{\pm m}$, $ml/2 - l + 3 \leq l(w) \leq ml/2$ and $|\nu| \leq ml/2 + l - 1$. As $m \geq 6$ this implies that there are at least $2l + 1$ non-boundary arcs incident at v . Hence v is incident to exactly 3 interior classes of arcs and 1 boundary class of arcs ν .

We denote the regions incident to v by Δ_1 , Δ_2 , Δ_3 and Δ_4 , as in Figure 41. (Note that these Δ 's are not *a priori* distinct.) Let α_i denote the angle on the corner of v in Δ_i for all i .

We have $\alpha_i \geq \pi/3$, for $i = 1, 2$, and $\alpha_i \geq \pi/2$ for $i = 3, 4$ (using Lemmas 8.2, 8.3 and 8.4 and the fact that v is incident to exactly one boundary class of arcs). If $\alpha_i \geq 5\pi/7$ for $i = 1$ or 2 then $\kappa(v) \leq -\pi/21$. If $\alpha_i \geq 17\pi/18$ for $i = 3$ or 4 then $\kappa(v) \leq -\pi/9$. If $\alpha_i = 11\pi/12$, for $i = 3$ or 4 , then v is of type $AB(u_i)$, a possibility that has already been excluded.

Hence we may assume $\alpha_i \leq 2\pi/3$, for $i = 1, 2$ and $\alpha_i \leq 5\pi/6$ for $i = 3, 4$. (Using Lemmas 8.2, 8.3 and 8.4 again). In particular $\chi(\Delta_i) = 1$, for $i = 1, 2, 3, 4$, which implies that all the Δ 's are distinct. Also $3 \leq \rho(\Delta_i) \leq 6$ for $i = 1, 2$ and $2 \leq \rho(\Delta_i) \leq 3$, for $i = 3, 4$. For $i = 3$ or 4 , write

$$T(\Delta_i) = \begin{cases} a & \text{if } \rho(\Delta_i) = 3 \text{ and } \beta(\Delta_i) = 1 \\ b & \text{if } \rho(\Delta_i) = 3 \text{ and } \beta(\Delta_i) = 2 \\ c & \text{if } \rho(\Delta_i) = 2 \end{cases}.$$

If $(T(\Delta_3), T(\Delta_4)) = (a, a)$, (a, b) , (b, a) or (b, b) then $\kappa(v) \leq -\pi/3$ so we must have $(T(\Delta_3), T(\Delta_4)) = (a, c)$, (b, c) , (c, c) , (c, a) or (c, b) . We consider the possible values of $\rho(\Delta_i)$, with $i = 1, 2$ in each of these cases.




Figure 41:

Consider first the case $(T(\Delta_3), T(\Delta_4)) = (c, a)$. If $\rho(\Delta_i) \geq 4$, for $i = 1$ or 2 , then $\kappa(v) \leq -\pi/6$. Hence $\rho(\Delta_i) = 3$ in this case, for $i = 1$ and $i = 2$. This gives configuration CC^- . The case $(T(\Delta_3), T(\Delta_4)) = (a, c)$ gives CC^+ . A similar argument shows that in the cases $(T(\Delta_3), T(\Delta_4)) = (c, b)$ and (b, c) the vertex v is of type $AC^\pm(v)$ or $AB(u_i)$ so these cases are already covered in (ii).

Now consider the case $(T(\Delta_3), T(\Delta_4)) = (c, c)$. We have $3 \leq \rho(\Delta_i) \leq 6$, for $i = 1, 2$ and it is easy to check that $\kappa(v) > -\pi/24$ implies $(\rho(\Delta_1), \rho(\Delta_2)) = (3, 3), (3, 4), (3, 5), (3, 6), (4, 4), (6, 3), (5, 3)$ or $(4, 3)$.

These values of $\rho(\Delta_i)$ give rise to configurations $C0$, $C1^\pm$, $C2^\pm$, $C3^\pm$ and $C4$. As v is distance more than 1 from all H^Λ -vertices it follows, from Lemma 9.11 that the boundary classes μ_i in these configurations all have width at most l .

Definition 10.2. . A configuration of type $C0$, $C1^\pm$, $C2^\pm$, $C3^\pm$, $C4$ or $CC^\pm$ is called a *C-configuration* (see Figures 42, 43, 44, 45, 46 and 47). A vertex u of Γ , with $\kappa(u) > -\pi/24$, which is distance at least 2 from all H^Λ -vertices and which appears as vertex v in a *C-configuration* is called a *C-vertex*.

We shall for convenience refer to configurations $Cj^\pm$, $j = 0, \dots, 4$, taking $Cj^+ = Cj^- = Cj$ when $j = 0$ or 4 .

11. CONFIGURATIONS D

Proposition 11.1. *If v is a C-vertex then either*

- (i) *we can adjust the angle assignment to corners of u_i and v so that $\kappa(v), \kappa(u_i) \leq -\pi/30$, for $i = 1, 2$, and $\kappa(\Delta)$ is unchanged for all regions Δ ; or*
- (ii) *v is the vertex a and u_1 is the vertex b in one of configurations $D0$, $D1$, $D2$, $D3$, $D4$, $D5$, $D6$, $D7$, $D8$ or $D9$.*

Such angle adjustments may be carried out over all C-configurations simultaneously.




 Figure 42: Configuration $C0$


Figure 43:

Proof. We shall define angle adjustments to satisfy (i) and (ii) for all C -configurations of Γ such that, considering each connected boundary component β of $\partial\Sigma$ in turn, these angle adjustments move angle from corner to corner of regions incident to




Figure 44:




Figure 45:






 Figure 46: Configuration $C4$


Figure 47:

β , only in the direction of orientation of β . Furthermore these angle adjustments leave $\kappa(\Delta)$ fixed, for all regions Δ of Γ .

Given a C -configuration with C -vertex v set $u = u_1$ and $\mu = \mu_1$ (if we're not in case CC^+). Denote the class of arcs joining u and v by ι . Denote the angle on the

Figure 48: Configuration $D0$ Figure 49: Configuration $D1$



corners of v and u in Δ_4 by α_4 and α_5 , respectively. Orient ∂u in the direction from μ to ι inside Δ_4 . Note that v is not of type $AA(v^i)$ these vertices have curvature $\kappa(v^i) \leq -\pi/9$. By definition v is not of type $AA(x^i)$, $AC^\pm(v)$, $AC^+(x)$, $AC^-(y)$, $AB(u_i)$ or $AB(v)$. Therefore no corners of v have had angles altered by previous angle adjustments.

To ensure that we can make angle adjustments simultaneously over all C -configurations we need to consider whether or not u and v may appear in more than one C -configuration. As v is incident to only one boundary class of arcs it cannot occur as the C -vertex of any other C -configuration. As $|\mu| \leq l$ and $|\nu| \geq ml/2 + 1$, u cannot be the C -vertex of a C -configuration in which v occurs as the vertex of type u_1 or u_2 . Suppose that u occurs as the C -vertex in some



Figure 50: Configuration $D2$ Figure 51: Configuration $D3$

C-configuration. Let C_0 be the original *C*-configuration, in which u is of type u_1 , and let C_1 be the configuration in which u occurs as the *C*-vertex. Since u is incident to a boundary class in C_1 it follows that C_0 is of type CC^+ , so u being the vertex of type $CC^+(u_1)$ in C_0 is not incident to a further boundary class of arcs. If C_1 is a configuration of type Cj then, since $\rho(\Delta_1) = 3$, it follows that C_1 is of type $C0$, with u of type $C0(v)$. This gives configuration $D'6$ of Figure 61. Otherwise C_1 is of type $CC^\pm$ and we obtain configuration $D8$ of Figure 56 or $D9$ of Figure 57, with u of type $Dj(b)$. In all three cases it can be easily verified that u is of type u_1 in C_0 alone.



In general u may occur as the vertex of type u_1 in more than one C -configuration. Suppose then that u is the vertex of type u_1 in p distinct C -configurations. We

Figure 52: Configuration $D4$ Figure 53: Configuration $D5$

show that (ii) holds only if $p = 1$ and that if $p = 1$, (ii) does not hold and u and v are not of type $D'6(u)$ and $D'6(v)$, respectively, then we can decrease the angle α_5 and increase α_4 to obtain $\kappa(u), \kappa(v) \leq -\pi/12$. If $p > 1$ we show that we can decrease the angle α_5 and increase α_4 to obtain $\kappa(u), \kappa(v) \leq -\pi/15$ over all p configurations, in which u occurs as the vertex of type u_1 , simultaneously.


 Figure 54: Configuration $D6$

 Figure 55: Configuration $D7$

Finally suppose that u and v are of type $D'6(u)$ and $D'6(v)$, respectively, and let β' be the boundary component to which u is incident in Figure 61. If β' is oriented from left to right in Figure 61 we show that we can adjust angles by increasing α_4 and α_7 and decreasing α_5 and α_6 so that $\kappa(u), \kappa(v), \kappa(x) \leq -\pi/30$. Similarly if β' is oriented from right to left in Figure 61 we show that we can adjust angles by

Figure 56: Configuration $D8$ Figure 57: Configuration $D9$

increasing α_4 and α_8 and decreasing α_5 and α_9 so that $\kappa(u), \kappa(v), \kappa(y) \leq -\pi/30$. These adjustments all leave curvature of regions unaltered. This is sufficient, given the remark at the beginning of the proof, to prove the theorem. We divide the proof into Case A, where $p = 1$, and Case B, where $p > 1$.




Figure 58:

Case A: $p = 1$.

In this case u occurs as the vertex of type u_1 in no C -configuration except that in which v occurs. Under this assumption we have two further cases to consider: Case A1 in which u is adjacent to no boundary classes (in configuration CC^+) except possibly μ (in configurations $Cj^\pm, j = 0, \dots, 4$ or CC^-) and Case A2 in which u is incident to boundary classes other than μ .

Case A1. As G is not of type $E(2, *, m)$ no interior class of arcs contains more than $l - 2$ arcs and μ contains no more than l arcs, from Theorem 10.1. It follows that in Case A1, configuration CC^+ , u is adjacent to at least 7 interior classes of arc and in Case A1, configuration $Cj^\pm$ or CC^- , u is adjacent to at least 6 interior classes of arc.

In configuration $Cj^\pm$, with $0 \leq j \leq 4$, or CC^- with u incident to 6 or more interior classes of arc we have $\sigma(u) \geq \pi + 5\pi/3$, so we can adjust α_5 , by $-\pi/2$, to 0 and α_4 , by $\pi/2$, to π giving $\kappa(u), \kappa(v) \leq -\pi/6$. In configuration CC^+ with u is incident to 7 or more classes of arc we have $\sigma(u) \geq 7\pi/3$, $\alpha_5 = \pi/3$ and $\alpha_4 = 5\pi/6$ so we may adjust to $\alpha_5 = \pi/6$ and $\alpha_4 = \pi$, to give $\kappa(u), \kappa(v) \leq -\pi/12$.

Hence in Case A1 it is always possible to adjust angles as required.

Case A2. Now suppose that u is incident to some boundary class of arcs ξ not meeting $\partial\Delta_4$. Suppose first we have one of configurations $C1^\pm, C2^\pm, C3^\pm, C4$ or CC^- (see Figure 58). In each case $\kappa(v) \leq \pi/6$. Let θ be the sum of angles on u between ι and ξ (that is, taken in the direction of Δ_1). If (in the same direction) there is 1 or more class of arcs incident at u between ι and ξ then it follows from Lemmas 8.2, 8.3 and 8.4 that $\theta \geq 5\pi/6$. If there are no arcs between ι and ξ then since $\rho(\Delta_1) \geq 3$ and, as is easily verified, Δ_1 is not a region of type $AA(\Delta_4^i)$, it




Figure 59:

Hence we need only consider configurations $C0$ and CC^+ .

Assume we have configuration CC^+ , so $\kappa(v) = 0$ (see Figure 59). Suppose first Δ_1 is an interior region, so triangular with all angles $\pi/3$. In this case if any region meeting u has Euler characteristic less than 1 then $\sigma(u) \geq 2\pi/3 + \pi + \pi/2 = 13\pi/6$, so we may decrease α_5 to $\pi/4$ and increase α_4 to $11\pi/12$ to obtain $\kappa(u), \kappa(v) \leq -\pi/12$. Hence we may assume all regions Δ incident to u have $\chi(\Delta) = 1$. Now under the current assumptions u is incident to 3 or more interior classes of arc and one or more boundary classes of arc. We consider four subcases. Subcase A2.1 where u is incident to more than one boundary class; Subcase A2.2 where u is incident to one boundary class and more than four interior classes; Subcase A2.3 where u is incident to 1 boundary class and four interior classes and Subcase A2.4 where u is incident to 1 boundary class and three interior classes.

Subcase A2.1: u is incident to more than one boundary class of arcs.

In this case $\sigma(u) \geq 2\pi/3 + \pi + \pi/2 + \pi/2$ so we may adjust angles α_4 and α_5 to give $\kappa(u), \kappa(v) \leq -\pi/3$.

Subcase A2.2: u is incident to one boundary class and more than four interior classes of arcs.

In this case $\sigma(u) \geq 4\pi/3 + \pi/2 + \pi/2$ so we may adjust angles α_4 and α_5 to obtain $\kappa(u), \kappa(v) \leq -\pi/6$.

Subcase A2.3: u is incident to one boundary class and four interior classes of arcs.

In this case either $\kappa(u) \leq -\pi/6$ or u is a vertex of type $B4(a)$ (see Figure 39) or u is of type $AA(u)$ (see Figure 30). If $\kappa(u) \leq -\pi/6$ then we can adjust α_4 and α_5 by $\pi/12$ to obtain $\kappa(u), \kappa(v) \leq -\pi/12$. If u is of type $AA(u)$ then v is of type $AA(v^i)$ or $AA(x_i)$, so this does not occur. Therefore we may assume u is a vertex of type $B4(a)$. In this case every incidence 0 corner on u must have angle $\pi/3$ and both incidence 1 corners must have angle $\pi/2$. There are therefore two possibilities as shown in Figures 52 and 53, configurations $D4$ and $D5$ respectively.




Figure 60: Configuration AC4




Figure 61: Configuration $D'6$

Subcase A2.4: u is incident to one boundary class and 3 interior classes of arcs.

In this case we have the situation of Figure 60. The angles θ and ϕ in Figure 60 are both either equal to $\pi/2$ or at least $5\pi/6$. If $\phi = \theta = \pi/2$ then u is of type $C0(v)$ and we have configuration $D'6$ of Figure 61. We




Figure 62:

defer further consideration of $D'6$ until the end of the proof. If $\phi = 5\pi/6$ and $\theta = \pi/2$ we have configuration $D7$, of Figure 55, with u of type $CC^\pm(v)$. If $\phi = \pi/2$ and $\theta = 5\pi/6$ then we have configuration $D8$, of Figure 56, with u of type $CC(v)^\pm$. If $\phi \geq 5\pi/6$ and $\theta \geq 5\pi/6$ then $\sigma(u) \geq 2\pi/3 + 5\pi/3 = 7\pi/3$ so we may adjust α_4 and α_5 by $\pi/6$ to give $\kappa(u), \kappa(v) \leq -\pi/6$.

This completes consideration of Case A2, configuration CC^+ with Δ_1 an interior region (except for configuration $D'6$). Now suppose that Δ_1 is a boundary region. As v is of type CC^+ we have $\rho(\Delta_1) = 3$ and $\beta(\Delta_1) = 1$. Hence the angle on the corner of u in Δ_1 is $5\pi/6$. If u is incident to a further boundary class of arcs (other than that in $\partial\Delta_1$) then $\sigma(u) \geq 5\pi/6 + \pi/3 + \pi/2 + \pi = 8\pi/3$ so we may adjust α_4 and α_5 by $\pi/3$ to give $\kappa(u), \kappa(v) \leq -\pi/3$. Hence we may assume u is incident to no further boundary classes of arcs. We have the situation of Figure 62, with ξ be the boundary class incident at u . Now, from Lemma 9.5 and the fact that v is not joined to an H^Λ -vertex, it follows that $|\xi| \leq ml/2 + l - 1$ and so u is adjacent to at least one more vertex. Hence $\theta \geq 5\pi/6$. If $\theta > 5\pi/6$ then $\theta \geq \pi$, using Lemmas 8.3 and 8.4, so $\sigma(u) \geq 13\pi/6$ and we may adjust α_4 and α_5 by $\pi/12$ to obtain $\kappa(u), \kappa(v) \leq -\pi/12$. Hence we may assume $\theta = 5\pi/6$ in which case we have configuration $D9$, with u of type $CC^\pm(v)$.

It remains to consider configurations $C0$ to complete Case A2 (except for $D'6$ which is treated later). In Case A2, configuration $C0$, we have the situation of Figure 63 with $\sigma(u) = \theta + \phi + \pi/2$ and $\kappa(v) \leq -\pi/6$. If u is incident to any region Δ with $\chi(\Delta) \leq 0$ then, since $\chi(\Delta_1) = 1$, Lemma 8.2 implies that either $\theta \geq 11\pi/6$ or $\phi \geq 2\pi$ so $\sigma(u) \geq 20\pi/6$. Hence we may adjust α_4 and α_5 by $\pi/2$ to obtain $\kappa(u), \kappa(v) \leq -\pi/6$. Hence we may assume $\chi(\Delta) = 1$, for all regions Δ meeting u .

We consider first the case that Δ_1 is an interior region, that is Δ_1 is triangular. In this case $\theta = \pi/3 + \psi$, for some angle ψ with $\psi \geq \pi/2$. As $\phi \geq \pi$ we have $\phi + \psi \geq 3\pi/2$. If $\phi + \psi = 3\pi/2$ then $\psi = \pi/2$ and $\phi = \pi$ and it is easy to check that we have one of configurations $D0$, $D1$ or $D2$. If $\psi > \pi/2$ then $\psi \geq 5\pi/6$, because






Figure 63:


 Figure 64: Configuration $D3'$

if $\psi = 2\pi/3$ then u is of type $AA(x^i)$ and v is forced to be of type $AA(v^i)$. Also $\phi \geq 7\pi/6$ (in fact $\phi \geq 4\pi/3$ unless u is of type $AC^+(x)$, $AC^-(y)$ or $AA(x^i)$). In both cases we may adjust α_4 and α_5 by $5\pi/12$ to obtain $\kappa(u), \kappa(v) \leq -\pi/12$.

Now suppose that Δ_1 is a boundary region. Then, in Figure 63, $\theta = 5\pi/6$ and $\phi \geq \pi$. If $\phi = \pi$ then we have configurations $D3$, $D3'$ or $D3''$ of Figures 51, 64 and 65, respectively. Theorem 9.11 implies that, in configurations $D3'$ and $D3''$, the boundary class incident to u in $\partial\Delta_4$ has size at most l so the boundary class incident to u in $\partial\Delta_1$ has size at least $(m-2)l > (ml/2) + l - 1$. Since u is not an H^Λ -vertex this contradicts Lemma 9.5. Hence the configurations $D3'$ and $D3''$ cannot occur. If $\phi \geq \pi$ then $\phi \geq 7\pi/6$ and we can adjust α_4 and α_5 by $5\pi/12$ to

Figure 65: Configuration $D3''$

give $\kappa(u), \kappa(v) \leq -\pi/12$. This completes Case A2 and so proves Case A: $p = 1$, of the theorem, with the exception of configuration $D'6$.

Case B: $p > 1$.

Suppose u is the vertex of type u_1 adjacent to C -vertices $v^1, \dots, v^p$, occurring in cyclic order around u , with $v = v^1$ and $p > 1$. Note that it is possible that u is a vertex of type $Cj^\pm(u_2)$ or $CC^\pm(u_2)$ but then the corresponding vertex $Cj^\pm(v)$ or $CC^\pm(v)$ is not among $v^1, \dots, v^p$. If v^k is the vertex v in one of configurations $C0$, $C1^\pm$, $C2^\pm$, $C3^\pm$, $C4$ or $CC^\pm$ we denote the corresponding vertices u_i , classes μ_i , ι_i and regions Δ_i by u_i^k , μ_i^k , ι_i^k and Δ_i^k , for $i = 1, 2$ and $k = 1, \dots, p$. Hence $u = u_1^k$, for $k = 1, \dots, p$. Write μ^k and ι^k for μ_1^k and ι_1^k , respectively and recall that ∂u is oriented from μ to ι inside Δ_4 . If μ^k precedes ι^k in the cyclic order around u then we say that v^k is *green* and otherwise that v^k is *blue*. (See Figure 66.)

We define

$$T(v^k) = \begin{cases} X, & \text{if } v^k \text{ is of type } CC(v)^+; \\ Y, & \text{if } v^k \text{ is of type } CC(v)^-; \\ Z, & \text{if } v^k \text{ is of type } Cj(v), \text{ with } 0 \leq j \leq 4 : \end{cases}$$

and say that v^k is of *type* X, Y or Z as appropriate.

We now compute a lower bound for the sum of angles θ^k on u between i^k and i^{k+1} , read in the direction of orientation of u (superscripts modulo p). We consider separately each of the four possible colourings, in green and blue, of v^k and v^{k+1} .

Case B1: v^k and v^{k+1} are green.

We treat the different combinations of types of v^k and v^{k+1} as distinct subcases.

Subcase B1.1: $T(v^k) = X, Y$ or Z , $T(v^{k+1}) = X$, see Figure 67. In the following Figures vertices with broken outlines may or may not be present. If they're absent their incident arcs are replaced by a single arc. In Figure 67 either one or none of the dotted vertices may be present.




Figure 66:

In this subcase as v^k is incident to only one boundary class the angle ϕ in Figure 67 is at least $\pi/3$, and so $\theta^k \geq 2\pi/3$.

Subcase B1.2: $T(v^k) = X, Y$ or Z , $T(v^{k+1}) = Y$ or Z , (see Figure 68, where at most one dotted vertex may be present).

In this case, as v^k is incident to only one boundary class, $\phi \geq 5\pi/6$ so that $\theta^k \geq 4\pi/3$.

Case B2: v^k and v^{k+1} are both blue.

Subcase B2.1: $T(v^k) = X$, (see Figure 69). In this case $\phi \geq \pi/3$ so $\theta^k \geq 2\pi/3$.

Subcase B2.2: $T(v^k) = Y$ or Z , (see Figure 70). As v^{k+1} is incident to only one boundary class we have $\phi \geq 5\pi/6$. Hence $\theta^k \geq 4\pi/3$.

Case B3: v^k is blue, v^{k+1} is green.

Subcase B3.1: $T(v^k) = X$, $T(v^{k+1}) = X$, (see Figure 71). Note that the vertex x in Figure 71 cannot be v^{k+1} , given that we have defined $u = u_1^{k+1}$




Figure 67:

with respect to the orientation of the boundary component β^{k+1} of $\partial\Sigma$ incident to v^{k+1} . We have $\phi \geq \pi/3$, so $\theta^k \geq 2\pi/3$.

Subcase B3.2: $T(v^k) = X$, $T(v^{k+1}) = Y$ or Z , (see Figure 72). As $\phi \geq \pi/2$ we have $\theta^k \geq 4\pi/3$.

Subcase B3.3: $T(v^k) = Y$ or Z , $T(v^{k+1}) = X$, (see Figure 73) Here $\phi \geq \pi/2$ so $\theta^k \geq 4\pi/3$.

Subcase B3.4: $T(v^k) = Y$ or Z , $T(v^{k+1}) = Y$ or Z , (see Figure 74) We have $\phi \geq \pi$ so $\theta^k \geq 2\pi$.

Case B4: v^k is green, v^{k+1} is blue, (see Figure 75). In this case $\theta^k \geq \pi/3$. We have therefore a lower bound for θ^k in all cases.

If v^k is green for all k with $c \leq k \leq d$ and v^{c-1} and v^{d+1} are blue (or $d+1-c \equiv 0 \pmod{p}$) then $v^c, v^{c+1}, \dots, v^d$ is called a *green component* (of u). *Blue components* are defined analogously.

Case B5: $v^1, \dots, v^p$ are all green. Suppose that $v^1, \dots, v^p$ are all green. Let $t = |\{k : T(v^k) = Y \text{ or } Z\}|$. Then $\sigma(u) \geq (2\pi/3)p + (2\pi/3)t$. If $p \geq 4$ then $\sigma(u) \geq 2\pi + (p-3)(2\pi/3) + (2\pi/3)t$ so

$$\sigma(u) \geq 2\pi + (p+1)2\pi/15 + (2\pi/3)t.$$




Figure 68:

Noting that $\kappa(v^k) \leq 0$ if $T(v^k) = X$ or Y and that $\kappa(v^k) \leq \pi/3$ if $T(v^k) = Z$ we see that we can adjust angles on u and v^k inside Δ_4^k so that $\kappa(u) \leq -2\pi/15$ and $\kappa(v^k) \leq -2\pi/15$, for all k . (This follows because $t \geq |\{k : T(v^k) = Z\}|$.)

Thus given that all v^k 's are green it remains to consider the cases $2 \leq p \leq 3$.

Subcase B5.1: $p = 3$.: In this case if $t \geq 1$ then $\sigma(u) \geq 2\pi + (2\pi/3)t \geq 2\pi + (\pi/3) + (\pi/3)t$. It follows that we can adjust angles α_4^k and α_5^k for $k = 1, 2, 3$ so that $\kappa(u)$ and $\kappa(v^k) \leq -\pi/12$. We may therefore suppose that $t = 0$. In this case we have the configuration of figure 76, and $\sigma(u) = 2\pi$ if and only if $\phi^1 = \phi^2 = \phi^3 = \pi/3$ (in figure 76). As v^i is incident to only one boundary class this occurs only if v^i is incident to x^{i+1} , for $i = 1, 2, 3$ (superscripts modulo 3). However the resulting configuration cannot occur unless G is of type $E(2, *, m)$, using Lemma 6.12. Hence $\phi^i > \pi/3$, for some i . As $\rho(\Delta_1^i) = 3$ it follows that $\phi^i \geq 2\pi/3$, for such i , so $\sigma(u) \geq 7\pi/3$ and we may adjust angles so that $\kappa(u), \kappa(v^i) \leq -\pi/12$.

Subcase B5.2: $p = 2$.: First suppose $t = 2$. Then we have the configuration of Figure 77 and $\sigma(u) = \pi + \phi^1 + \phi^2$, with $\phi^i \geq 5\pi/6$, for $i = 1, 2$ (as u is not of type $AA(x^i)$, Figure 30). As $|\mu^i| \leq l$ we have $\phi^i = 5\pi/6$ only if u is adjacent to w^i , for $i = 1, 2$. (Otherwise u is incident to fewer than ml




Figure 69:

arcs.) However if u is adjacent to w^i and incident to no further classes of arcs then we have a contradiction to Lemma 6.12. Hence $\phi^i \geq 7\pi/6$, for $i = 1$ or 2 and $\sigma(u) \geq 3\pi$. Therefore we may adjust angles so that $\kappa(u)$, $\kappa(v^i) \leq -\pi/9$.

Next suppose $t = 1$. Then we have the configuration of Figure 78, where $\phi^1 \geq \pi/3$ and $\phi^2 \geq 5\pi/6$ and $\sigma(u) = 5\pi/6 + \phi^1 + \phi^2$ (as u is not of type $AA(x^i)$, Figure 30). If $\phi^2 = 5\pi/6$ either w^2 is adjacent to β^1 and there are no arcs incident to u between ι^2 and μ^1 or u is adjacent to w^2 via the unique class of arcs incident to u between ι^2 and μ^1 . In both of these cases, since $|\mu^i| \leq l$ there must be arcs incident to u between ι^1 and μ^2 . If none of these arcs are boundary arcs then there must be at least two classes of arcs incident to u between ι^1 and μ^2 from which it follows that $\phi^1 \geq \pi$. On the other hand if there is a boundary arc incident to u between ι^1 and μ^2 then $\phi^1 \geq 4\pi/3$. Hence $\phi^2 = 5\pi/6$ implies that $\sigma(u) \geq 8\pi/3$.

If $\phi^2 > 5\pi/6$ then, as $\alpha_1^2 = \pi/3$, it follows that $\phi^2 \geq 7\pi/6$. If $\phi^2 = 7\pi/6$ then Δ_1^2 must be triangular and there can be at most 2 classes of arc incident to u between ι^2 and μ^1 , all of which must be interior. In this case there must be a class of arcs incident to u between ι^1 and μ^2 so $\phi^1 \geq 2\pi/3$. Hence $\phi^2 = 7\pi/6$ implies that $\sigma(u) \geq 8\pi/3$. If $\phi^2 > 7\pi/6$ then $\phi^2 \geq 5\pi/4$, with equality only if Δ_1^2 is triangular, there is only one class of arcs incident to u between ι^2 and μ^1 , and u is a vertex of type $AB(u_1)$. In this case there must be at least 2 classes of arc incident to u between ι^1 and μ^2 , so




Figure 70:

$\phi^1 \geq \pi$ and $\sigma(u) \geq 37\pi/12$. If $\phi^2 > 5\pi/4$ then $\phi^2 \geq 3\pi/2$ so, as $\phi^1 \geq \pi/3$, $\sigma(u) \geq 8\pi/3$. Therefore $t = 1$ implies $\sigma(u) \geq 8\pi/3$ and we may adjust angles so that $\kappa(u), \kappa(v^i) \leq -\pi/9$.

Finally suppose $t = 0$, in which case we have the configuration of figure 79. If u is incident to 3 more classes of arc (than those shown in Figure 79) then $\sigma(u) \geq 7\pi/3$. If u is incident to a boundary class of arcs then $\phi^i \geq 4\pi/3$, since $\alpha_1^i = \pi/3$, for $i = 1$ or 2 . In this case $\sigma(u) \geq 7\pi/3$ again. Hence, using Lemma 6.12, we may assume $\sigma(u) \geq 7\pi/3$ and adjust angles so that $\kappa(u), \kappa(v^i) \leq -\pi/9$.

Hence in Case B5 where all v^k 's are green we may adjust angles suitably. It follows, by symmetry, that this is also possible if all v^k 's are blue.

Case B6: there are q green and q blue components. Suppose that there are $q \geq 1$ green and $q \geq 1$ blue components, $G_1, \dots, G_q$ and $B_1, \dots, B_q$ respectively, in cyclic order round u . Let $g_j(A)$ be the number of vertices of type A in G_j and $b_j(A)$ be the number of vertices of type A in B_j , where A is X , Y or Z . Let $g_j = g_j(X) + g_j(Y) + g_j(Z)$ and $b_j = b_j(X) + b_j(Y) + b_j(Z)$; so

$$p = \sum_{j=1}^q g_j + b_j.$$

Furthermore let

$$g(A) = \sum_{j=1}^q g_j(A), \quad b(A) = \sum_{j=1}^q b_j(A) \quad \text{and}$$




Figure 71:

$$p(A) = g(A) + b(A), \text{ where } A = X, Y \text{ or } Z.$$

To complete the proof of case B of the theorem we aim to show that

$$(11.1) \quad \sigma(u) \geq 2\pi + p(Z)\pi/3 + (p+1)/24.$$

This is sufficient to allow a suitable angle adjustment to be made. In fact transferring angle of a from the corner of u in Δ_4^i to the corner of v^i in Δ_4^i , where

$$a = \begin{cases} 9\pi/24, & \text{if } T(v^i) = Z \\ \pi/24, & \text{if } T(v^i) = X \text{ or } Y \end{cases},$$

we have $\sigma(v^i) \geq 2\pi + \pi/24$, for all i , and

$$\begin{aligned} \sigma(u) &\geq 2\pi + p(Z)\pi/3 + (p+1)\pi/24 - p(Z)9\pi/24 - (p(X) + p(Y))\pi/24 \\ &= 2\pi + \pi/24. \end{aligned}$$

Hence we may adjust angles as described to obtain $\kappa(u), \kappa(v^i) \leq -\pi/24$, for all i .

A green or blue component $v^c, \dots, v^{c+d}$ is said to *begin* with v^c and *end* with v^{c+d} . We examine the effect of the type of the extremal vertices of a component on the sum of angles between its C -vertices.

If a component C begins with v^c and ends with v^{c+d} then the sum of angles around u between ι^c and ι^{c+1} , ι^{c+1} and ι^{c+2} , $\dots$, ι^{c+d-1} and ι^{c+d} is called the *angle* of the component and is denoted $\sigma(C)$.

Consider a green component G_j . If G_j begins with a Y or Z vertex then

$$\sigma(G_j) \geq (4\pi/3)(g_j(Y) + g_j(Z) - 1) + (2\pi/3)g_j(X).$$




Figure 72:

If G_j begins with an X vertex then

$$\sigma(G_j) \geq (4\pi/3)(g_j(Y) + g_j(Z)) + (2\pi/3)(g_j(X) - 1).$$

Now consider a blue component B_j . If B_j ends with an Y or Z vertex then

$$\sigma(B_j) \geq (4\pi/3)(b_j(Y) + b_j(Z) - 1) + (2\pi/3)b_j(X).$$

If B_j ends with an X vertex then

$$\sigma(B_j) \geq (4\pi/3)(b_j(Y) + b_j(Z)) + (2\pi/3)(b_j(X) - 1).$$

Let $e(S, T)$ denote the number of blue components which end with a vertex of type S and such that the following green component begins with a vertex of type T , where $S, T \in \{X, Y, Z\}$.

Then $q = e(X, X) + e(X, Y) + e(X, Z) + e(Y, X) + e(Y, Y) + e(Y, Z) + e(Z, X) + e(Z, Y) + e(Z, Z)$,

the number of blue components ending with a vertex of type A is

$$e(A, X) + e(A, Y) + e(A, Z)$$

and the number of green components beginning with a vertex of type A is

$$e(X, A) + e(Y, A) + e(Z, A).$$

Now $\sigma(u) = [\text{sum of angles of green components}]$

+ [sum of angles of blue components]

+ [sum of angles between green and blue components]

+ [sum of angles between blue and green components]




Figure 73:




Figure 74:




Figure 75:

$$\begin{aligned}
&\geq \left[(4\pi/3) \sum_{j=1}^q (g_j(Y) + g_j(Z) - 1) + (2\pi/3) \sum_{j=1}^q (g_j(X) - 1) \right. \\
&\quad + (2\pi/3) (e(X, Y) + e(Y, Y) + e(Z, Y) + e(X, Z) + e(Y, Z) + e(Z, Z)) \\
&\quad + (4\pi/3) (e(X, X) + e(Y, X) + e(Z, X)) \left. \right] \\
&\quad + \left[(4\pi/3) \sum_{j=1}^q (b_j(Y) + b_j(Z) - 1) + (2\pi/3) \sum_{j=1}^q (b_j(X) - 1) \right. \\
&\quad + (2\pi/3) (e(Y, X) + e(Y, Y) + e(Y, Z) + e(Z, X) + e(Z, Y) + e(Z, Z)) \\
&\quad + (4\pi/3) (e(X, X) + e(X, Y) + e(X, Z)) \left. \right] \\
&\quad + [q\pi/3] \\
&\quad + [(2\pi/3) e(X, X) + (4\pi/3) (e(X, Y) + e(X, Z) + e(Y, X) + e(Z, X)) \\
&\quad + 2\pi (e(Y, Y) + e(Y, Z) + e(Z, Y) + e(Z, Z))] \left. \right] \\
&= (2\pi/3) \sum_{j=1}^q (g_j(Y) + g_j(Z) + g_j(X) + b_j(Y) + b_j(Z) + b_j(X)) \\
&\quad + (2\pi/3) \sum_{j=1}^q (g_j(Y) + g_j(Z) + b_j(Y) + b_j(Z)) \\
&\quad - 4\pi q + (\pi/3)q \\
&\quad + (10\pi/3) (e(X, X) + e(X, Y) + e(X, Z) + \\
&\quad e(Y, X) + e(Y, Y) + e(Y, Z) + \\
&\quad e(Z, X) + e(Z, Y) + e(Z, Z)) \\
&= (2\pi/3) p - (\pi/3) q + (2\pi/3) (p(Z) + p(Y)).
\end{aligned}$$

Note that $p \geq 2q$ and if p is odd $p \geq 2q + 1$. Hence if $p \geq 5$ then $p - 3 \geq q$ and so

$$\begin{aligned}
\sigma(u) &\geq 2\pi + (p - 3) (2\pi/3) - (\pi/3) q + (2\pi/3) p(Z) \\
&\geq 2\pi + (\pi/3) (p - 3) + (2\pi/3) p(Z) \\
&\geq 2\pi + (\pi/9) (p + 1) + (\pi/3) p(Z),
\end{aligned}$$

since $3(p - 3) \geq (p + 1)$, when $p \geq 5$.




Figure 76:

Hence (11.1) holds when $p \geq 5$ and it remains to check the cases $2 \leq p \leq 4$.

Subcase B6.1: $p = 4$.

If $p = 4$ then $q = 1$ or 2 and

$$\sigma(u) \geq (8\pi/3) - (\pi/3)q + (2\pi/3)(p(Z) + p(Y)).$$

We shall show that, when $p = 4$, we have

$$\sigma(u) \geq 7\pi/3 + p(Z)(\pi/3)$$

so that

$$\sigma(u) \geq 2\pi + p(Z)(\pi/3) + (p+1)\pi/15$$

as required.

Subcase B6.1.1: If $p(Y)$ is non-zero then we have

$$\sigma(u) \geq 8\pi/3 + (2\pi/3)p(Z) > 7\pi/3 + (\pi/3)p(Z).$$

Subcase B6.1.2: Suppose $p(Y) = 0$. Then, if $p(Z) \geq 1$, we have

$$\sigma(u) \geq 2\pi + (2\pi/3)p(Z) \geq 7\pi/3 + (\pi/3)p(Z).$$




Figure 77:

If $p(Y) = p(Z) = 0$ and $q = 1$ then

$$\sigma(u) \geq 7\pi/3 = 7\pi/3 + (\pi/3)p(Z).$$

If $p(Y) = p(Z) = 0$ and $q = 2$ then we have the configuration of figure 80 where v^1 and v^3 are green, of type X , and v^2 and v^4 are blue of type X . (Note that, in Figure 80, ϕ^2 may equal zero if $x^2 = x^3$ and ϕ^4 may equal zero if $x^1 = x^4$.) Suppose first $\phi^2 = \phi^4 = 0$. As




Figure 78:

$\alpha_1^1 = \alpha_1^2 = \alpha_1^3 = \alpha_1^4 = \pi/3$ we have $\phi^1 \geq 2\pi/3$ and $\phi^3 \geq \pi/3$, or vice-versa, otherwise G is of type $E(2, 3, 6)$. Hence $\sum_{i=1}^4 \phi^i \geq \pi$. Now suppose $\phi^2 \neq 0$ or $\phi^4 \neq 0$, so $\phi^2 + \phi^4 \geq \pi/3$, and again $\sum_{i=1}^4 \phi^i \geq \pi$. We have then

$$\sigma(u) \geq 7\pi/3 = 7\pi/3 + p(Z)\pi/3.$$

This completes Subcase B6.1.

Subcase B6.2: $p = 3$.




Figure 79:

In this case $q = 1$ and

$$\sigma(u) \geq (5\pi/3) + (2\pi/3)(p(Z) + p(Y)).$$

If $p(Y) \neq 0$ then

$$\sigma(u) \geq (7\pi/3) + (2\pi/3)p(Z) \geq 2\pi + (\pi/3)p(Z) + \pi/3,$$

as required.




Figure 80:

Assume then that $p(Y) = 0$. If $p(Z) \geq 2$ then

$$\sigma(u) \geq (7\pi/3) + (\pi/3)p(Z) \geq 2\pi + (\pi/3)p(Z) + \pi/3.$$

Next consider the case $p(Z) = 1$. In this case either we have two blue and one green C -vertices adjacent to u or two green and one blue.

Subcase B6.2.1: $p(Y) = 0$, $p(Z) = 1$ and u is adjacent to two blue and one green C -vertices.




Figure 81:

Suppose first that $g(Z) = 1$ in which case $b(Z) = 0$ and $b(X) = 2$. Then we have the configuration of Figure 81. From Lemma 9.5 and Assumption 9.7, u must be incident to some class of arcs other than those shown in Figure 81. It follows that $\sum_{i=1}^3 \phi^i \geq 3\pi/2$ so $\sigma(u) \geq 8\pi/3$.

Now suppose $b(Z) = 1$ so that $g(X) = b(X) = 1$. This gives rise to two possible configurations depending on the cyclic order of the blue types X and Z around u . These two configurations correspond to Figures 82 and 83. In the configuration of Figure 82 it is possible that $\phi^3 = 0$ if $x^1 = x^3$. In this case u is incident to at least 2 classes of arcs not shown in Figure 82, using Lemma 9.5 and Assumption 9.7,




Figure 82:

as before. Therefore if $\phi^3 = 0$ we have $\sum \phi^i \geq 3\pi/2$. If $\phi^3 > 0$ then as $\phi^2 \geq 5\pi/6$ we have again $\sum \phi^i \geq 3\pi/2$. Hence $\sigma(u) \geq 8\pi/3$.

In the configuration of Figure 83 again u must be incident to at least one more class than shown and so as before $\sigma(u) \geq 8\pi/3$. Hence in Subcase B6.2.1 we have

$$\sigma(u) \geq 8\pi/3 = 2\pi + p(Z)\pi/3 + \pi/3.$$

Subcase B6.2.2: $p(Z) = 1$ and u is adjacent to two green and one blue C -vertices.

This subcase follows by symmetry from Subcase B6.2.1.

Subcase B6.2.3: The remaining Subcase of Subcase B6.2 is that in which $p(Y) = p(Z) = 0$. We consider only the case in which u is




Figure 83:

adjacent to two blue and one green C -vertices. The case in which u is adjacent to one blue and two green C -vertices follows by symmetry. We have $b_1(X) = 2$ and $g_1(X) = 1$ giving the configuration of Figure 84. If $\phi^2 = 0$ (that is $x^2 = x^3$) then u must be incident to at least 2 classes of arc which are not shown, so $\sum \phi^i \geq 4\pi/3$. If $\phi^2 > 0$ then $\phi^2 \geq \pi/3$. As u is incident to at least 7 classes of arcs $\sum \phi^i \geq 4\pi/3$ again. Hence, when $p(Z) = 0$, we have $\sigma(u) \geq 7\pi/3 = 2\pi + p(Z)\pi/3 + \pi/3$.

This completes Subcase B6.2.

Subcase B6.3: $p = 2$.:

In this case $q = 1$ and

$$\sigma(u) \geq \pi + 2\pi/3(p(Z) + p(Y)),$$




Figure 84:

with $p(Z) = 0, 1$ or 2 . We shall show that

$$(11.2) \quad \sigma(u) \geq 2\pi + p(Z)(\pi/3) + \pi/3 \geq 2\pi + p(Z)(\pi/3) + (p+1)\pi/9,$$

as required.

First suppose $p(Z) = 2$. Then (up to reflection) we have the configuration of Figure 85. As $|\mu^i| \leq l$, for $i = 1, 2$, it follows that there must be additional classes of arcs other than μ^1 and μ^2 then u is incident to at least 7 classes of arcs. If u is incident to 7 or more classes of arcs including exactly 2 boundary classes then either $\phi^2 \geq \pi/3$ and $\phi^1 \geq 5\pi/3$ or $\phi^2 \geq 2\pi/3$ and $\phi^1 \geq 4\pi/3$ or $\phi^2 \geq \pi$ and $\phi^1 \geq \pi$. Hence $\phi^1 + \phi^2 \geq 2\pi$. On the other hand if u is incident to a boundary class of arcs other than μ^1 and μ^2 then




Figure 85:

either $\phi^2 \geq \pi/3$ and $\phi^1 \geq 2\pi$ or $\phi^2 \geq \pi$ and $\phi^1 \geq \pi$. Therefore, in Figure 85, $\phi^1 + \phi^2 \geq 2\pi$ so $\sigma(u) \geq 3\pi = 2\pi + p(Z)\pi/3 + \pi/3$.

Now suppose $p = 2$ and $p(Z) = 1$. Assume that v^1 is blue and v^2 is green. If $b(Y) = g(Z) = 1$ or $b(Z) = g(Y) = 1$ then the argument above, for the case $p = p(Z) = 2$, can be used to show that $\sigma(u) \geq 2\pi + p(Z)\pi/3 + 2\pi/3$. Hence we may assume that either $b(X) = g(Z) = 1$ or $b(Z) = g(X) = 1$.

Subcase B6.3.1: $p = 2$, $p(Z) = 1$ and $b(X) = g(Z) = 1$.

We have the configuration of Figure 86. Again u must be incident to at least 3 further non-boundary classes of arc or 1 further boundary class.

If u is incident to three or more further classes of arcs then $\phi^1 \geq \pi/2$ and $\phi^2 \geq 4\pi/3$ or $\phi^1 \geq 5\pi/6$ and $\phi^2 \geq \pi$ or $\phi^1 \geq 7\pi/6$ and $\phi^2 \geq 2\pi/3$ or $\phi^1 \geq 3\pi/2$ and $\phi^2 \geq \pi/3$. Thus $\phi^1 + \phi^2 \geq 11\pi/6$.

If u is incident to more than one boundary class then either $\phi^1 \geq \pi/2$ and $\phi^2 \geq 4\pi/3$ (since $\alpha_1^1 = \pi/3$) or $\phi^1 \geq 3\pi/2$ and $\phi^2 \geq \pi/3$. Thus $\phi^1 + \phi^2 \geq 11\pi/6$ in Subcase B6.3.1. Therefore $\sigma(u) > 8\pi/3 \geq 2\pi + p(Z)\pi/3 + \pi/3$.

Subcase B6.3.2: $p = 2$, $p(Z) = 1$ and $b(Z) = g(X) = 1$.

This follows by symmetry from Subcase B6.3.1.

Finally we consider the case $p = 2$ and $p(Z) = 0$. Assume that v^1 is blue and v^2 is green. Then in the cases $g(Y) = b(Y) = 1$, $g(Y) = b(X) = 1$ and $g(X) = b(Y) = 1$ the arguments used above in the cases $p(Z) = 1$ and




Figure 86:

2 go through to give (11.2). Suppose then that $g(X) = b(X) = 1$, giving the configuration of Figure 87. Note that $\phi^1 = 0$ if and only if $x^1 = x^2$.

If u is incident to more than 6 classes of arc then $\phi^1 + \phi^2 \geq 5\pi/3$. If u is incident to a boundary class then either $\phi^2 = \pi/3$ and $\phi^1 = \pi$ or $\phi^2 = \pi/3$ and $\phi^1 \geq 4\pi/3$ or $\phi^2 \geq 2\pi/3$ and $\phi^1 \geq \pi$ or $\phi^2 \geq 5\pi/3$ and $\phi^1 \geq 0$. In all except the first case $\phi^1 + \phi^2 \geq 5\pi/3$, so $\sigma(u) \geq 7\pi/3 \geq 2\pi + p(Z)\pi/3 + \pi/3$. Also the first case, with $\phi^2 = \pi/3$ and $\phi^1 = \pi$, gives (up to reflection) configuration AA of Figure 30, in which the v^i are not C -vertices as angles have previously been adjusted, so this does not occur.

This completes consideration of Subcase B6.3 and shows that if u is incident to more than one vertex of type $Cj^\pm(v)$ then an angle adjustment may be made to give all vertices involved negative curvature of at most $-\pi/15$, hence completing Case B of the proof.

To complete the proof of the Proposition we consider configuration $D'6$, where u is of type $C0(v)$, v of type $CC^+(v)$ and $p = 1$ as in case A2.4. Let β' be the boundary component of Σ to which u is incident. Suppose first that β' is oriented, in Figure 61, from left to right. Assume first that x occurs as vertex u_1 in only 1 C -configuration (namely that appearing in Figure 61). Then in Case A2 with u of type $C0(v)$ either we have adjusted angles in Δ_5 , increasing α_7 and decreasing α_6 , so that $\kappa(x), \kappa(u) \leq -\pi/12$, or we found that u was of type $Dj(a)$, with $j = 0, 1, 2$ or 3. If we adjusted angles α_6 and α_7 in Case A2 then we make further angle adjustments, increasing α_4 by $\pi/18$ and decreasing α_5 and α_{10} by $\pi/36$, so that $\kappa(x), \kappa(u), \kappa(v) \leq -\pi/18$. On the other hand if u is of type $Dj(a)$ then, since β' is




Figure 87:

oriented from left to right, x is of type $Dj(b)$ and the only possibility is that $j = 3$. This gives rise to configuration $D6$, with v of type $D6(a)$ and u of type $D6(b)$.

Now with β' still oriented from left to right assume that x occurs as u_1 in $q > 1$ C -configurations. If the C -vertices of these C -configurations are $u, v_2, \dots, v_q$ then we have the configuration of Figure 88. In Case B above angles on $u, x, v_2, \dots, v_q$ were adjusted so that each of these vertices had curvature at most $-\pi/15$. Hence we may further adjust angles α_4 and α_5 so that $\kappa(u), \kappa(v) \leq -\pi/30$. If say v_i is also a vertex of type $D'6(u)$ and x is the corresponding vertex of type $D'6(x)$ then similar adjustments are made to the angles of v_i and the corresponding vertex of type $D'6(v)$.

Now suppose that β' is oriented from right to left in Figure 61. If y occurs as vertex u_1 in only one C -configuration then, from Case A2 with u of type $C0(v)$, either we may adjust angles α_8 and α_9 in Figure 61 so that $\kappa(u), \kappa(y) \leq -\pi/12$ or u is of type $Dj(a)$, with $j = 0, 1, 2$ or 3 . The latter cannot occur as β' is oriented from right to left. Therefore we may make further adjustments to angles $\alpha_4, \alpha_5, \alpha_8$ and α_9 so that $\kappa(u), \kappa(v), \kappa(y) \leq -\pi/18$. If on the other hand y occurs as vertex u_1 in more than one C -configuration then, as in the case of β' oriented from left to right, we may adjust angles α_4 and α_5 so that $\kappa(u), \kappa(v) \leq -\pi/30$. Furthermore we may make such adjustments simultaneously over all configurations of type $D'6$ in which y occurs as the vertex of type $D'6(y)$. This completes the proof of Proposition 11.1.

12. FINAL ANGLE ADJUSTMENT

We now make final adjustments to the assignment of angles to corners of vertices. These adjustments are carried out on those configurations of type Dj where




Figure 88:

the angle of type $Dj(a)$ has curvature greater than $-\pi/30$: that is, those arising in the proof of Proposition 11.1 in which angles were not adjusted. In all such configurations the vertex b is incident only to one C -vertex (namely the vertex $Dj(a)$). In configurations Dj , with $0 \leq j \leq 3$ we transfer angle of $\pi/3$ from the corner of b in Δ_1 to the corner of a in Δ_1 . Note that since configuration $D6$ contains configuration $D3$ this results in a transfer angle of $\pi/3$ from the corner of f in Δ_4 of $D6$ to the corner of b in Δ_4 of $D6$. Having made all the above adjustments to the angle assignments we make the following definitions.

Definition 12.1. Let $\mathcal{V}_N = \{v \in \mathcal{V} : \kappa(v) \leq -\pi/30\}$, let $\mathcal{V}_G = \mathcal{V}_N \cup \mathcal{V}_B$, let


$$\mathcal{V}_{k,B} = \{v \in \mathcal{V} : v \text{ is distance at most } k \text{ from } \mathcal{V}_B\},$$

and let $\mathcal{V}_Z = \mathcal{V} \setminus \mathcal{V}_G$ (where $\mathcal{V}_B$ is defined in Definition 6.4).

Lemma 12.2. *After final angle adjustment $\kappa(v) \leq 0$, for all $v \in \mathcal{V} \setminus \mathcal{V}_{1,B}$ and $\kappa(v) \leq \pi$, for all $v \in \mathcal{V}_{1,B}$. Furthermore if $v \in \mathcal{V}_Z \setminus \mathcal{V}_{1,B}$ then either*

- (i) v is of type $B4(a)$ or
- (ii) v is of type $Dj(a)$, $j = 0, \dots, 9$ or
- (iii) v is of type $Dj(b)$, $j = 0, \dots, 9$ or
- (iv) v is of type $Ij(a)$ or its mirror image, $j = 0, \dots, 6$.

Proof. It follows from Theorem 9.8 and Corollary 9.4 that before the angle adjustments made in the proof of Proposition 11.1 either v is interior and $\kappa(v) \leq 0$ or v is a boundary vertex and $\kappa(v) \leq \pi$, for all $v \in \mathcal{V}$. Thus $\kappa(v) \leq \pi$, for all $v \in \mathcal{V}_{1,B}$. From Theorem 9.8, if v is an interior vertex not in $\mathcal{V}_{1,B}$ then $\kappa(v) \leq -\pi/3$, so $v \in \mathcal{V}_N$. If v is a boundary vertex not in $\mathcal{V}_{1,B}$ and v meets $\partial\Sigma$ in 2 or more classes

Figure 89: Configuration $F1$

of arcs then Corollary 9.4 and Proposition 9.6 imply that either $v \in \mathcal{V}_N$ or (iv) of the Lemma holds. If v is a boundary vertex not in $\mathcal{V}_{1,B}$ and v meets $\partial\Sigma$ in exactly 1 class of arcs then, from Theorem 10.1, either $v \in \mathcal{V}_N$, (i) of the Lemma holds or v is a C -vertex. Moreover $\kappa(v) \leq 0$, when (i) or (iv) of the Lemma holds.

The angle adjustments made in the proof of Proposition 11.1 affect only C -vertices and their adjacent vertices of type u_1 and result in curvature being at most $-\pi/30$ for all vertices affected. Hence all of the above holds after these adjustments have been made with the following two exceptions. Firstly, if v is a boundary vertex not in $\mathcal{V}_{1,B}$ and v meets $\partial\Sigma$ in 2 or more classes of arcs then either $v \in \mathcal{V}_N$; or (iv) of the Lemma holds; or v is of type $Dj(b)$, $0 \leq j \leq 3$. Secondly, if v is a boundary vertex not in $\mathcal{V}_{1,B}$ and v meets $\partial\Sigma$ in exactly one class of arcs then either $v \in \mathcal{V}_N$ or (i), (ii) or (iii) of the Lemma holds.

Final angle adjustment affects only vertices in (ii) or (iii) of the Lemma, so all of the above still holds and in addition $\kappa(v) \leq 0$ when (ii) or (iii) of the Lemma holds.


Definition 12.3. We say that a vertex v is of type $Fj(a)$ if

- (i) v occurs as vertex a in a configuration $F1$ or $F2$ of Figures 89 or 90, respectively, and
- (ii) the configuration Fj in which v occurs contains no H^Λ -arcs.

Theorem 12.4. If $v \in \mathcal{V}$ then either

- (i) v is of type $F1(a)$ or $F2(a)$ or
- (ii) v is distance at most 3 from $\mathcal{V}_G$.

Proof. We may assume that $v \in \mathcal{V}_Z$ otherwise (ii) holds trivially. Let $v \in \mathcal{V}_Z$ be distance more than 1 from $\mathcal{V}_G$. From Theorem 12.2 it follows that v and all its adjacent vertices are of type $Ij(a)$ or its mirror image, with $j = 1, \dots, 6$, type $B4(a)$, type $Dj(a)$ or type $Dj(b)$ with $0 \leq j \leq 9$. Note that vertices of type $Dj(b)$, with $j = 4$ or 5 are also of type $B4(a)$ and that vertices of type $D6(b)$ are of type

Figure 90: Configuration $F2$

$D3(a)$. Thus once we've shown that a particular vertex is not of type $B4(a)$ or $D3(a)$ then this vertex cannot be of type $Dj(b)$, $j = 4, 5, 6$.

Case : v of type $B4(a)$:

We show that in this case v is either of type $F2(a)$ or is distance at most 2 from $\mathcal{V}_G$. Assume then that v is distance at least 3 from $\mathcal{V}_G$. Let the vertices incident to v be v_i of type $B4(a_i)$, $i = 1, \dots, 4$, as in Figure 39. Consider first v_1 . If v_1 were of type $Dj(a)$ then v would necessarily be of type $Dj(c)$ so, from Theorem 10.1, the boundary class incident at v would have width at most l . As this boundary class has width more than $2l$, v_1 is not of type $Dj(a)$. Furthermore v_1 is incident to a region Δ_1 with $\rho(\Delta_1) = 3$, so v_1 cannot be of type $Ij(a)$. Therefore v_1 can only be of type $Dj(b)$, $j = 0, \dots, 9$ or $B4(a)$. Moreover v_1 cannot be of type $D3(b)$, since such vertices are not incident to any vertex of type $B4(a)$. Suppose that $v_1 = Dj(b)$ with $0 \leq j \leq 2$. Since v is neither of type $D0(f)$ or $Dj(a)$ it follows that v is of type $Dj(d)$, forcing the configuration of Figure 91(a). Since we consider the case where v_1 is of type $B4(a)$ below and v_1 cannot be of type $D3(a)$, we need not consider the possibilities that v_1 is of type $Dj(b)$, with $j = 4, 5$ or 6 . If v_1 is of type $D7(b)$ then v must be of type $D7(d)$ and we obtain the configuration of Figure 91(b). If v_1 is of type $D8(b)$ then, as v is of type $B4(a)$, the only possibility is that v is the corresponding vertex of type $D8(d)$. However v_1 cannot then be of type $B4(a_1)$ so this cannot occur. If v is of type $D9(b)$ then v must be of type $D9(e)$. However in this case v_1 is a C -vertex and the class of boundary arcs incident at v can have width at most l . As v is of type $B4(a)$ this class is in fact of width at least $2l$, so v_1 cannot be of type $D9(b)$.

If v_1 is of type $B4(a)$ then we have the configuration of Figure 92(a), and we consider vertex v_2 . Clearly v_2 is not of type $D3(b)$. If v_2 is not adjacent to y then, as v_1 is of type $B4(a)$, y and v_2 must both be incident to the same boundary component, as shown in Figure 92(b). As v_2 must be incident to some further arcs and the angle ϕ in Figure 92(b) must be $\pi/3$ it follows that v_2 and v_3 are adjacent. By inspection of the regions incident at v_2 we see that v_2 cannot be of type $B4(a)$, $Ij(a)$ or $Dj(b)$. Therefore v_2 is of type $Dj(a)$ and we obtain the configuration of Figure 93(a). Suppose, on the other hand that, in Figure 92(a), v_2 is adjacent




Figure 91:




Figure 92:

to y . Then if v_2 is of type $B4(a)$ we obtain the configuration of Figure 94(a). If v_2 is of type $Dj(a)$ then, as v_2 is already adjacent to v , v_1 and y , it follows that v_2 must be of type $Dj(a)$ with $j \geq 4$ and v_3 must be incident to $\partial\Sigma$, as shown in Figure 93(b). It follows that v_2 must have type $D4(a)$ and y must have type $D4(c)$. Inspection of regions incident to v_2 , in Figure 92(a), show that v_2 cannot be adjacent to y and of type $Dj(b)$, with $0 \leq j \leq 9$.




Figure 93:




Figure 94:

We conclude that if v is of type $B4(a)$ then we have one of the configurations of Figures 91(a), 91(b), 93(a), 93(b) or 94(a). In Figures 91(a), 91(b) and 93(a) the vertex v_2 is a C -vertex and this forces the boundary class incident at v_3 in Figures 91(a), 91(b) and 93(a) to have width at most l . Hence v_3 can only be of type $Dj(b)$, with $0 \leq j \leq 3$. Consideration of its incident regions shows that v_3 cannot be any of these types so Figures

91(a), 91(b) and 93(a) do not occur. In Figure 93(b) as, by assumption, v is distance more than 2 from $\mathcal{V}_G$ we have $y \in \mathcal{V}_Z$. However, as y is of type $D4(c)$ the boundary class incident at y in Figure 93(b) has width at most l . Hence y must be of type $Dj(b)$, with $0 \leq j \leq 3$ or $7 \leq j \leq 9$. As this is clearly not the case the configuration of Figure 93(b) does not arise.

Hence the only possible configuration is that of Figure 94(a). Clearly v_3 is not then of type $Ij(a)$. If v_3 were of type $Dj(a)$ then the boundary class incident at v_2 could have width at most l , whereas it has, in this case, width greater than $2l$. As above v_3 is not of type $Dj(b)$ so v_3 must be of type $B4(a)$. A similar argument shows that v_4 is of type $B4(a)$. This implies that v is of type $F2(a)$.

Case : v of type $Dj(a)$, $0 \leq j \leq 9$:

Now consider the case where v is of type $Dj(a)$, for $0 \leq j \leq 9$, and is distance more than 1 from $\mathcal{V}_G$. Let the vertices adjacent to v be v_1, v_2 and v_3 of type $Dj(b), Dj(d)$ and $Dj(c)$ respectively. Consideration of regions to which v_i is incident, shows that v_i cannot be of type $Ik(a)$, for all i, j and k . Furthermore v_3 cannot be of type $B4(a)$ or a C -vertex, as v_3 meets $\partial\Sigma$ in a class of width at most l . Hence v_3 must be of type $Dj(b)$, with $0 \leq j \leq 3$.

Subcase : v of type $D0(a)$:

Suppose v_3 is of type $D0(b)$. Then consideration of regions incident to v_3 shows that v_2 and v_3 are adjacent and that v_2 is the vertex of type $D0(a)$ corresponding to the type $D0(b)$ vertex v_3 . We therefore have the configuration of Figure 94(b), including the dashed vertex and arc. However in this figure both boundary classes incident to v_1 are of width less than l , so v_1 is incident to fewer than ml arcs, a contradiction. A similar argument shows that v_3 cannot be of type $Dj(b)$ with $j = 1$ or 2 . If v_3 is of type $D3(b)$ then v must be of type $D3(a)$, which is not the case. Therefore v cannot be of type $D0(a)$.

Subcase: v of type $Dj(a)$, $j = 1$ or 2 :

The argument of the previous case applies to show that v cannot be of these types.

Subcase: v of type $D3(a)$:

If v_3 is of type $Dj(b)$ with $j \leq 2$ then v_2 must be of type $Dj(a)$, with $j \leq 2$, and v_2 and v_3 must be adjacent. Here v_3 is incident to fewer than ml arcs, so this cannot happen. If v_3 is of type $D3(b)$ then v must be the corresponding vertex of type $D3(a)$, but this is impossible, given the orientation of boundary components and the existence of a vertex of type $D3(f)$. Therefore v cannot be of type $D3(a)$.

Subcase : v of type $D4(a)$ or $D5(a)$:

In this case v_1 is of type $B4(a)$ and hence, using the Case: v of type $B4(a)$, is a distance at most 2 from $\mathcal{V}_G$. Therefore v is distance at most 3 from $\mathcal{V}_G$.

Subcase : v of type $D6(a)$:

As v_1 is of type $D3(a)$ the distance of v_1 from $\mathcal{V}_G$ is at most 1 so that v is distance at most 2 from $\mathcal{V}_G$.

Subcase : v of type $D7(a)$:

Since v_1 is of type $CC^\pm(v)$ it follows that v_2 is of type $Dj(b)$ and v_2 cannot be a vertex of type $B4(a)$ or a C -vertex, so $0 \leq j \leq 3$. It is easy to check that v_2 cannot be of these types, so v cannot be of type $D7(a)$.

Subcase : v of type $D8(a)$:

If v_3 is of type $Dj(b)$, with $0 \leq j \leq 2$, then v_3 is adjacent to v_2 which must therefore be of type $B4(a)$. In this case v is distance at most 3 from $\mathcal{V}_G$. If v_3 is of type $D3(b)$ then v is of type $D3(a)$, which it cannot be. Therefore if v is of type $D8(a)$ then v is distance at most 3 from $\mathcal{V}_G$.

Subcase : v of type $D9(a)$:

If v_3 is of type $Dj(b)$, with $j \leq 3$, then v_2 is incident to fewer than ml arcs. Therefore v is not of type $D9(a)$.

Case : v of type $Dj(b)$, with $0 \leq j \leq 3$:

The vertex v is adjacent to a vertex of type $Dj(a)$, with $0 \leq j \leq 3$, which, from the above, is distance at most 1 from $\mathcal{V}_G$. Hence v is distance at most 2 from $\mathcal{V}_G$.

Case : v of type $Dj(b)$, with $4 \leq j \leq 6$:

In this case v is of type $B4(a)$ or $D3(a)$ so, from the above it follows that v is distance at most 2 from $\mathcal{V}_G$.

Case : v of type $Dj(b)$, with $7 \leq j \leq 9$:

If $j = 7$ or 9 then $Dj(a)$ is distance at most 1 from $\mathcal{V}_G$ so v is distance at most 2 from $\mathcal{V}_G$. If $j = 8$ then, from the case v of type $D8(a)$, if v is distance more than 2 from $\mathcal{V}_G$ then the corresponding vertex of type $D8(d)$ is also of type $B4(a)$, so is distance at most 2 from $\mathcal{V}_G$. Hence v is distance at most 3 from $\mathcal{V}_G$.

Case : v of type $Ij(a)$:

Assume v is distance more than 3 from $\mathcal{V}_G$. As vertices of type $B4(a)$ and $Dj(a)$ are neither adjacent nor paired to a vertex of type $Ij(a)$ it follows that v can be adjacent or paired only to vertices of type $Ij(a)$ or $Dj(b)$ with $j = 0, \dots, 9$. If v is adjacent or paired to a vertex of type $Dj(b)$ then, by inspection, $0 \leq j \leq 3$ and from the above v is distance at most 3 from $\mathcal{V}_G$, a contradiction. Hence all vertices to which v is adjacent or paired are of type $Ij(a)$. It follows that v is of type $F1(a)$, Figure 89, and the result follows.

This completes the proof of the theorem.

We partition $\mathcal{V}_Z$ into $\mathcal{V}_F = \{v \in \mathcal{V}_Z : v \text{ is of type } F1(a) \text{ or } F2(a)\}$ and $\mathcal{V}_D = \mathcal{V}_Z \setminus \mathcal{V}_F$.

Lemma 12.5. *The number of vertices in $\mathcal{V}_D$ is bounded above by*

$$|\mathcal{V}_D| \leq (ml)^3 |\mathcal{V}_G|.$$

Proof. If $v \in \mathcal{V}_D$, then $v \notin \mathcal{V}_G$ but the distance of v from $\mathcal{V}_G$ is at most 3 (Theorem 12.4).

13. ISOPERIMETRY

Let $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ be a positive 4-partition of $(h, n, t, p) \in \mathbb{Z}^4$, $\mathcal{L}$ a consistent system of parameters and $\mathbf{z}$ a special element of $(H^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h})$ and let $\mathcal{H}$ be the homogeneous system of parameters associated with $\mathbf{z}$ and $\mathcal{L}$. Then $\bar{\mathbf{z}}$ is a special element of $(H^\Lambda, \mathcal{H}, \mathbf{n}, \mathbf{h})$. Throughout this section Γ is assumed to be a reduced, minimalistic picture over G on a compact surface Σ of type $(\mathbf{n}, \mathbf{t}, \mathbf{p})$ with boundary partition $\mathbf{b} = (b_1, \dots, b_{W_1})$, prime labels $\bar{\mathbf{z}}$ and labelled by $\hat{\alpha}(\bar{\mathbf{z}})$, for some solution α to $\mathcal{H}$. Let $\mathcal{D}$ denote the set of boundary components of Σ . Then

$$\begin{aligned}
& \sum_{\mathcal{V}} \kappa(v) + \sum_{\mathcal{R}} \kappa(\Delta) + \sum_{\mathcal{D}} \kappa(\beta) \\
&= \sum_{\mathcal{V}} (2\pi - \sigma(v)) + \sum_{\mathcal{R}} (\sigma(\Delta) - t(\Delta)\pi + 2\pi\chi(\Delta)) \\
&\quad + \sum_{\mathcal{D}} -\sigma(\beta) \\
&= \sum_{\mathcal{V}} 2\pi + \sum_{\mathcal{R}} (-t(\Delta)\pi + 2\pi\chi(\Delta)) \\
&= 2\pi \left(|\mathcal{V}| + \sum_{\mathcal{R}} (-(1/2)t(\Delta) + \chi(\Delta)) \right) \\
&= 2\pi \left(|\mathcal{V}| - |\mathcal{A}| + \sum_{\mathcal{R}} \chi(\Delta) \right) = 2\pi\chi(\Sigma).
\end{aligned}$$

That is

$$(13.1) \quad \sum_{v \in \mathcal{V}} \kappa(v) + \sum_{\Delta \in \mathcal{R}} \kappa(\Delta) + \sum_{\beta \in \mathcal{D}} \kappa(\beta) = 2\pi\chi(\Sigma)$$

Definition 13.1. The Z -subspace of Γ is the union of all maximal designated corridors of Γ .

Definition 13.2. Let $C = (D, \Gamma_D, \mu)$ be a maximal designated simple j -corridor, where $j = 0, 1$ or 2 . Assume that the image of C under μ is a disk, let c be the width of C and let $\Gamma_\beta = \Gamma_D(0)$ or $\Gamma_D(c-1)$. If Γ_β is a picture of type square- j then $\mu(\Gamma_\beta)$ is called an j -corridor-boundary. Let E be a maximal designated 2-corridor with foundation $C_0 = (D_0, \Gamma_{D_0}, \mu_0)$ and extension $C_1 = (D_1, \Gamma_{D_1}, \mu_1)$ and with image a disk. Then E has one compartment Γ_0 which is not a compartment of C_1 and one compartment Γ_1 which is not a compartment of C_0 . The image $\mu(\Gamma_i)$ is called a 2-corridor-boundary, for $i = 1, 2$. A j -corridor-boundary with $j = 0, 1$ or 2 is called a corridor-boundary.

The image of every corridor is an annulus a Möbius band or a disk. Every corridor, of width at least 2, which has image a disk has two corridor-boundaries or consists of a unique arc of type I .

Proposition 13.3. The number of 0-corridor-boundaries is bounded above by

$$3(ml(ml+1)^3|\mathcal{V}_G| + W_0 - 2\chi(\Sigma)) + (6ml + 10)W_1.$$

Proof. Let $\mathcal{R}_1 = \{\Delta \in \mathcal{R} : \Delta \text{ is not contained in the } Z\text{-subspace of } \Gamma\}$. For $\Delta \in \mathcal{R}_1$ define $I(\Delta) = \text{number of } 0\text{-corridor-boundaries meeting } \partial\Delta$. Let $\mathcal{R}_2 = \{\Delta \in \mathcal{R}_1 : 0 < I(\Delta) \leq \rho(\Delta)\}$. Then

$$(13.2) \quad \sum_{\Delta \in \mathcal{R}_2} I(\Delta) \leq \sum_{\Delta \in \mathcal{R}_2} \rho(\Delta).$$

Let

$$\mathcal{R}_3 = \{\Delta \in \mathcal{R}_1 : I(\Delta) > \rho(\Delta) \text{ and } \Delta \text{ is not collapsible}\}$$

and

$$\mathcal{R}_4 = \{\Delta \in \mathcal{R}_1 : I(\Delta) > \rho(\Delta) \text{ and } \Delta \text{ is collapsible}\}.$$

Then $\Delta \in \mathcal{R}_3 \cup \mathcal{R}_4$ implies that $\varepsilon(\Delta) \geq I(\Delta) > \rho(\Delta)$. Let $\Delta \in \mathcal{R}_3$. If $\rho(\Delta) \geq 1$ then

$$\kappa(\Delta) \leq (\rho(\Delta) - \varepsilon(\Delta))\pi/2 \leq (\rho(\Delta) - I(\Delta)/3)\pi,$$

using the proof of Lemma 9.2. If $\rho(\Delta) = 0$, then

$$\kappa(\Delta) \leq (-\varepsilon(\Delta) + 2\chi(\Delta))\pi.$$

If $\rho(\Delta) = 0$ and $\chi(\Delta) \leq 0$ then

$$\begin{aligned} \kappa(\Delta) &\leq -\varepsilon(\Delta)\pi \leq -I(\Delta)\pi < -I(\Delta)\pi/3 \\ &= (\rho(\Delta) - I(\Delta)/3)\pi. \end{aligned}$$

If $\rho(\Delta) = 0$ and $\chi(\Delta) = 1$, then $t(\Delta) = \varepsilon(\Delta) \geq 2$, since Γ has no trivial boundary labels. If $\varepsilon(\Delta) = 2$ then Δ is collapsible so $\Delta \notin \mathcal{R}_3$. Hence $\varepsilon(\Delta) \geq 3$ and

$$\kappa(\Delta) \leq (-\varepsilon(\Delta) + 2)\pi \leq -\varepsilon(\Delta)\pi/3 \leq (\rho(\Delta) - I(\Delta)/3)\pi.$$

Therefore if $\Delta \in \mathcal{R}_3$ then

$$\kappa(\Delta) \leq (\rho(\Delta) - I(\Delta)/3)\pi.$$

As $\kappa(\Delta) \leq 0$, for all regions Δ (Lemma 9.2), we have

$$\sum_{\mathcal{R}} \kappa(\Delta) \leq \sum_{\Delta \in \mathcal{R}_3} \kappa(\Delta) \leq \sum_{\Delta \in \mathcal{R}_3} (\rho(\Delta) - I(\Delta)/3)\pi.$$

Also, from Lemma 12.2 and Lemma 6.6

$$\sum_v \kappa(v) \leq |\mathcal{V}_{1,B}|\pi \leq 2(ml+1)W_1\pi,$$

as the number of vertices at distance at most 1 from an H^Λ -vertex is at most ml .

Hence, using (13.1) and Lemma 9.1,

$$2\pi\chi(\Sigma) \leq 2(ml+1)W_1\pi + \sum_{\Delta \in \mathcal{R}_3} (\rho(\Delta) - I(\Delta)/3)\pi + W_0\pi,$$

so

$$(13.3) \quad \sum_{\Delta \in \mathcal{R}_3} I(\Delta) \leq 6(ml+1)W_1 + \sum_{\Delta \in \mathcal{R}_3} 3\rho(\Delta) + 3W_0 - 6\chi(\Sigma)$$

Now let $\Delta \in \mathcal{R}_4$. Then $\Delta \in \mathcal{R}_1$ implies that $\partial\Delta$ contains an H^Λ -arc so there are at most $2W_1$ regions in $\mathcal{R}_4$. As Δ is collapsible we have $\varepsilon(\Delta) = t(\Delta) = 2$ and so $I(\Delta) \leq 2$. Therefore

$$(13.4) \quad \sum_{\Delta \in \mathcal{R}_4} I(\Delta) \leq 4W_1.$$

Now consider a vertex v which meets $\partial\Delta$, for some region $\Delta \in \mathcal{R}_2 \cup \mathcal{R}_3$. Then v cannot be of type $F2(a)$ as no region incident to a in $F2$ meets a 0-boundary. If v is of type $F1(a)$ then Δ and v both belong to the same j -corridor, contrary to choice of $\Delta \in \mathcal{R}_1$, unless v is an H^Λ -vertex (that is $v \in \mathcal{V}_B \subseteq \mathcal{V}_G$). Hence, from Theorem 12.4, v is distance at most 3 from $\mathcal{V}_G$. Every vertex has degree ml , so if a vertex is paired to p vertices and joined to q vertices then $p + q \leq ml$. Hence the number of vertices which are distance at most 3 from $|\mathcal{V}_G|$ is less than or equal to $(ml + 1)^3 |\mathcal{V}_G|$. A vertex meets at most ml regions so

$$(13.5) \quad \sum_{\Delta \in \mathcal{R}_2 \cup \mathcal{R}_3} \rho(\Delta) \leq ml(ml + 1)^3 |\mathcal{V}_G|.$$

From (13.2), (13.3) and (13.4), it follows that

$$\sum_{\Delta \in \mathcal{R}_2 \cup \mathcal{R}_3 \cup \mathcal{R}_4} I(\Delta) \leq \sum_{\Delta \in \mathcal{R}_2 \cup \mathcal{R}_3} 3\rho(\Delta) + 6(ml + 1)W_1 + 3W_0 - 6\chi(\Sigma) + 4W_1.$$

As the number of 0-corridor-boundaries is at most $\sum_{\Delta \in \mathcal{R}_2 \cup \mathcal{R}_3 \cup \mathcal{R}_4} I(\Delta)$ the result follows from (13.5).

Lemma 13.4. *There are fewer than*

$$n + 3((ml + 1)^4 |\mathcal{V}_G| + W_0 - 2\chi(\Sigma)) + (6ml + 10)W_1$$

maximal designated corridors of Γ .

Proof. Suppose C is a maximal designated j -corridor and let C_D be the image of the C . If C_D is an annulus or Möbius band then C_D is a connected component of Σ . As Σ has n boundary components there are at most n maximal designated corridors of this form. Assume then that C_D is not an annulus or Möbius band, so C_D is a disk. If $j = 0$ then C has a 0-corridor-boundary. From Proposition 13.3 there are at most

$$c_0 = 3(ml(ml + 1)^3 |\mathcal{V}_G| + W_0 - 2\chi(\Sigma)) + (6ml + 10)W_1$$

corridors with a 0-corridor-boundary. If $j \geq 1$ then either C has 2 0-corridor boundaries or a j -corridor-boundary. If C has a j -corridor-boundary Γ_β with vertex v then v cannot be of type $Fj(a)$. Hence v is distance at most 3 from $\mathcal{V}_G$, from Theorem 12.4. The number of vertices distance at most 3 from $\mathcal{V}_G$ is less than or equal to $c_1 = (ml + 1)^3 |\mathcal{V}_G|$ so there are at most c_1 corridors with a j -corridor-boundary, $j \geq 1$. The result follows.

Lemma 13.5. *The number of vertices in $\mathcal{V}_N$ is bounded above by*

$$|\mathcal{V}_N| \leq 30(2(ml + 1)W_1 + W_0 - 2\chi(\Sigma)).$$

Proof. From (13.1) and Lemma 12.2 we have

$$2\pi\chi(\Sigma) \leq \sum_{v \in \mathcal{V}_N} \kappa(v) + \sum_{v \in \mathcal{V}_{1,B}} \kappa(v) + \sum_{\Delta \in \mathcal{R}} \kappa(\Delta) + \sum_{\beta \in \mathcal{D}} \kappa(\beta)$$

and, using Lemma 6.6, Lemma 9.1 and Lemma 9.2, this gives

$$2\pi\chi(\Sigma) \leq -(\pi/30)|\mathcal{V}_N| + 2(ml + 1)W_1\pi + W_0\pi,$$

from which the result follows.

Definition 13.6. We define

$$\begin{aligned}\mathcal{B}_1 &= \mathcal{B}_1(\mathbf{z}) = 2W_1 + 30(2(ml+1)W_1 + W_0 - 2\chi(\Sigma)) \text{ and} \\ \mathcal{B} &= \mathcal{B}(\mathbf{z}) = 2W_1 + ml((ml)^3 + 1)\mathcal{B}_1 + M(\mathbf{z})(n + 3((ml+1)^4\mathcal{B}_1 + W_0 - 2\chi(\Sigma)) \\ &\quad + (6ml + 10)W_1).\end{aligned}$$

Theorem 13.7. Let Γ be a picture over G as described at the beginning of this section. Then the number of arcs $|\mathcal{A}|$ of Γ satisfies $|\mathcal{A}| \leq \mathcal{B}$.

Proof. Let y be an arc of Γ and assume first that y is not contained in the Z -subspace. If y is incident to no vertex then the conditions on Γ imply that y is an H^Λ -arc. If y is incident to a vertex v then $v \in \mathcal{V}_G \cup \mathcal{V}_D$, as if v is of type $Fj(a)$ then y is contained in the Z -subspace. Hence the number of arcs of Γ which are not contained in the Z -subspace of Γ is at most

$$2W_1 + ml|\mathcal{V}_G \cup \mathcal{V}_D|.$$

From Corollary 7.11 and Lemma 13.4 there are at most

$$M(\mathbf{z})(n + 3((ml+1)^4|\mathcal{V}_G| + W_0 - 2\chi(\Sigma)) + (6ml + 4)W_1)$$

arcs in the Z -subspace (note that $M(\mathbf{z}) = M(\bar{\mathbf{z}})$: see Section 3.1). The result now follows from Lemma 12.5 and Lemma 13.5.

14. PROOF OF THEOREM 5.9

Let $\mathcal{N} = \mathcal{N}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ and $l = l(r)$. Consider first the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem when $\mathbf{h}$ is a partition of $h = 0$. In this case the problem reduces to the $\mathcal{Q}$ -problem and, as in Example 5.7, under the current hypotheses is solvable in $(G, A, B, H_i : i \in I)$.

We may therefore assume that $\mathbf{h}$ is a partition of a positive integer h , that $\mathbf{n}$ is a partition of a positive integer and that there is some i such that $h_i n_i > 0$, since otherwise the system reduces to the $\mathcal{Q}$ -problem again. It follows that the positive 4-partition, of length 1, $(1, 1, 0, 0) \in \mathcal{N}$. The hypothesis on $(A, B, H_i : i \in I)$ now implies that A, B, H_i all have soluble word problem. It follows from Lemma 5.5 that, in $(G, A, B, H_i : i \in I)$, the $\mathcal{GE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is solvable if the $\mathcal{SGE}(\mathbf{h}'', \mathbf{n}'', \mathbf{t}'', \mathbf{p}'')$ -problem is solvable for all positive 4-partitions $(\mathbf{h}'', \mathbf{n}'', \mathbf{t}'', \mathbf{p}'')$ of (h, n'', t, p) such that $n'' \in \mathbb{Z}$ and $(\mathbf{h}'', \mathbf{n}'', \mathbf{t}'', \mathbf{p}'') \leq (\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$. Such partitions $(\mathbf{h}'', \mathbf{n}'', \mathbf{t}'', \mathbf{p}'')$ are in $\mathcal{N}$. It therefore suffices, using the argument at the beginning of the proof of Theorem 5.8 in Section 5.1, to show that the $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is solvable in $(G, A, B, H_i : i \in I)$, where $\mathbf{n}$ is a partition of an integer $n \geq 1$.

Let i_A, i_B and i_* be symbols not in I and let $J = I \cup \{i_A, i_B\}$. Let $K = J \cup \{i_*\}$, $X_{i_*} = \{i_A, i_B\}$ and $X_j = \{j\}$, for all $j \in J$. Set $H_{i_A} = A$ and $H_{i_B} = B$ so $H_{*, i_*} = H$ and, with $\mathbf{s}_{i_*} = \{s\}$ we have $G_{i_*} = G$. Set $\mathbf{s}_j = \emptyset$, for all $j \in J$. Then $H_{*, i_A} = A$, $H_{*, i_B} = B$, $H_{*, i} = H_i$, for all $i \in I$ and $G_j = H_{*, j}$, for all $j \in J$. If $\mathcal{L}$ is a consistent system of parameters and

$$\mathbf{z} \in \left(\bigcup_{k \in K} H_{*, k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h} \right)$$

then the equation Q associated to $\mathbf{z}$ has environment

$$((X_k)_{k \in K}; (H_j)_{j \in J}; (G_k)_{k \in K})^\Lambda.$$

The $\mathcal{SGE}(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$ -problem is solvable if, given such $\mathbf{z}$, which is in addition special, there exists an algorithm to decide whether or not Q has a solution and if so to output one.

Assume then that $\mathbf{z}$ is a special element of $\left(\dot{\bigcup}_{k \in K} H_{*,k}^\Lambda, \mathcal{L}, \mathbf{n}, \mathbf{h}\right)$. Reordering the elements of $\mathbf{z}$ if necessary we may assume that we have a positive integer u such that $\mathbf{z}$ is supported over $\mathbf{n}$ with basis $(a_1, \dots, a_k)$, where $a_i = i_*$, for $i = 1, \dots, u$ and $a_i \in J$, for $i > u$. Then $\mathbf{z} = \mathbf{z}_1, \mathbf{z}_2$, $\mathbf{h} = \mathbf{h}_1, \mathbf{h}_2$ and $\mathbf{n} = \mathbf{n}_1, \mathbf{n}_2$ where

$$\begin{aligned} \mathbf{z}_1 &= (z_1, \dots, z_{\mu_{u+1}}) \in (H^\Lambda, \mathcal{L}, \mathbf{n}_1, \mathbf{h}_1), \\ \mathbf{z}_2 &= (z_{\mu_{u+1}+1}, \dots, z_n) \in \left(A^\Lambda \dot{\cup} B^\Lambda \dot{\cup} \bigcup_{i \in I} H_i^\Lambda, \mathcal{L}, \mathbf{n}_2, \mathbf{h}_2\right), \\ \mathbf{h}_1 &= (h_1, \dots, h_u), \\ \mathbf{h}_2 &= (h_{u+1}, \dots, h_k), \\ \mathbf{n}_1 &= (n_1, \dots, n_u) \text{ and} \\ \mathbf{n}_2 &= (n_{u+1}, \dots, n_k), \end{aligned}$$

where $\mu_i = \mu_i(\mathbf{n})$ as defined in (3.9). Define

$$\begin{aligned} \mathbf{t}_1 &= (t_1, \dots, t_u), \\ \mathbf{t}_2 &= (t_{u+1}, \dots, t_k), \\ \mathbf{p}_1 &= (p_1, \dots, p_u) \text{ and} \\ \mathbf{p}_2 &= (p_{u+1}, \dots, p_k), \end{aligned}$$

so that $\mathbf{t} = \mathbf{t}_1, \mathbf{t}_2$ and $\mathbf{p} = \mathbf{p}_1, \mathbf{p}_2$. Then

$$(\mathbf{h}_i, \mathbf{n}_i, \mathbf{t}_i, \mathbf{p}_i) \in \mathcal{N}, \text{ for } i = 1, 2.$$

Let

$$Q = Q(\mathbf{z}, \mathcal{L}, \mathbf{n}, \mathbf{t}, \mathbf{p}) = (\mathbf{q} = 1, \beta, \mathcal{L}),$$

let

$$\begin{aligned} \mathbf{q}_1 &= (q(\xi_j, n_j, t_j, p_j) : j = 1, \dots, u) \text{ and } \mathbf{q}_2 = (q(\xi_j, n_j, t_j, p_j) : j = u+1, \dots, k), \\ \beta_i &= \beta|_{L_D(\mathbf{q}_i)} \text{ and } Q_i = (\mathbf{q}_i = 1, \beta_i, \mathcal{L}), \text{ for } i = 1, 2. \end{aligned}$$

Let $Q_{\mathcal{H}}$ be the homogeneous equation associated to $\mathbf{z}$ and $\mathcal{L}$ and suppose that $Q_{\mathcal{H}} = (\mathbf{q} = 1, \beta_{\mathcal{H}}, \mathcal{H})$. Define $\mathcal{H}_i$ to be the set of elements of $\mathcal{H}$ containing parameters which occur in $\overline{\mathbf{z}_i}$ and

$$Q_{\mathcal{H}_i} = (\mathbf{q}_i = 1, \beta_{\mathcal{H}_i}, \mathcal{H}_i),$$

where $\beta_{\mathcal{H}_i} = \beta_{\mathcal{H}}|_{L_D(\mathbf{q}_i)}$, for $i = 1, 2$. Then $Q_{\mathcal{H}_1}$ is the homogeneous equation associated to $\mathbf{z}_1$. Up to relabelling of parameters it is also the case that $Q_{\mathcal{H}_2}$ is the homogeneous equation associated to $\mathbf{z}_2$. As $\mathbf{z}$ is special so is $\mathbf{z}_i$ and hence also, using Lemma 4.11, is $Q_{\mathcal{H}_i}$. The equations Q_1 and $Q_{\mathcal{H}_1}$ have environment

$$(X_{i_*}; H_{i_A}, H_{i_B}; G_{i_*}) = (X_{i_*}; A, B; G)^\Lambda,$$

while Q_2 and $Q_{\mathcal{H}_2}$ have environment

$$((X_j)_{j \in J}; (H_j)_{j \in J}; (H_j)_{j \in J})^\Lambda.$$

If (ϕ, α) is a solution to the equation Q then define

$$\phi_i = \phi|_{L(\mathbf{q}_i)}, \text{ for } i = 1, 2.$$

Then (ϕ_1, α) is a solution to Q_1 and (ϕ_2, α) is a solution to Q_2 . Conversely suppose we are given solutions (ϕ_i, α) to Q_i , for $i = 1, 2$. Define

$$(14.1) \quad \begin{aligned} \phi : F(L(\mathbf{q})) &\longrightarrow H * A * B * *_{i \in I} H_i \\ \text{by } \phi(x) &= \phi_i(x), \text{ for } x \in L(\mathbf{q}_i). \end{aligned}$$

Then (ϕ, α) is a solution to Q .

Assuming that a solution (ϕ, α) to Q exists then there is a picture Γ on a surface Σ of type $(\mathbf{n}_1, \mathbf{t}_1, \mathbf{p}_1)$ such that (α, Γ) corresponds to the solution (ϕ_1, α) to Q_1 . A minimal picture on Σ with the same boundary labels as Γ then exists, and corresponds to a, possibly different, solution to Q_1 . Together with ϕ_2 this solution to Q_1 can be used, as in (14.1), to construct a new solution to Q . Hence we may assume that Γ is a minimal picture. (See Definition 6.16, Definition 6.9 and Corollary 6.15.) Let Γ_s be a Z -reduced picture obtained from Γ by Z -cancellation. Then Γ_s is reduced and minimalistic and, using Lemma 7.17(ii), determines a solution α_s to $\mathcal{H}_1$ such that (α_s, Γ_s) corresponds to some solution (ϕ_s, α_s) to $Q_{\mathcal{H}_1}$: that is Γ_s is a basic picture for $(\mathbf{z}_1, \mathcal{L})$. From Theorem 7.19 there exists a solution α_0 to $\mathcal{L} \cup \mathcal{L}(\Gamma_s, P)$, where P is the subgraph of some path in the Z -graph $G(\mathbf{z}_1)$ of $\mathbf{z}_1$, with initial vertex $v(\Gamma_s)$. Furthermore α_0 may be chosen so that $\alpha_0(\lambda) = \alpha(\lambda)$, for all $\lambda \in \Lambda$ occuring in Q .

Since partisan regions are not involved in Z -cancellation, the partisan regions of Γ and Γ_s are the same. Let Δ be the closure of a partisan region of Γ . Assume that Δ is a V -region, where $V = A$ or B and that Δ is the sum of $t' = t'(\Delta)$ torii with $p' = p'(\Delta)$ projective planes and has $k' = k'(\Delta)$ boundary components $\beta'_1, \dots, \beta'_{k'}$. Assume further that $\beta'_1, \dots, \beta'_{d'}$ contain partisan boundary intervals of Γ , whilst $\beta'_{d'+1}, \dots, \beta'_{k'}$ do not. Fix as base point in β'_j an end point of an arc of Γ (or a point of a boundary partition of a component of $\partial\Sigma$, if β'_j does not meet an arc of Γ). Starting at the base point and reading prime labels of boundary corners and labels of vertex corners as they occur in order around β'_j gives the *prime label*

$$v_j = v_j(\Delta) = \prod_{i=1}^{n'_j} (h_{j_i}, f_{j_i}) \in V^\Lambda$$

of β'_j . Reading labels of both boundary corners and vertex labels we obtain the label of β'_j which, by construction, is equal to $\hat{\alpha}(v_j)$. If we identify $(v, 1) \in V^\Lambda$ with $v \in V$ then, for $j = d' + 1, \dots, k'$, the prime label is the same as the label of β'_j , since β'_j contains no partisan boundary intervals for such j . Thus, for $j = d' + 1, \dots, k'$, the label (and prime label) of β'_j is a fixed element v_j of V . The prime labels of boundary components of Δ are called the *prime boundary labels* of Δ .

Setting

$$\mathbf{v}(\Delta) = v_1, \dots, v_{k'}$$

and $h'(\Delta) = \sum_{j=1}^{k'} |v_j|_\Lambda$ we have $\mathbf{v}(\Delta) \in (V^\Lambda, \mathcal{L}, (k'), (h'))$ and, using property P3 of pictures, Δ determines a solution to $Q(\mathbf{v}(\Delta), \mathcal{L}, (h'), (k'))$ in V .

We have

$$\Gamma = \Gamma_1 \dot{\cup} \dots \dot{\cup} \Gamma_u,$$

where Γ_i is a picture on a surface Σ_i of type (n_i, t_i, p_i) . Let Γ_0 be the connected component of Γ containing Δ . Then Γ_0 is a component of a sub-picture Γ_i of Γ , for some i with $1 \leq i \leq u$. Thus Γ_0 is a picture on a connected surface Σ_0 of type

(n_0, t_0, p_0) , where $n_0 \leq n_i$, $t_0 \leq t_i$ and $p_0 \leq p_i$. As Δ is a region of a picture on Σ_0 we have

$$(14.2) \quad 2t' + p' \leq 2t_0 + p_0 \leq 2t_i + p_i.$$

As Γ is minimal so is Γ_0 and so, from Proposition 6.11, we have $\chi(\Delta) \geq \chi(\Sigma_0)$. That is

$$(14.3) \quad \begin{aligned} 2 - 2t' - p' - k' &\geq 2 - 2t_0 - p_0 - n_0 \geq 2 - 2t_i - p_i - n_i \\ \text{so } k' &\leq n_i + 2(t_i - t') + p_i - p'. \end{aligned}$$

Let Γ have a total of n' partisan regions $\Delta_1, \dots, \Delta_{n'}$. Assume that Δ_j is a V_j -region, where $V_j = A$ or B , and that Δ_j is the sum of $t'(\Delta_j)$ torii with $p'(\Delta_j)$ projective planes and has $k'(\Delta_j)$ boundary components of which $d'(\Delta_j)$ are partisan whilst the others are non-partisan. Let l_i be the number of partisan regions of Γ_i , let

$$\omega_0 = 0 \text{ and let } \omega_i = \sum_{j=1}^i l_j, \text{ for } i = 1, \dots, u,$$

so $\omega_u = n'$. We may assume that Δ_j is a region of Γ_i for all j such that $\omega_{i-1} + 1 \leq j \leq \omega_i$. Set

$$\begin{aligned} \mathbf{v}_j &= \mathbf{v}(\Delta_j), \text{ for } j = 1, \dots, n', \\ \mathbf{v} &= \mathbf{v}_1, \dots, \mathbf{v}_{n'}, \\ \mathbf{h}' &= h'(\Delta_1), \dots, h'(\Delta_{n'}), \\ \mathbf{k}' &= k'(\Delta_1), \dots, k'(\Delta_{n'}), \\ \mathbf{t}' &= t'(\Delta_1), \dots, t'(\Delta_{n'}) \text{ and} \\ \mathbf{p}' &= p'(\Delta_1), \dots, p'(\Delta_{n'}). \end{aligned}$$

Then

$$(14.4) \quad \mathbf{v} \in (A^\Lambda \dot{\cup} B^\Lambda, \mathcal{L}, \mathbf{h}', \mathbf{k}').$$

Also $0 \leq l_i \leq h_i$, as the boundary of every partisan region contains at least one proper exponential H -letter. Moreover

$$\sum_{j=\omega_{i-1}+1}^{\omega_i} h'(D_j) \leq h_i$$

and (14.2) and (14.3) hold, with $k'(D_j), t'(D_j), p'(D_j)$ in place of k', t', p' , for $\omega_{i-1} + 1 \leq j \leq \omega_i$ and $i = 1, \dots, u$. Hence

$$(14.5) \quad (\mathbf{h}', \mathbf{k}', \mathbf{t}', \mathbf{p}') \in \mathcal{N}(\mathbf{h}_1, \mathbf{k}_1, \mathbf{t}_1, \mathbf{p}_1) \subseteq \mathcal{N}.$$

The equation associated to $\mathbf{v}$ is $Q_\Delta = Q(\mathbf{v}, \mathcal{L}, \mathbf{h}', \mathbf{k}')$, an equation with environment $(X_{i_A}, X_{i_B}; A, B; A, B)^\Lambda$. Set $\phi_\Delta = \phi|_{L(\mathbf{v})}$. Then, since $\alpha_0(\lambda) = \alpha(\lambda)$ for all λ occurring in Q , the pair (ϕ_Δ, α_0) is a solution to $Q_\Delta = Q(\mathbf{v}, \mathcal{L}, \mathbf{h}', \mathbf{k}')$. Let $\mathbf{q}_\Delta$ be the system of quadratic words in standard form and β_Δ the map from $L_D(\mathbf{q}_\Delta)$ to $A^\Lambda \dot{\cup} B^\Lambda$ such that $Q_\Delta = (\mathbf{q}_\Delta = 1, \beta_\Delta, \mathcal{L})$.

Let $(h_1, f_1), \dots, (h_{W_1}, f_{W_1})$ be the letters of $\mathbf{z}_1$, where $W_1 = W_1(\mathbf{z}_1)$. Form a system of parameters $\mathcal{L}^-(\Gamma_s, P)$ from $\mathcal{L}(\Gamma_s, P)$ by deleting those equations $f_q = \alpha_s(\lambda_q)$ from $\mathcal{L}(\Gamma_s, P)$, for all q such that (h_q, f_q) is a proper exponential H -letter

with $h_q \in A \dot{\cup} B$. Now, since elements of Λ occurring in $\mathcal{L}(\Gamma_s, P)$ do not occur in Q_1 and we may choose then so that they do not occur in Q , we have

$$\mathbf{z}'' = \mathbf{v}, \mathbf{z}_2 \in (A^\Lambda \dot{\cup} B^\Lambda \dot{\cup} \bigcup_{i \in I} H_i^\Lambda, \mathcal{L} \cup \mathcal{L}^-(\Gamma_s, P), \mathbf{h}'', \mathbf{n}''),$$

where $\mathbf{h}'' = \mathbf{h}', \mathbf{h}_2$, and $\mathbf{n}'' = \mathbf{n}', \mathbf{n}_2$. Define $\mathbf{t}'' = \mathbf{t}', \mathbf{t}_2$ and $\mathbf{p}'' = \mathbf{p}', \mathbf{p}_2$ and

$$\begin{aligned} Q'' &= Q(\mathbf{z}'', \mathcal{L} \cup \mathcal{L}^-(\Gamma_s, P), \mathbf{n}'', \mathbf{t}'', \mathbf{p}'') \\ &= (\mathbf{q}'' = 1, \beta'', \mathcal{L} \cup \mathcal{L}^-(\Gamma_s, P)), \end{aligned}$$

a system of equations with environment $((X_j)_{j \in J}; (H_j)_{j \in J}; (H_j)_{j \in J})^\Lambda$. Define

$$\phi_0 : F(L(\mathbf{q}'')) \longrightarrow A * B * *_{i \in I} H_i$$

by $\phi_0|_{L(\mathbf{q}_\Delta)} = \phi_\Delta$ and $\phi_0|_{L(\mathbf{q}_2)} = \phi|_{L(\mathbf{q}_2)} = \phi_2$. Then (ϕ_0, α_0) is a solution to Q'' and $(\mathbf{h}'', \mathbf{n}'', \mathbf{t}'', \mathbf{p}'') \in \mathcal{N}$.

Now Γ_s is a picture over G on a surface of type $(\mathbf{n}_1, \mathbf{t}_1, \mathbf{p}_1)$ with partisan regions $\Delta_1, \dots, \Delta_{n'}$. Removing boundary labels from partisan boundary intervals of Δ_j we replace these labels with the prime labels of the corresponding boundary intervals of Γ_s . We call the result a picture with *partisan boundary labels stripped* and denote it Γ_s^- . Note that given $(\mathbf{h}, \mathbf{n}, \mathbf{t}, \mathbf{p})$, $\mathbf{z}$ and $\mathcal{L}$ the graph $G(\mathbf{z}_1)$ is uniquely determined. Given Γ_s the vertex $v(\Gamma_s)$ of $G(\mathbf{z}_1)$ is unique and, since {active left markings of Γ_s } = {active left markings of Γ_s^- } we may determine $v(\Gamma_s)$ from Γ_s^- . We may therefore refer to this vertex as $v(\Gamma_s^-)$. Furthermore, from the above and Theorem 7.19, if there exists a solution (ϕ, α) to $Q(\mathbf{z}, \mathcal{L}, \mathbf{h}, \mathbf{n})$ then there exists a Z -path in $G(\mathbf{z}_1)$ with path-subgraph P and with initial vertex $v(\Gamma_s^-)$ and a solution α_0 to $\mathcal{L} \cup \mathcal{L}^-(\Gamma_s, P)$. Again $\mathcal{L}^-(\Gamma_s, P)$ may be constructed using only Γ_s^- , since the equation of $\mathcal{L}(\Gamma_s, P)$ corresponding to a partisan boundary interval with prime label (h_q, f_q) , the q th letter of $\mathbf{z}_1$, on partisan boundary component of Γ has the form $f_q = \alpha_s(\lambda_q)$, and so has been deleted. Hence we may refer to $\mathcal{L}^-(\Gamma_s, P)$ as $\mathcal{L}^-(\Gamma_s^-, P)$. Also, given $\mathbf{z}$, $\mathcal{L}$, Γ_s^- and $\mathcal{L}^-(\Gamma_s^-, P)$ we may construct the system Q'' , which is again uniquely determined by these data. The existence of the solution (ϕ, α) then guarantees the existence of the solution (ϕ_0, α_0) to Q'' . Hence given a solution (ϕ, α) to Q we obtain the following.

- (a) A picture Γ_s^- with partisan boundary labels stripped which is obtained (by stripping partisan boundary labels) from a Z -reduced, reduced, minimalistic, picture Γ_s such that (α_s, Γ_s) corresponds to a solution to $Q_{\mathcal{H}_1}$ for some retraction $\alpha_s : M \longrightarrow \mathbb{Z}$.
- (b) The path-subgraph P of some Z -path in $G(\mathbf{z}_1)$ with initial vertex $v(\Gamma_s^-)$ and a corresponding system of parameters $\mathcal{L}^-(\Gamma_s^-, P)$.
- (c) A system of equations Q'' as above and solution (ϕ_0, α_0) to Q'' .

Conversely, given the data of (a), (b) and (c) we may form a picture Γ_t from Γ_s^- as follows. Let $(h_1, f_1), \dots, (h_{W_1}, f_{W_1})$ be the letters of $\mathbf{z}_1$ and let b_q be the boundary interval of Γ_s^- with prime label (h_q, λ_q) . Form Γ_t by assigning to the partisan boundary interval b_q of Γ_s^- the label $\hat{\alpha}_t(h, f)$. Define a retraction $\alpha_t : M \longrightarrow \mathbb{Z}$ by $\alpha_t(\lambda_q) = \alpha_0(f_q)$, if b_q is a partisan boundary interval and $\alpha_t(\lambda_q) = |b_q|$ otherwise. Then α_t is a solution to $\mathcal{H}_1$ and the boundary interval b_q of Γ_t has label $\hat{\alpha}_t(h_q, \lambda_q)$, for $q = 1, \dots, W_1 = W_1(\mathbf{z}_1)$. Hence (α_t, Γ_t) corresponds to a solution (ϕ_t, α_t) to $Q_{\mathcal{H}_1}$. Note that $v(\Gamma_t) = v(\Gamma_s^-) = v(\Gamma_s)$ in $G(\mathbf{z}_1)$ and that $\mathcal{L}^-(\Gamma_s, P) = \mathcal{L}^-(\Gamma_t, P)$. Therefore α_0 is a solution to $\mathcal{L}^-(\Gamma_t, P)$ and so also to $\mathcal{L}(\Gamma_t, P)$. It follows from Theorem 7.19 that there exists a solution α_c to $\mathcal{L}$ and a picture Γ_c obtained from Γ_t

by Z -insertion along a path with path-subgraph P such that (α_c, Γ_c) corresponds to a solution (ϕ_c, α_c) to Q_1 . In fact, since α_0 is a solution to $\mathcal{L} \cup \mathcal{L}(\Gamma_t, P)$ and $\alpha_0(\lambda) = \alpha_c(\lambda)$, for all λ occurring in $\mathbf{z}_1$ or $\mathcal{L}$ we may take $\alpha_c = \alpha_0$. From (α_0, Γ_0) we obtain a solution (ϕ_c, α_0) to Q_1 . If $\phi_2 = \phi_0|_{L(\mathbf{q}_2)}$ then (ϕ_2, α_0) is a solution to Q_2 . Using (14.1) we obtain a solution (ϕ, α_0) to Q .

It therefore suffices to determine whether the data of (a), (b) and (c) exist and if so to find it. We describe an algorithm, similar to those described in [7], [8] and [9], which solves this problem. First note that given $\mathbf{z}$ (which we are assuming is special) we may construct $\mathbf{z}_1$ and calculate $\mathcal{B} = \mathcal{B}(\mathbf{z}_1)$ of Definition 13.6, since the word problem is solvable in A and B . The number of arcs of Γ_s , if it exists, is at most $\mathcal{B}$, using Theorem 13.7. The same is true of Γ_s^- . Assume that Γ_s exists and that the set of arcs of Γ_s^- is $\mathcal{A}$, so $0 \leq |\mathcal{A}| \leq \mathcal{B}$. There are, up to homeomorphism, finitely many surfaces of type $(\mathbf{n}_1, \mathbf{t}_1, \mathbf{p}_1)$ which have no closed components. We fix one such surface, call it Σ and assume that Γ_s is a picture on Σ . Then Σ is a disjoint union of surfaces $\Sigma_1, \dots, \Sigma_u$, where Σ_j is of type (n_j, t_j, p_j) , for $j = 1, \dots, u$. Note that Σ_j need not be connected and each connected component of Σ is a connected component of Σ_j , for some j with $1 \leq j \leq u$. Assume then that Σ has c connected components $\Omega_1, \dots, \Omega_c$, for some c such that $u \leq c \leq \mu_{u+1}$. If Ω is a connected component of Σ we denote by $\Gamma_s^-(\Omega)$ the restriction of Γ_s^- to Ω . There are finitely many partitions $(A_1, \dots, A_c)$ of $|\mathcal{A}|$ and (if Γ_s^- exists) one such partition satisfies $A_j =$ the number of arcs of $\Gamma_s^-(\Omega_j)$, $j = 1, \dots, c$. Fix some j , with $1 \leq j \leq c$ and write $\Omega = \Omega_j$. Let Ω be the connected sum of t_Ω tori and p_Ω projective planes and have n_Ω boundary components with prime boundary labels z_i , for $i = d+1, \dots, d+n_\Omega$ and an appropriate integer d .

Let Ω_0 be the surface

$$\Omega_0 = \Omega \setminus \{\text{int}(v) : v \text{ is a vertex of } \Gamma_s^-(\Omega)\}.$$

Let V_j be the number of vertices of $\Gamma_s^-(\Omega)$. As $\Gamma_s^-(\Omega)$ has A_j arcs and its vertices all have index ml it follows that $V_j \leq A_j/ml$.

Let Ω_{00} be the surface formed by cutting Ω_0 along arcs of $\Gamma_s^-(\Omega)$. Each connected component of Ω_{00} is the closure of a region of $\Gamma_s^-(\Omega)$. Since Γ_s is minimalistic so is $\Gamma_s^-(\Omega)$, so for each region Δ of $\Gamma_s^-(\Omega)$ we have $\chi(\Delta) \geq \chi(\Omega)$. As Ω_0 is connected Ω_{00} has at most $A_j + 1$ connected components and, by construction,

$$\begin{aligned} \chi(\Omega_{00}) &= \chi(\Omega) - V_j + A_j \\ &= 2 - t_\Omega - p_\Omega/2 - n_\Omega - V_j + A_j \\ &\geq 2 - (t_\Omega + p_\Omega/2) - n_\Omega + A_j(1 - 1/ml). \end{aligned}$$

Set $\chi_j = 2 - (t_\Omega + p_\Omega/2) - n_\Omega - A_j(1 - 1/ml)$. There are finitely many surfaces of Euler characteristic χ_j with at most $A_j + 1$ connected components each of Euler characteristic at least $\chi(\Omega)$. The algorithm takes each such surface and attempts to construct $\Gamma_s^-(\Omega)$ from it. Given a such a surface which we take to be Ω_{00} suppose that $\partial\Omega_{00}$ has connected components $c_1, \dots, c_D$. There are finitely many ways of assigning a positive integer ρ_i to c_i , for $i = 1, \dots, D$, so that $\sum_{i=1}^D \rho_i = 2(A_j + 1)$. Choose one such assignment and divide c_i into ρ_i intervals, $i = 1, \dots, \rho_i$. There are now finitely many ways of identifying intervals in pairs to obtain a surface of type $(n_\Omega + V_0, t_\Omega, p_\Omega)$, for each choice of V_0 such that $0 \leq V_0 \leq A_j/ml$. If Γ_s^- exists then one such gives rise to a surface homeomorphic to Ω_0 . Fix such a choice of V_0 and of pairings and assume the resulting surface is Ω_0 . The paired intervals become

properly embedded arcs on Ω_0 which we refer to as arcs, as they become arcs of Γ_s^- if the process is successful. To obtain Ω all but n_Ω of the boundary components of Ω_0 are capped off with disks, which we refer to as vertices (they'll become the vertices of $\Gamma_s^-(\Omega)$). This must be done in such a way that each vertex meets exactly ml arcs. There are finitely many ways of doing this, one of which results in a surface homeomorphic to Ω . Fix one such choice and assume the resulting surface is Ω and that the vertices and arcs are those of $\Gamma_s^-(\Omega)$. Images of connected components of Ω_{00} in Ω are called regions of Ω . Choose one of the finitely many possible orientations of $\Gamma_s^-(\Omega)$ consistent with the definition of orientation of a sketch in Section 6. The union of Ω with these arcs, vertices, regions and orientation is called Γ_b .

To recover $\Gamma_s^-(\Omega)$ the components of $\{\text{vertices of } \Gamma_b\} \setminus \{\text{arcs of } \Gamma_b\}$ and of $\{\text{boundary components of } \Omega\} \setminus \{\text{arcs of } \Gamma_b\}$ must be labelled to give a picture with partisan boundary labels stripped. There are finitely many ways of labelling corners of vertices with elements of $A \dot{\cup} B$ so that each vertex has label $r^{\pm m}$. There are finitely many ways of assigning the z_j 's to the boundary components of Ω . This must be done in such a way that if $z_{d+j} \in A^\Lambda \dot{\cup} B^\Lambda$ then z_{d+j} is assigned to a boundary component which meets no arc of Γ_b . Suppose z is assigned to β . There are then finitely many ways of assigning a boundary partition to β which $\mathcal{H}_1$ admits z , since the arcs of Γ_b are already fixed. Suppose that a boundary partition of β , which $\mathcal{L}$ -admits z has been chosen. There are finitely many ways of assigning the letters of z as prime labels of the boundary intervals of this boundary partition. Assume then that β has prime label z with some boundary partition. If $[b, c]$ is a non-partisan boundary interval of this boundary partition then the label of $[b, c]$ and of all components of $[b, c] \setminus \Gamma_b$ is fixed by the assignments made so far. Choosing boundary partitions and prime labels in this way for all boundary components of $\partial\Omega$ we obtain labels for all corners of Γ_b except those containing partisan boundary intervals.

Since A and B have solvable word problem, the conditions $P1$, $P2$ and $P4$, for Γ_b to be a picture with partisan boundary labels stripped can now be checked, as prime labels are enough to verify these conditions. Furthermore $P3$ can be verified for every non-partisan region of Γ_b . To see this note that, since Γ_s is obtained from Γ by Z -cancellation and Γ_s is minimalistic, by construction $\chi(\Delta) \geq \chi(\Omega)$, for all regions Δ of Γ_b .

Suppose Δ is a region of Γ_b which has s' boundary components all of which are non-partisan, with labels $(u_1, \dots, u_{s'})$ in V , for $V = A$ or B . Suppose that Δ is a union of t' torii and p' projective planes. Since $\chi(\Delta) \geq \chi(\Omega)$, (14.2) and (14.3) hold, with j in place of i . Setting

$$\mathbf{u}' = u_1, \dots, u_{s'}$$

we have $\mathbf{u} \in (V^\Lambda, \mathcal{H}_1, s', t', p')$. Furthermore $(0, s', t', p') \in \mathcal{N}$ and so, by hypothesis, the $\mathcal{GE}(0, s', t', p')$ -problem is solvable in (A, B) . Therefore we may check whether or not $(s', t', p') \in \mathcal{H}_1\text{-genus}(\mathbf{u}')$ in V : that is whether condition $P3$ holds for Δ . If all these conditions are satisfied then Γ_b is a candidate for $\Gamma_s^-(\Omega)$. We now repeat this process for all connected components $\Omega = \Omega_1, \dots, \Omega_c$ of our candidate Σ . If successful, this gives rise to a candidate for Γ_s^- .

Assume we have a candidate for Γ_s^- , which we continue to call Γ_b . Next we consider the partisan regions of Γ_b . Using the same notation for partisan regions of Γ_b as used for Γ_s above, the prime labels boundary components of partisan

boundary regions of Γ_b give rise to an element

$$\mathbf{v} \in (A^\Lambda \cup B^\Lambda, \mathcal{L}, \mathbf{h}', \mathbf{k}')$$

as in (14.4) and furthermore (14.5) holds, as before.

Since the word problem is solvable in A and B the graph $G(\mathbf{z}_1)$ may be constructed and the vertex $v(\Gamma_s^-)$ located, using Γ_s^- . There are finitely many path-subgraphs of Z -paths in $G(\mathbf{z}_1)$, since $G(\mathbf{z}_1)$ is a finite graph, and we fix one such and call it P . This allows construction of the system of parameters $\mathcal{L}^-(\Gamma_s^-, P)$. We may now construct the system of equations $Q'' = Q(\mathbf{z}'', \mathcal{L} \cup \mathcal{L}^-(\Gamma_s^-, P), \mathbf{h}'', \mathbf{n}'')$ described above. As $(\mathbf{h}'', \mathbf{n}'', \mathbf{t}'', \mathbf{p}'') \in \mathcal{N}$ and Q'' has environment $((X_j)_{j \in J}; (H_j)_{j \in J}; (H_j)_{j \in J})^\Lambda$, there is an algorithm to determine whether or not Q'' has a solution and if so to find one.

If a solution (ϕ_0, α_0) to Q'' is found then, as above, a solution to Q may be constructed. If no such solution exists then either there is no solution to Q or one of the choices above was not the right one. At each stage there are finitely many choices so once they've all been tried either we have a solution or there is none. This completes the description of the algorithm.

REFERENCES

- [1] G. Baumslag. Some aspects of groups with unique roots. *Acta Math.*, 104:217–303, 1960.
- [2] G. Baumslag. On free Q-group. *Commun. on Pure and Appl. Math.*, 18:25–30, 1965.
- [3] W.A. Bogley and S.J. Pride. Aspherical relative presentations. *Proc. Edinburgh Math. Soc.*, 35:1–39, 1992.
- [4] Leo P. Comerford and Charles C. Edmunds. Quadratic equations over free groups and free products. *J. Algebra*, pages 276–297, 1981.
- [5] Leo P. Comerford and Charles C. Edmunds. Quadratic parametric equations over free groups. *Cont. Math.*, 33:159–196, 1984.
- [6] M. Culler. Using surfaces to solve equations in free groups. *Topology*, pages 237–300, 1981.
- [7] Andrew J. Duncan and J. Howie. The genus problem for one-relator products of locally indicable groups. *Math. Z.*, 208:225–237, 1991.
- [8] Andrew J. Duncan and J. Howie. The nonorientable genus problem for one-relator products. *Commun. in Alg.*, 19:2547–2556, 1991.
- [9] Andrew J. Duncan and J. Howie. One relator products with high-powered relators. In Graham A. Niblo and Martin A. Roller, editors, *Geometric Group Theory, Volume 1 of Proceedings of the Symposium held in Sussex 1991*, volume 181 of *London Mathematical Society Lecture Note Series*, pages 48–74. Cambridge University Press, 1994.
- [10] M. Edjvet and R.M. Thomas. The groups $(l, m|n, k)$. *J. Pure Applied Algebra*, 114:175–208, 1997.
- [11] R.A. Fenn. *Techniques of Geometric Topology*. Number 57 in London Math. Soc. Lecture Notes. Cambridge University Press, Cambridge, 1983.
- [12] A. Myasnikov G. Baumslag and V. Remeslennikov. Algebraic geometry over groups I. Algebraic sets and ideal theory. *J. of Algebra.*, 219:16–79, 1999.
- [13] S.M. Gertsten. Reducible diagrams and equations over groups. In S.M. Gersten, editor, *Essays in group theory*, volume 8 of *MSRI Publications*, pages 15–73. Springer, 1987.
- [14] R.Z. Goldstein and E.C. Turner. Applications of topological graph theory to group theory. *Math. Z.*, 165:1–10, 1979.
- [15] R. I. Grigorchuk and I. G. Lysionok. A description of solutions of quadratic equations in hyperbolic groups. *Internat. J. Algebra Comput.*, 2(3):237–274, 1992.
- [16] J Howie. The quotient of a free product of groups by a single high-powered relator. I. Pictures. Fifth and higher powers. *Proc. London Math. Soc. (3)*, 59:507–540, 1989.
- [17] J Howie. The quotient of a free product of groups by a single high-powered relator. II. Fourth powers. *Proc. London Math. Soc. (3)*, 61:33–62, 1989.
- [18] J Howie. The quotient of a free product of groups by a single high-powered relator. III. the word problem. *Proc. London Math. Soc. (3)*, 62:590–606, 1991.

- [19] J. Howie and R.M. Thomas. The groups $(2, 3, p, q)$; asphericity and a conjecture of Coxeter. *J. Algebra*, 154:289–309, 1993.
- [20] O. Kharlampovich and A. Myasnikov. Equations in a free Q -group. *Trans. Amer. Math. Soc.*, 350(3):947–974, 1998.
- [21] O. Kharlampovich and A. Myasnikov. Irreducible affine varieties over a free group. I: Irreducibility of quadratic equations and nullstellensatz. *J. of Algebra*, 200(2):472–516, 1998.
- [22] O. Kharlampovich and A. Myasnikov. Irreducible affine varieties over a free group. II: Systems in triangular quasi-quadratic form and description of residually free groups. *J. of Algebra*, 200(2):517–570, 1998.
- [23] Seymour Lipschutz and Charles F. Miller III. Groups with certain solvable and unsolvable decision problems. *Comm. Pure Appl. Math.*, XXIV:7–15, 1971.
- [24] R.C. Lyndon. Equations in free groups. *Trans. Amer. Math. Soc.*, 96:445–457, 1960.
- [25] R.C. Lyndon. Groups with parametric exponents. *Trans. Amer. Math. Soc.*, 96:518–553, 1960.
- [26] R.C. Lyndon and M.P. Schützenberger. The equation $a^m = b^n c^p$ in a free group. *Michigan Math. J.*, 9:289–298, 1962.
- [27] G.S. Makanin. Equations in a free group. (russian). *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(6):1199–1273, 1982. English translation in: *Math. USSR-Izv.* 21 (1983), no. 3, 546–582.
- [28] I.U. Matiyasevič. *Hilbert's tenth problem*. Foundations of Computing. MIT Press, Cambridge, Mass., 1993.
- [29] A.G. Myasnikov and V.N. Remeslennikov. Exponential groups II: extensions of centralisers and tensor completion of CSA-groups. *Int. J. of Algebra and Comp.*, 6(6):687–711, 1996.
- [30] S.J. Pride. Identities among relations of group presentations. In (E. Ghys, A. Haefliger, and A. Verjovsky, editors, *Group Theory from a Geometric Viewpoint, Conference Proceedings, ICTP*), pages 687–716. World Scientific, 1991.
- [31] E. Rips and Z. Sela. Canonical representatives and equations in hyperbolic groups. *Invent. Math.*, 120(3):489–512, 1995.
- [32] C.P. Rourke. Presentations and the trivial group. In R. Fenn, editor, *Topology of Low Dimensional Manifolds*, number 722 in *Lecture Notes in Mathematics*, pages 134–143. Springer, 1979.
- [33] Zlil Sela. Diophantine geometry over groups. II. Completions, closures and formal solutions. to appear.
- [34] Zlil Sela. Diophantine geometry over groups. I. Makanin–Razborov diagrams. *Publ. Math. Inst. Hautes Études Sci.*, 93:31–105, 2001.
- [35] Hamish Short. *Topological methods in group theory: the adjunction problem*. PhD thesis, University of Warwick, 1984.

SCHOOL OF MATHEMATICS AND STATISTICS, MERZ COURT, UNIVERSITY OF NEWCASTLE UPON TYNE, NEWCASTLE UPON TYNE, NE1 7RU, UK

E-mail address: `a.duncan@ncl.ac.uk`

URL: `http://www.ncl.ac.uk/math/`